Wireless Radiation and Electromagnetic Field Studies Published since 2024 (Volume 3)

Joel M. Moskowitz, Ph.D.
School of Public Health
University of California, Berkeley

Electromagnetic Radiation Safety

November 18, 2025 Edition

Links hereby incorporated by reference.

Non-thermal biological effects of radiofrequency electromagnetic radiation: Mechanistic insights into male reproductive vulnerability in the era of ubiquitous exposure

Jangid P, Rai U, Ahmed S, Singh S, Singh R. Non-thermal biological effects of radiofrequency electromagnetic radiation: Mechanistic insights into male reproductive vulnerability in the era of ubiquitous exposure. Reproductive Toxicology, Volume 138, 2025. doi:10.1016/j.reprotox.2025.109087.

Abstract

The rapid proliferation of wireless technologies has led to continuous, low-intensity exposure to radiofrequency electromagnetic radiation (RF-EMR) from devices such as mobile phones, Wi-Fi routers, and wearable electronics. Although RF-EMR is non-ionizing, a growing body of evidence indicates that chronic exposure can induce non-thermal biological effects with significant implications for male reproductive health. This review synthesizes current findings on the mechanistic pathways through which RF-EMR may impair fertility, focusing on oxidative stress, mitochondrial dysfunction, DNA damage, apoptosis, autophagy dysregulation, and hormonal disruption via hypothalamic-pituitary-gonadal (HPG) axis interference. Experimental and clinical studies consistently link RF-EMR exposure to reduced sperm count and motility, altered testicular architecture, blood-testis barrier compromise, and suppressed testosterone synthesis, often at specific absorption rates below current safety thresholds. These outcomes are underpinned by redox imbalance, mitochondrial collapse, and steroidogenic impairment, with emerging evidence for epigenetic and transgenerational effects. Existing regulatory frameworks, based largely on thermal safety limits, fail to address the complexity of real-world, multi-source, long-term exposures. The review highlights critical knowledge gaps, particularly regarding 5 G and millimeter-wave frequencies, and underscores the need for standardized research protocols, biologically relevant exposure metrics, and precautionary public health measures to mitigate reproductive risks in an increasingly wireless environment.

Highlights

- RF-EMR triggers oxidative stress in male reproductive cells.

- Leydig cell mitochondria are highly sensitive to RF exposure.
- RF-EMR impairs testosterone synthesis and steroidogenesis.
- Sperm quality declines after non-thermal RF-EMR exposure.
- ROS and autophagy mediate RF-EMR-induced cell damage.

Limitations of SAR-based regulatory paradigms

Regulatory bodies like the International Commission on Non-Ionizing Radiation Protection (ICNIRP) and the Federal Communications Commission (FCC) still rely on SAR as the main safety benchmark for RF-EMR exposure [14], [74]. But SAR mainly reflects how much tissue heats up over short periods, it was originally designed to prevent thermal damage in lab settings [15]. While that works for avoiding immediate overheating, it doesn't capture the kinds of long-term, non-thermal effects that research is now increasingly reporting. For example, oxidative stress, hormonal imbalances, mitochondrial dysfunction, and even DNA fragmentation have been observed at SAR levels still considered "safe" under current guidelines [4], [19]. These kinds of findings make it clear that we need to rethink how exposure is measured, especially for reproductive health, and move toward including more biologically relevant markers like oxidative stress indicators, endocrine changes, and genotoxic outcomes.

The need for a precautionary framework

A proactive, precautionary approach is essential to safeguard public health amid ongoing scientific uncertainty. The absence of acute thermal injury must not be equated with biological safety. Multiple expert groups have advocated for the inclusion of non-thermal biological endpoints, such as oxidative stress, hormonal disruption, and reproductive toxicity, into current safety standards [5], [17].

To align public health policy with emerging evidence, reforms should prioritize:

- Mandatory RF-EMR labeling on consumer electronics;
- Promotion of low-exposure practices (e.g., using speaker mode or wired headphones, maintaining device distance, limiting phone use in low-signal areas, etc.);
- Creation of low-EMF zones in schools, hospitals, and transport systems;
- Dedicated funding for longitudinal studies on fertility and epigenetic impacts;
- Regular RF exposure mapping and transparent public communication.

Incorporating these elements into policy will better reflect biological realities and ensure protection for high-risk populations. Without such action, regulatory frameworks may continue to underestimate the long-term reproductive risks of chronic low-intensity RF-EMR exposure.

Conclusion

With wireless technologies now embedded in almost every part of daily life, we're constantly exposed to low levels of RF-EMR, often without even realizing it. This kind of ongoing exposure has raised real concerns about biological effects that go well beyond just heating tissues. Studies using cell lines, animals, and even human subjects have pointed to consistent changes in things like redox balance, mitochondrial health, programmed cell death pathways, autophagy, and hormonal regulation, and these effects seem especially pronounced in the male reproductive system. These disturbances have been

linked to reduced sperm quality, disrupted testicular architecture, and compromised fertility, with some studies pointing to potential impacts on future generations.

Existing safety standards, primarily based on thermal effects, do not capture the complexity of these non-thermal, long-term exposures. As technologies like 5 G introduce novel frequencies and modulations, there is a critical need for biologically relevant models, standardized endpoints, and human-focused cohort studies. A multidisciplinary approach integrating mechanistic research and policy updates is essential to ensure reproductive safety amid expanding RF-EMR exposure.

https://www.sciencedirect.com/science/article/pii/S0890623825002588

--

Radiofrequency radiation-induced gene expression

Lai H, Levitt BB. Radiofrequency radiation-induced gene expression. Rev Environ Health. 2025 Oct 29. doi: 10.1515/reveh-2025-0104.

Abstract

Genes are differentially expressed in cells in response to changes in the internal or external environment. The response is generally an adaptive mechanism to the environmental challenge to restore cell functions to homeostasis. There are numerous studies reporting changes in gene expression in cells exposed to radiofrequency radiation (RFR), the type of energy emitted by wireless communication devices. The major genes affected are those involved in: repair of damaged proteins, response to stress, oxidative changes, apoptosis, DNA damage detection and repair, and changes in neural functions. Gene expression data supports the notion that RFR is a stressor that causes oxidative changes and DNA and protein damage in cells under different exposure conditions, in many biological systems. Changes in all these significant gene expression effects are supported by results of other biological studies of RFR exposure in the literature. They should be considered in the setting of RFR-exposure guidelines.

Excerpts

"Gene expressions are under feedback control in response to external challenges with numerous variables. RFR has a complex interaction with living organisms depending on various parameters of the field. With these considerations in mind, response oscillation and non-linear dose-response could occur. But gene expression is not expected under all exposure conditions and biological systems studied. Thus, inevitably, in some studies, no significant effects would be expected. There are studies that reported no significant effect of RFR on gene expression related to protein damage and removal, DNA, oxidative processes, stress, and brain functions (Table 7)."

"Forty gene expression effects (marked as low intensity (LI) in Tables 1–6) were observed at a SAR less than 0.4 W/kg. This is relevant to the effects of exposure to RFR in the environment. 0.4 W/kg is the dose rate assumed to be safe by international radiofrequency radiation guidelines (e.g., the International Commission on Non-Ionizing Radiation Protection (ICNIRP) and the International Electronics and Electrical Engineers (IEEE)). This is important to understand regarding the validity of those organization's high-intensity standards-setting models. (For readers who are interested in 'low-

intensity' RFR effects, there are more than 200 studies that reported biological effects at SAR <0.4 W/kg [see the 'RFR-low-intensity studies' file in reference [1]]).

Lastly, one may wonder whether up-regulation of gene expression would lead to translation into functional proteins. Several RFR studies support increased translation (e.g., [33], [34], [35], [36]). It is an important complement to the gene-expression studies to look at translation levels of proteins as these are the functional units that promote/sustain both health and illness. Proteomic studies are an attractive next step in understanding the biological effects of RFR exposure. This was suggested more than a decade ago [37]. Changes in transcription and protein functions point to a need to investigate molecular signaling pathways after RFR exposure [38]. In addition, the data on gene-expression suggests two areas of mechanism research that are worth pursuing. The first area is the involvement of stress in RFR-induced biological effects. Involvements of 'cellular stress responses' and the hypothalamic-pituitary-adrenal axis and, particularly, effects on the limbic system should be further investigated [3]. The second area is the cellular oxidative processes. Particularly, the mechanism of induction of free radicals by RFR should be systematically studied. Present data indicates that low-frequency modulations in RFR may play a role [39]. It is, however, not known how much these low-frequency components contribute to the overall effect. Similar effects on gene expression have been reported after exposure to extremely-low frequency electromagnetic fields (see 'static and ELF EMF genetic effects' file in reference [1])."

Open access: https://www.degruyterbrill.com/document/doi/10.1515/reveh-2025-0104/html

--

Electromagnetic fields and oxidative stress: The link to the development of cancer, neurological diseases, and behavioral disorders

Leković Ž. Electromagnetic fields and oxidative stress: The link to the development of cancer, neurological diseases, and behavioral disorders. Electromagn Biol Med. 2025 Oct 21:1-25. doi: 10.1080/15368378.2025.2567872.

Abstract

Background: Epidemiological studies suggest an association between exposure to electromagnetic fields (EMFs) and an increased incidence of malignant, cardiovascular, and neurodegenerative diseases. This study aims to elucidate the fundamental principles and plausible mechanisms by which EMFs may influence physiological and pathological processes that lead to disease development.

Materials and methods: Published reports of oxidative stress, DNA damage, and disease risk related to EMF exposure were examined. The literature review provided the foundation for building a new conceptual model called the Electromagnetic Pathogenesis (EMP) model.

Mechanisms: The EMP model proposes an increase in the probability of electron tunneling through the mitochondrial electron transport chain as the primary pathophysiological mechanism triggered by non-ionizing EMFs. Induced electric fields and quantum tunneling may enhance electron leakage during mitochondrial respiration, which is a major source of free radicals.

Findings: There is a deep connection between quantum tunneling, entropy, and Heisenberg's principle. As a direct consequence of Heisenberg's principle, electron tunneling is essentially involved in free radical production and entropy generation in cells. Both normal aging and chronic diseases may be considered as the biologic manifestations of increasing entropy. Heisenberg's principle underlies normal aging and sets the limit to life expectancy.

Social implications: The human brain, particularly the structural and functional networks that support social communication, is highly vulnerable to oxidative stress associated with EMF exposure. Long-term exposure may negatively affect social and reproductive behaviors in both men and women, potentially contributing to a decline in fertility rates and the acceleration of population aging.

Concluding remarks

As a direct consequence of Heisenberg's principle, quantum tunneling is essential for energy conversion in mitochondria and many physiological processes in cells, including the production of free radicals.

The external time-varying magnetic field brings energy into the biological system and transfers it to charged particles, especially electrons, via induced electric field (electromotive force). During exposure, the induced electric field increases the probability of electron tunneling through the mitochondrial ETC. This facilitates electron leakage during mitochondrial respiration, which is the major source of free radicals in cells.

The oxidative stress associated with man-made EMF exposures represents one of the primary mechanisms that can lead to disease development. Under oxidative stress conditions, excess superoxide releases iron from ferritin and iron-sulfur proteins and increases intracellular free iron. In addition, peroxynitrite, formed when nitric oxide reacts with the superoxide, can liberate iron from iron-sulfur proteins and copper from ceruloplasmin, which destroys an important part of the antioxidant defense – transition metal binding proteins. The elevation in free iron and copper and the ability of DNA to bind metal ions leads to damage by the site-specific Fenton-type reaction, which generates hydroxyl radicals immediately near the site of metal fixation in the DNA.

Autoxidation of dopamine is relatively slow under normal physiological conditions but is accelerated by free transition metals like iron and copper. Oxidative stress-induced dysfunction of the dopamine system has been implicated in different nervous system diseases and neuropsychiatric behavioral disorders, including autism, addiction, and depression.

Embryonic and fetal brain tissues are especially susceptible to peroxidative injury. Maternal oxidative stress may compromise fetal development, leading to congenital malformations or the pathogenesis of neurodevelopmental disorders. EMFs as an environmental factor have the potential to increase the rate of somatic mutations and incidence of brain disorders, including autism.

The human brain, particularly the structural and functional networks that support social communication, is highly vulnerable to oxidative stress associated with EMF exposure. Disturbance in social communication affects partnership, marital status, reproductive behavior, and ultimately childbearing.

Oxidative stress can also initiate degenerative processes in other tissues and organs, which explains the association between social relationships, including marital status, and disease risk. At the population level, long-term EMF exposure can contribute to a decline in the total fertility rate and the acceleration of population aging.

In establishing a causal relationship between electromagnetic fields and associated diseases, the key and final step would be to reduce population exposure. The EMP model predicts that measures to reduce the level of electromagnetic fields in the environment based on the precautionary principle would lead to a decrease in the incidence of almost all diseases associated with oxidative stress, including malignant, cardiovascular, neurodevelopmental, and neurodegenerative diseases.

The EMP model indicates possible links between long-term EMF exposure, oxidative stress, brain disorders, and social and reproductive behavior, providing a solid foundation for future research.

https://pubmed.ncbi.nlm.nih.gov/41117390/

--

Transition Pathways Towards Electromagnetic Sustainability in the Built and Lived Environment

Habash R, Baho GY. Transition Pathways Towards Electromagnetic Sustainability in the Built and Lived Environment. *Sustainability*. 2025; 17(22):10252. doi: 10.3390/su172210252

Abstract

Electromagnetic (EM) fields, as one of the basic forms of energy in the built and lived environment (BLE), present an environmental health challenge, yet they often remain an overlooked concern, particularly with the development of information and communication technologies (ICT) and energy systems. Although these fields are essential for the contemporary infrastructure, society needs to engage in a thorough discussion regarding their potential impact on health. In light of this, a commitment should be made to design and manage technologies and infrastructure that strive to lower EM pollution, while ensuring optimal functionality. Achieving this goal requires viable urban planning and sustainability strategies. The motivation of this study is to examine various instances to foster a deeper understanding of the EM in the BLE. It explores significant sources of exposure and major safety guidelines. A literature review and EM field audits in three locations within two cities in Canada and the UK have been provided to understand the trends and serve as a comparative sample. Key transition pathways towards EM sustainability have been proposed, including the establishment of observatory systems in urban locations, hygiene practices, risk governance, and an interplay between sustainability and technology.

Open access: https://www.mdpi.com/2071-1050/17/22/10252

--

Brain Tumor and Mobile Phone Risk Among Young People: Analysis of Japanese People Using the MOBI-Kids International Case-Control Study

Kojimahara N, Kiyohara K, Sato Y, Wake K, Taki M, Castano-Vinyals G, Yamaguchi N, The Study Group of MOBI-Kids Japan. Brain Tumor and Mobile Phone Risk Among Young People: Analysis of Japanese

People Using the MOBI-Kids International Case-Control Study. Bioelectromagnetics. 05 November 2025. doi: 10.1002/bem.70032.

Abstract

This study investigated the association between mobile phone use and brain tumors in young people aged 10–29 years, specifically within the Japanese population. Building upon the international MOBI-Kids study, this Japanese case-control study included 120 brain tumor patients and 360 controls treated in the Kanto area, broadening the age range and including all primary brain tumors, unlike the international study. The data were evaluated using unconditional logistic regression, both unadjusted and adjusted for age and sex, with respect to the duration and intensity of mobile phone use. The analysis accounted for differences in the output power of various mobile phone generations prevalent in Japan, particularly the high-output 2G PDC system. Despite this rigorous exposure modeling, the study found no increased risk of brain tumors associated with mobile phone use, even when factoring in the weighted power output. The findings align with other recent international studies that also report no causal association. While acknowledging limitations such as potential recall bias and a relatively small sample size for sub-analyses, the study's strengths include its expanded age range and a realistic assessment of Japanese mobile phone usage patterns. Due to the ever-changing exposure conditions from wireless technology, further studies are needed to address their relation to young people's health.

Summary

Our study, focusing on young people in Japan, found no association between brain tumors and mobile phone use, based on a rigorous case-control design expanding the MOBI-Kids international case control study.

Analysis accounted for mobile phone generation differences, including Japan's high-output 2G PDC system with different phone types and power outputs.

https://onlinelibrary.wiley.com/doi/10.1002/bem.70032

--

The Influence of Mobile Technologies on the Quality of Sleep

Slezáková Z, Baláž J, Wimmerová S, Valachovičová M. The Influence of Mobile Technologies on the Quality of Sleep. J Biomed Phys Eng. 2025 Oct 1;15(5):491-496. doi: 10.31661/jbpe.v0i0.2311-1685.

Abstract

The extensive usage of mobile phones across all age brackets, including children and teenagers, leads to significant exposure to Radiofrequency Electromagnetic Fields (RF-EMF). This exposure raises concerns about potential adverse effects on sleep. The current study aimed to explore the influence of mobile phone-emitted RF-EMFs on the duration and quality of sleep in a cohort of medical students. Participants alternated between sleeping with and without a mobile phone for two-week intervals, while their sleep patterns were monitored using smartwatches. The study results indicated no statistically significant disparities in sleep quality between sleeping with and without a mobile phone. However,

exposure to radiofrequency electromagnetic fields had a notable impact on the minimum and average blood oxygen saturation levels.

Excerpt

During sleep, the deep sleep phase is considered crucial for physical recovery as it facilitates the release of growth hormones and regeneration of organs, muscles, and cells. In our study, we did not observe any statistically significant differences in the durations of wakefulness, REM sleep, light sleep, and deep sleep. However, we did find that RF EMF had a significant impact on minimum and average blood oxygen saturation levels. These findings emphasize the importance of conducting further comprehensive research in this area.

Open access: https://jbpe.sums.ac.ir/article_50575_eb6407c3fe71aef14e1f2dbc24bbcadb.pdf

--

Symptoms associated with environmental factors are positively related to sensory-processing sensitivity

Reddington K, Witthöft M, Köteles F, Szemerszky R. Symptoms associated with environmental factors are positively related to sensory-processing sensitivity. J Psychosom Res. 2025 Oct 25;199:112425. doi: 10.1016/j.jpsychores.2025.112425.

Abstract

Objective: Symptoms associated with environmental factors (SAEFs) are conditions characterized by chronic somatic symptom distress and impaired everyday functioning. SAEFs are positively related to certain psychological characteristics, such as somatosensory amplification (SSA), worries about the negative health consequences of modern technologies (MHWs), and body awareness. It was proposed recently that SAEFs might also be associated with sensory-processing sensitivity (SPS), i.e., the trait-like perceived sensitivity to internal and external sensory information.

Methods: Participants of a cross-sectional survey (n = 491; M_{age} = 34.7 yrs., SD_{age} = 14.2 yrs.; 76.4 % women) completed questionnaires assessing four SAEFs (perceived sensitivity to chemicals, buildings, sounds, electromagnetic fields) with multiple questions, and somatic symptom distress, body awareness, SSA, MHWs, and SPS.

Results: Somatic symptoms distress, SSA, body awareness, and SPS were significantly positively related to all four SAEFs. A very strong association ($r = 0.73$, $p < .001$) between SSA and SPS was found.

Conclusion: SPS can be considered a newly described (although not independent) psychological concomitant of SAEFs. Understanding the background of the association between SSA and SPS can be helpful to better understand SAEFs.

https://pubmed.ncbi.nlm.nih.gov/41160934/

--

Effects of Simultaneous In-Vitro Exposure to 5G-Modulated 3.5 GHz and GSM-Modulated 1.8 GHz Radio-Frequency Electromagnetic Fields on Neuronal Network Electrical Activity and Cellular Stress in Skin Fibroblast Cells

Hurtier A, Patrignoni L, Canovi A, Orlacchio R, Tjiou H, Gannes FP, Garenne A, Lévêque P, Arnaud-Cormos D, Lagroye I, Lewis N, Percherancier Y. Effects of Simultaneous In-Vitro Exposure to 5G-Modulated 3.5 GHz and GSM-Modulated 1.8 GHz Radio-Frequency Electromagnetic Fields on Neuronal Network Electrical Activity and Cellular Stress in Skin Fibroblast Cells. Bioelectromagnetics. 2025 Oct;46(7):e70026. doi: 10.1002/bem.70026.

Abstract

The widespread deployment of 5G wireless networks alongside existing GSM technologies has increased the need to assess potential biological effects of co-exposure to multiple radiofrequency electromagnetic fields (RF-EMF). This study evaluates the in-vitro impact of simultaneous exposure to 5G-modulated 3.5 GHz and GSM-modulated 1.8 GHz signals on neuronal electrical activity, mitochondrial reactive oxygen species (ROS) production, and cellular stress protein responses in neurons and skin fibroblasts. Primary cortical neurons and human immortalized skin fibroblasts were exposed to RF-EMF at specific absorption rates (SAR) of 1 or 4 W/kg for 15 min or 24 h, respectively. Neuronal activity was analyzed using multi-electrode arrays (MEAs), mitochondrial ROS production was measured using MitoSOX Red, and stress protein activity was assessed using bioluminescence resonance energy transfer (BRET) assays targeting RAS, PML, and HSF1 proteins. The results indicate no significant effects on the mean bursting rate (MBR) or mean firing rate (MFR) of cortical neurons, consistent with previous findings at similar SAR levels. Mitochondrial ROS production in fibroblasts also remained unaffected by RF-EMF co-exposure. BRET assays detected minor variations in the basal activity of RAS and PML and in the maximal efficacy of PMA and As_2O_3 to activate these pathways. However, these effects were small, near the detection threshold, and showed no consistent pattern across different tests or chemical treatments. No change was observed in HSF1 basal activity or responsiveness to MG132. These findings suggest that co-exposure to 5G- and GSM-modulated RF-EMF at SAR levels up to 4 W/kg does not produce conclusive evidence of marked biological effects under the tested conditions. Observed variations, when present, are of low amplitude and likely to fall within the range of experimental variability.

Summary

- Co-exposure to 5G-modulated 3.5 GHz and GSM-modulated 1.8 GHz at a total SAR of 4 W/kg (15 min) did not alter spontaneous bursting or firing rates in cortical neuronal networks.
- Twenty four hours co-exposure to 5G-modulated 3.5 GHz and GSM-modulated 1.8 GHz at a total SAR of 1 or 4 W/kg did not change mitochondrial ROS levels in human skin fibroblasts.
- BRET assays in fibroblasts showed that 24 h co-exposure to 5G-modulated 3.5 GHz and GSM-modulated 1.8 GHz at a total SAR of 1 or 4 W/kg produced only small changes, with slight increases in basal RAS and PML activity and minor variations in their chemical responses, while HSF1 remained unaffected.

Excerpts

Knowing the widespread deployment of the 5G wireless network at 3.5 GHz alongside the Global System for Mobile Communications (GSM) at 1.8 GHz, this study aims to investigate in-vitro effects of simultaneous exposure to both signals at SAR levels up to 4 W/kg on neuronal electrical activity, as well as ROS production and stress protein responses in skin fibroblasts. At such carrier frequencies, the

penetration depth into biological tissues is approximately 1 cm. As a result, the primary targets of interest are the skin cells and the central nervous system, particularly cortical neurons....

As illustrated in Figure 1, two exposure setups were used for 5G-modulated 3.5 GHz and GSM-modulated 1.8 GHz co-exposure depending on the cell model of interest. The setups were based on a cell culture incubator converted into a reverberation chamber exposure system (Figure 1A) for exposing skin fibroblasts, and an open transverse electromagnetic cell (TEM cell) containing a microelectrode array (MEA) placed in a classical cell culture incubator for neurons exposure (Figure 1B). The signal generation components were located outside incubators. For both exposure systems, the external signal unit was composed of two RF signal generators (SMBV100A, Rohde & Schwarz, Munich), a power combiner (Mini-circuits, ZB2PD-63-N+, NY, USA) combining 5G-modulated 3.5 GHz and GSM-modulated 1.8 GHz signals, a 25-dB gain amplifier (Mini-circuits, ZHL-4W-422+, NY, USA), a bidirectional coupler (Mini-circuits, ZGBDC30-372HP+, NY, USA), and a power meter (Keysight N1912A, USA) connected to two power sensors (Keysight N1921A, USA). The 3.5 GHz signal emitted by the radiofrequency generator is a real 5G NR signal (release 15, Digital Standards SMBVB-K444; Rohde & Schwarz) with FDD duplexing, QPSK modulation and 100 MHz channel bandwidth. The signal is also characterized by the power level and the peak envelope power (PEP), for example, for a power level of -10 dBm the peak envelope power is 1.26 dBm.

https://pubmed.ncbi.nlm.nih.gov/41104878/

--

Male Reproductive and Cellular Damage After Prenatal 3.5 GHz Radiation Exposure: One-Year Postnatal Effects

Dolanbay EG, Mert T, Bender GC, Bektas H, Uslu U, Fernandez-Rodriguez CE, Dasdag S. Male Reproductive and Cellular Damage After Prenatal 3.5 GHz Radiation Exposure: One-Year Postnatal Effects. 23 October 2025. doi: 10.1111/nyas.70116.

Abstract

This study investigates the long-term effects of prenatal exposure to 3.5 GHz radiofrequency radiation (RFR) on male reproductive health. Pregnant Wistar Hannover rats were divided into sham control, full-gestation exposure (3T RFR), and late-gestation exposure (2T RFR) groups (2 h/day). Male offspring were euthanized at 12 months for testicular analysis. In the 3T RFR group, seminiferous tubule diameter and epithelial height were significantly reduced compared to controls (adjusted $p = 0.03$ and 9.71×10^{-8}), along with lower Johnsen scores (adjusted $p = 0.022$). Abnormal sperm morphology increased significantly (adjusted $p = 0.036$). γ-H2AX immunostaining scores were elevated in the 2T and 3T groups (adjusted $p = 0.012$ and 6.36×10^{-9}). Beclin-1 expression was significantly higher in the 3T group versus sham and 2T groups (adjusted $p = 8.55 \times 10^{-4}$ and 4.51×10^{-6}). TUNEL-positive cell counts were significantly higher in both RFR groups than in sham (adjusted $p = 8.77 \times 10^{-18}$ for 3T, 6.42×10^{-17} for 2T), as was the apoptosis index (adjusted $p = 8.77 \times 10^{-18}$ for 3T, 5.66×10^{-17} for 2T). All p values were Holm–Bonferroni corrected. These findings indicate that prenatal exposure to 3.5 GHz RFR results in persistent testicular damage, impaired spermatogenesis, and increased DNA damage, autophagy, and apoptosis in adult male rats.

Graphical Abstract

Despite the accumulated knowledge of its effects, the long-term consequences of prenatal radiofrequency radiation (RFR) exposure on male reproductive health remain underexplored. Widespread usage makes understanding the potential impact of RFR on molecular pathways crucial. Thus, the purpose of this study is to investigate whether the reproductive systems of male rats exposed to 3.5 GHz RFR exclusively in the prenatal period were affected in the first year after birth.

Excerpts

A GSM signal generator (model 3500 PM10, Everest Comp., Adapazarı, Turkey), which generates a radiofrequency waveform modulated in the 3.5 GHz GSM band, was utilized to expose rats to RFR resembling the frequency to that generated by mobile phones. The power output of the signal generator was kept at 1 W during the exposure period. The piece of the generator that functions as the antenna was designed to replicate that of a conventional mobile phone....

EMR 300 (NARDA, Pfullingen, Germany) was used to measure the electric fields. E field measurements were performed with an EMR 300 in the corners and at the center of the cages we used in our study, which are the same size as standard animal cages. E fields were measured as 24 V/m, 26 V/m, 26.3 V/m, and 27 V/m at the corners, while 28 V/m was measured at the center of the cage....

The SAR calculations were conducted using the IEEE/IEC 62704-1 standard to assess the distribution of SAR values. The psSAR for 1 g was 0.06622 mW/g, and for 10 g was 0.03825 mW/g....

Conclusion

This study involved monitoring the male offspring of pregnant rats exposed to 3.5 GHz RFR exclusively during the prenatal period, with no exposure to electromagnetic waves for a period of one year following birth. The most original aspect of this study is that young male rats were kept alive for 1 year without any RF radiation exposure except during the prenatal period, and at the end of this period, their reproductive systems were examined. The examinations conducted in the first year after birth showed that exposure to 3.5 GHz RFRs only during the prenatal period poses significant risks to male rat reproductive health. Another original aspect of the study is that the first of the pregnant groups was exposed to RFRs during the three trimesters and the other during the last two trimesters.

Histological changes, increased abnormal sperm morphology, DNA damage, increased autophagic activity, and increased apoptosis are among the observed effects. These findings revealed that the reproductive systems of male offspring exposed to RFRs exclusively during pregnancy were negatively affected, even though they were not exposed to any EMFs for a year after birth, and the observed effects were even more pronounced in the full trimester group. In summary, this study showed that exposure to radiofrequency radiation (RFR) during pregnancy may have a negative impact on the reproductive health of subsequent male generations. Therefore, further molecular and cellular studies are needed to improve our understanding of the relationship between prenatal exposure to 3.5 GHz radiofrequency radiation (RFR) and subsequent male reproductive toxicity, and the mechanisms involved. Future investigations should extend the experimental timeline by incorporating multiple postexposure time points (e.g., 1, 6, and 12 months) to allow for a more comprehensive evaluation of the long-term effects of prenatal RFR exposure on reproductive function. Additionally, detailed assessments of sperm parameters such as motility, as well as organ weights, were not included in the current study. These aspects represent limitations and are planned to be addressed in ongoing and

future studies to provide a more comprehensive understanding of the systemic and functional consequences of prenatal RFR exposure. Finally, such research is crucial for developing evidence-based guidelines and implementing precautionary measures to safeguard reproductive health.

https://nyaspubs.onlinelibrary.wiley.com/doi/10.1111/nyas.70116

--

Synergistic Effects of 2600 MHz Radiofrequency Exposure and Indomethacin on Oxidative Stress and Gastric Mucosal Injury in Rats

Oruç S, Aral BS, Eraslan E, *et al.* Synergistic Effects of 2600 MHz Radiofrequency Exposure and Indomethacin on Oxidative Stress and Gastric Mucosal Injury in Rats. *Bratisl. Med. J.* (2025). doi: 10.1007/s44411-025-00382-4.

Abstract

Background/Aim With the widespread use of mobile technologies and NSAIDs, concerns have emerged regarding their potential combined impact on gastric health. This study investigated the synergistic effects of 2600 MHz RFF exposure and IND, a commonly used NSAID, on oxidative stress and gastric injury in rats.

Materials and Methods Forty male Wistar albino rats were randomly assigned to five groups: control, sham, IND, RFF, and IND + RFF. Animals were exposed to 2600 MHz RFF (1 h/day, 5 days/week) for 15 days, and/or received IND (4 mg/kg) via intragastric gavage. Gastric tissues were evaluated biochemically for oxidative stress markers (TOS, OSI, MDA, GSSG), antioxidant parameters (TAC, SOD, CAT, GSH, GPx), histologically for tissue damage, and immunohistochemically for HSP70 expression and apoptosis.

Results Single exposure to IND or RFF significantly elevated oxidative stress markers and decreased antioxidant levels in gastric tissues ($p < 0.001$). Combined IND + RFF exposure resulted in markedly greater gastric injury, with pronounced increases in HSP70 expression and apoptotic cell counts compared to single exposures ($p < 0.001$).

Conclusion Co-exposure to 2600 MHz RFF and IND markedly intensified oxidative stress, apoptosis, and gastric mucosal injury in rats. These findings suggest that the combined exposure produces a synergistic detrimental effect, exacerbating gastric tissue damage beyond the impact of each factor alone.

link.springer.com/article/10.1007/s44411-025-00382-4

--

Effect of Electromagnetic Field on Oral Tissues: A Narrative Review

Mallipattana V, Satelur KP, Shah MN, Thirugnana P, Ramnarayan BK.(2025). Effect of Electromagnetic Field on Oral Tissues: A Narrative Review. J Clin of Diagn Res. *19*(10), ZE01-ZE04. doi: 10.7860/JCDR/2025/79533/21893.

Abstract

An emerging pattern in clinical practice involves patients presenting with unexplained systemic and neurological symptoms despite normal diagnostic workups, with metallic or dissimilar-metal dental restorations being a recurring feature. The present narrative review explores potential links between Electromagnetic Field (EMF) exposure, dental materials and systemic health effects, with a particular focus on neurological manifestations. Drawing on principles from quantum biology and physics, the review examines unique clinical cases that defy conventional medical explanations yet may be interpreted through interactions between EMFs and metal restorations at the quantum level. These cases suggest that the human body may act as a biological interface where classical and quantum physical phenomena intersect. While the review emphasises evidence-based findings, it also presents quantum-level hypotheses—such as spin dynamics and Radical-Pair Mechanisms (RPM)—for interdisciplinary consideration, underscoring the complexity of EMF-biological interactions and the need for further investigation.

Open access: https://www.jcdr.net/article_fulltext.asp?issn=0973-709x&year=2025&month=October&volume=19&issue=10&page=ZE01-ZE04&id=21893

--

Electromagnetic Field Stimulation Effects on Intrinsically Disordered Proteins and Their Role in Aging and Neurodegeneration

Perez FP, Bandeira J, Morisaki J, Kanakri H, Rizkalla M. Electromagnetic Field Stimulation Effects on Intrinsically Disordered Proteins and Their Role in Aging and Neurodegeneration. J Biomed Sci Eng. 2025 Oct;18(10):408-438. doi: 10.4236/jbise.2025.1810030.

Abstract

There is increasing evidence from preclinical studies. There is growing evidence from preclinical studies in cell cultures and small organisms that exposure to Electromagnetic Fields (EMFs) produces beneficial biological effects. However, controversy persists due to the absence of a clearly defined mechanism. Classical physics, constrained by the non-ionizing nature of these exposures, cannot account for these effects, which do not involve the breaking of chemical bonds to induce conformational changes in proteins. Emerging studies suggest that these effects are mediated through quantum mechanical phenomena-specifically, quantum tunneling and particle-wave duality-acting on the water surrounding proteins at their interfaces. Furthermore, we present evidence of EMF-induced conformational changes in Intrinsically Disordered Proteins (IDPs), including beta-amyloid, tau, alpha-synuclein, and Heat Shock Factor 1 (HSF1). These findings offer a new framework for understanding EMF bioeffects and open promising avenues for research in biophysics and quantum biology. In this context, we address the challenge of reproducibility by examining how variables such as frequency, intensity, Specific Absorption Rate (SAR), and exposure time windows interact, along with how parameters like polarization, phase, pulse modulation, and scheduling influence outcomes. Experimental data identify specific RF frequencies and SAR levels that activate proteostasis and autophagy in cell cultures and small animal

models, with potential applications in human treatments that remain consistent with safety thresholds established by regulatory agencies.

Open access: https://www.scirp.org/journal/paperinformation?paperid=146794

--

RF-EMF Exposure Assessment: Comparison of Measurements in Airports and Flights with and Without Wi-Fi Service

Arribas E, Escobar I, Martinez-Plaza A, Rufo-Pérez M, Jimenez-Barco A, Paniagua-Sánchez JM, Marín P, Ramirez-Vazquez R. RF-EMF Exposure Assessment: Comparison of Measurements in Airports and Flights with and Without Wi-Fi Service. *Sensors*. 2025; 25(21):6710. doi: 10.3390/s25216710.

Abstract

This paper presents the results of personal exposure measurements to Radiofrequency Electromagnetic Fields from 2.4 GHz and 5.85 GHz Wi-Fi frequency bands. Measurements were taken in several specific scenarios: within international airports terminals, during takeoff, inside airplanes while flying with and without onboard Wi-Fi service (including while actively using a Wi-Fi connection), and during landing. Data were recorded onboard four international flights (two-round trip flights), from Spain to Mexico, and from Spain to Belgium. Two personal exposimeters, EME SPY 140 and EME Spy Evolution, were used to collect intensity level measurements in each scenario. During the outbound, the mean exposure value inside the airplane flight was 93.9 µW/m2 in the 2.4 GHz Wi-Fi frequency band and 46.4 µW/m2 in the 5.85 GHz Wi-Fi band (Spain to Mexico), and 7.29 µW/m2 in the 2.4 GHz Wi-Fi band and 2.40 µW/m2 in the 5.85 GHz Wi-Fi band (Spain to Belgium). For the return flight, the average value was 26.7 µW/m2 in the 2.4 GHz Wi-Fi band and an average of 9.87 µW/m2 in the 5.85 GHz Wi-Fi band (Mexico to Spain), and 3.24 µW/m2 in the 2.4 GHz Wi-Fi band and 1.23 µW/m2 in the 5.85 GHz Wi-Fi band (Belgium to Spain). Personal exposure levels to RF-EMFs from the Wi-Fi frequency band inside an airplane, even at the airport, are very low and well below the reference levels established by the international guidelines (10 W/m2).

Open access: https://www.mdpi.com/1424-8220/25/21/6710

--

Exposure Variability Between 1- or 6-Minute and 30-Minute Averaging Time Lengths in Radiofrequency-Electromagnetic Field Exposure Monitoring

Bhatt CR, Henderson S, Sanagou M, Loughran S. Exposure Variability Between 1- or 6-Minute and 30-Minute Averaging Time Lengths in Radiofrequency-Electromagnetic Field Exposure Monitoring. Bioelectromagnetics. 2025 Dec;46(8):e70030. doi: 10.1002/bem.70030.

Abstract

Different averaging time lengths (ATLs) are widely used in radiofrequency-electromagnetic field (RF-EMF) exposure monitoring. This study evaluated variability in RF-EMF exposure associated with the ATLs of 1- and 6-min normalised to 30-min data. For 15 frequency bands of interest, RF-EMF exposures were collected in contiguous 1-min blocks over 30 min at four sites (two outdoor and two indoor). Frequency-

band and site-specific variability in exposure between the three ATLs was assessed. First, the variability in terms of relative deviation (in dB) between 1- or 6-min and 30-min were estimated. Second, the overall exposure variability (in $\mu W/m^2$) were compared between 1- or 6-min and 30-min blocks statistically using the quantile regression method. The overall exposure variability on ATL of 1-min or 6-min was considered different to 30-min if a majority of respective sub-pair comparisons across different percentiles (P_5 , P_{25} , P_{50} , P_{75} , P_{95}) were significantly different. The study largely showed that the exposure variability (i.e., relative deviation) of different ATLs was within +/− 3 dB. Further, the overall exposure variability between 1- or 6-min and 30-min ATLs showed inconsistent outcomes. Frequency bands of broadcast and most of the mobile services < 2 GHz demonstrated overall similar exposure variability for 1-min and 6-min ATLs.

https://onlinelibrary.wiley.com/doi/10.1002/bem.70030

--

Visualizing radiofrequency electromagnetic field exposure through Voronoi-based maps

Arribas E, Ramirez-Vazquez R, Escobar I. Visualizing radiofrequency electromagnetic field exposure through Voronoi-based maps. Environ Sci Pollut Res Int. 2025 Nov 15. doi: 10.1007/s11356-025-37188-4.

Abstract

Measuring the exposure of radiofrequency electromagnetic fields in a city is a very laborious task. To simplify this process for a city of 200 thousand inhabitants, Voronoi diagrams were used. The city was divided into cells based on the Euclidean distance to a point on the map, and each cell was assigned the value of the electric field measured at that point using a personal exposimeter. The number of cells varied from five to 30 and, finally, one 100. The maps obtained are commented on, using a palette of four uniform colors for the cells, to facilitate their perception. The more cells are considered, the smaller the cells will be. A stabilization of the process is observed as the points within the city map increase. The colors represent the RMS (root mean square) electric field measured at each seed point, extended geometrically to its Voronoi cell for visualization purposes. The colors of the areas remain the same; and in some other areas, cells appear with slightly distinct colors, due to the addition of new measured points. Some cells change color due to these new measurements of the new points. In this study, the predominant color is green, which is the measured field at the seed point was 1.9 V/m RMS (while the maximum allowed by the ICNIRP guidelines is 61.4 V/m). There are three cells in the city with E values above 3.9 V/m, reaching the highest value of 11.4 V/m. The entire city is within the recommended maximum. The Voronoi diagram method is shown to be useful and interesting for the presentation of radiofrequency electromagnetic fields exposure measurements in a medium-sized city. The Voronoi cells do not contain information about the intracell spatial variation of the electric field; each polygon represents the RMS value measured at its generating point (seed), offering a discrete visualization of the spatial distribution of the measurements and illustrating how the electric field levels vary across the study area.

https://pubmed.ncbi.nlm.nih.gov/41239101/

--

Empowering the Serbian EMF RATEL System for Monitoring RF-EMF Through Drive Test

Djuric N, Kavecan N, Otasevic V, Kljajic D, Djuric S. Empowering the Serbian EMF RATEL System for Monitoring RF-EMF Through Drive Test. *IEEE Access*, vol. 13, pp. 180918-180940, 2025, doi: 10.1109/ACCESS.2025.3618714.

Abstract

The Serbian EMF RATEL system has been monitoring radiofrequency (RF) electromagnetic fields (EMFs) since 2017. As one of five active European systems for continuous monitoring of RF-EMF, it has been able to provide insight into the long-term, temporal variability of the field, especially in areas with increased sensitivity to RF-EMF exposure. Although it has proven its benefits for the general public and the scientific community, the relatively small number of RF-EMF sensors installed at fixed locations reveals its weakness – the inability to cover larger urban areas and provide adequate information on the spatial distribution of the overall field. The distribution of sensors at scattered points is a consequence of the limited costs of system development, as well as the practical difficulties in obtaining an authorized permit to install sensors at a public or private location, which is crucial for the use of any large number of sensors. Consequently, the EMF RATEL network has been upgraded to perform spatial field analysis by integrating the functionality of the drive test, for the first time among existing monitoring systems, developing it to a higher level and achieving progress in RF-EMF research by establishing a single EMF entity for observing both temporal and spatial changes of RF-EMF in the territory of interest. This paper presents some technical details of the integration of the drive test into EMF RATEL, as well as preliminary quantitative results of the corresponding drive test measurements to validate the approach. The aim of this paper is to encourage other, similar EMF monitoring systems to further develop the topic of RF-EMF observation.

Open access: https://ieeexplore.ieee.org/document/11195077

--

Thermal and SAR-Based Limits for Human Skin Exposed to Terahertz Radiation

Elmaadawy S, Jornet JM. Thermal and SAR-Based Limits for Human Skin Exposed to Terahertz Radiation. *2025 International Telecommunications Conference (ITC-Egypt)*, Cairo, Egypt, 2025, pp. 388-393, doi: 10.1109/ITC-Egypt66095.2025.11186625.

Abstract

Terahertz technology holds strong potential for wireless communication and sensing and has become a growing focus of research in recent years. However, current safety guidelines for wireless communication do not fully address the entire terahertz spectrum, as they only cover the sub-terahertz band (100−300GHz). Establishing safe exposure limits is a necessary prerequisite for the practical deployment of terahertz systems. In this paper, a COMSOL Multiphysics model was developed to analyze the thermal impact of terahertz exposure on human skin and to determine the maximum permissible power limits across the frequency range from 0.1 THz to 5 THz. The current state of terahertz wireless communication technology was also explored by analyzing a representative system to assess present capabilities and exposure safety. The results indicate that the existing power levels defined for the sub-terahertz band may no longer be adequate, either for the sub-terahertz range itself

or for the broader terahertz region evaluated in this study. This research lays the groundwork for safer terahertz technology standards, facilitating its broader use in wireless communication.

Conclusions

This study established the first comprehensive thermal and SAR safety analysis for human skin exposed to terahertz radiation across the 0.1−5THz spectrum. Our COMSOL Multiphysics ® simulations revealed that maintaining SAR below the 1.6 W/kg safety threshold requires power densities not exceeding 0.226 W/m2, which induced negligible temperature increases. Temporal SAR analysis demonstrated that even stricter limits may be necessary for prolonged exposures. Notably, while power densities up to 100 W/m2 produced minimal thermal effects (approximately 0.03 K), they generated SAR values that far exceeded safety standards. These findings directly informed safe operating parameters for emerging terahertz wireless communication systems. Our results demonstrated that current sub-THz safety standards could not be extrapolated to the broader THz spectrum, necessitating new exposure guidelines.

Our analysis was limited to the 0.1−5THz range and did not extend to frequencies up to 10 THz, where additional thermal effects may emerge. Moreover, our model assumed uniform skin structure, whereas actual skin exhibits complex layered heterogeneity that could affect radiation absorption patterns.

Future work will incorporate experimental validation of these findings through phantom studies and extend the analysis to investigate potential long-term biological effects of terahertz radiation.

https://ieeexplore-ieee-org.libproxy.berkeley.edu/document/11186625

--

Prolonged 3.5 GHz and 24 GHz RF-EMF Exposure Alters Testicular Immune Balance, Apoptotic Gene Expression, and Sperm Function in Rats

Syed Taha SMA, Jaffar FHF, Hairulazam A, Vijay S, Jamaludin N, Zulkifli AF, Mat Ros MF, Osman K, Zakaria Z, Mohd Bahar MAA, et al. Prolonged 3.5 GHz and 24 GHz RF-EMF Exposure Alters Testicular Immune Balance, Apoptotic Gene Expression, and Sperm Function in Rats. *Biomedicines*. 2025; 13(10):2471. https://doi.org/10.3390/biomedicines13102471

Abstract

Background/Objectives: The rapid rollout of 5G has renewed interest in potential reproductive effects of mid-band (3.5 GHz) and millimeter-wave (24 GHz) radiofrequency electromagnetic fields (RF-EMF). We examined frequency- and duration-dependent changes in testicular cytokines, apoptosis-related genes, and sperm quality in rats.

Methods: Male Sprague Dawley rats (n = 6 per group) were exposed for 60 days to 3.5 GHz or 24 GHz RF-EMF for 1 h/day or 7 h/day. The sham controls were housed identically. Testicular expressions of IL-10, IL-6, IL-1β, and TNF-α were quantified; Tp53, Bax, Bcl2, and Casp3 mRNA expressions were measured; and sperm concentration, viability, and motility were evaluated.

Results: IL-10 was significantly reduced in the 24 GHz group at both 1-h and 7-h exposure duration. At 7

h, TNF-α was also lower at 24 GHz. Casp3 expression was higher and Tp53 was lower at 3.5 GHz at 1-h exposure duration. Sperm concentration and viability were reduced after 24 GHz exposure at 7 h, while sperm motility was reduced after 3.5 GHz exposure at both durations.

Conclusions: Exposure to RF-EMF 3.5 GHz primarily impacts sperm motility via extrinsic pro-apoptotic pathways, while exposure to 24 GHz impacts sperm concentration and viability potentially through immune–apoptotic mechanisms, with all negative effects amplified by 7-h daily exposure.

Open access: https://www.mdpi.com/2227-9059/13/10/2471

--

The effect of Wi-Fi on elastic and collagen fibres in the blood vessel wall of the chorioallantoic membrane

Holovská K, Andrašková S, Petrovová E, Molnár J, Vince T, Almášiová V. The effect of Wi-Fi on elastic and collagen fibres in the blood vessel wall of the chorioallantoic membrane. Acta Vet. Brno 2025, 94: 137-144. doi: 10.2754/avb202594020137

In this study we observed the effect of Wi-Fi on various fibrous components that form the wall of large blood vessels in the chorioallantoic membrane (CAM) of the chicken embryo. Chicken embryos in the experimental groups (Wi-Fi) were continuously exposed (24 h per day) to Wi-Fi radiation at a frequency of 2.4 GHz and an average power density of 300 μW/m^2 for 9 and 14 embryonic days (ED). Subsequently, the CAM samples from the control (CO) and experimental (Wi-Fi) groups were histologically processed and evaluated. The samples stained with haematoxylin and eosin did not reveal any notable differences in the structure of large blood vessels between the CO and Wi-Fi groups. However, the use of special staining methods showed adverse effect of Wi-Fi on the fibrous elements within the blood vessel wall. The optical density (OD) of elastic fibres significantly decreased from 0.294 ± 0.025 (CO 9) to 0.197 ± 0.022 (Wi-Fi 9) at ED 9 and from 0.706 ± 0.028 (CO 14) to 0.271 ± 0.031 (Wi-Fi 14) at ED 14. On the other hand, at ED 9, the OD of collagen fibres exhibited a notable increase from 0.178 ± 0.023 (CO 9) to 0.334 ± 0.020 (Wi-Fi 9). However, at ED 14, there was a significant decline of collagen fibres from 0.418 ± 0.033 (CO 14) to 0.327 ± 0.031 (Wi-Fi 14). Our results support the hypothesis that Wi-Fi could affect the fibrous elements of the large vessel wall and may play a role in the development of different cardiovascular disorders.

Open access: https://actavet.vfu.cz/media/pdf/actavet_2025094020137.pdf

--

Investigating the Effects of Occupational Noise and Extremely Low-Frequency Electromagnetic Field Exposure on Oxidative Response in Power Plant Workers

Jafarimanesh S, Ehsani H, Shaki F, Moosazadeh M, Samaei SE. Investigating the Effects of Occupational Noise and Extremely Low-Frequency Electromagnetic Field Exposure on Oxidative Response in Power Plant Workers. Bioelectromagnetics. 2025 Oct;46(7):e70027. doi: 10.1002/bem.70027.

Abstract

Occupational noise and extremely low-frequency electromagnetic fields (ELF-EMF) are common in power plants and represent important risk factors that may contribute to oxidative stress. This study examined how simultaneous exposure to these hazards affects oxidative stress biomarkers in workers under real-world conditions. Participants were assigned to one of four exposure groups: Control (C), Noise (N), ELF-EMF (E), or a combined Noise and ELF-EMF group (NE). Occupational noise and ELF-EMF exposures were measured according to ISO 9612 and IEEE Std C95.3.1, respectively. To assess oxidative stress, venous blood samples were collected from all participants, and plasma levels of malondialdehyde (MDA), glutathione (GSH), superoxide dismutase (SOD), and total antioxidant capacity (TAC) were analyzed using validated biochemical assays. The NE group showed the highest MDA levels, indicating elevated lipid peroxidation compared with controls ($p < 0.001$). GSH concentrations were lower in NE relative to controls ($p < 0.001$). SOD activity was significantly reduced in both the N and NE groups compared with the control group ($p < 0.005$). TAC was lowest in the NE group, showing a significant decrease compared with both the control and Noise-only groups ($p < 0.05$). While these findings suggest that concurrent exposure to noise and ELF-EMF can influence oxidative stress biomarkers, they do not provide direct evidence to mandate specific workplace monitoring or interventions. Further studies are needed to clarify potential health risks and to guide evidence-based occupational safety measures.

Summary

- This study investigated oxidative stress biomarkers in power plant workers exposed to occupational noise, ELF-EMF, and their combined exposure.
- Concurrent exposure to noise and ELF-EMF was linked to a significant increase in the lipid peroxidation marker malondialdehyde compared with unexposed workers.
- Other oxidative stress markers, including glutathione, superoxide dismutase, and total antioxidant capacity, exhibited less consistent changes across the exposure groups.
- These findings indicate potential oxidative stress effects associated with combined occupational exposures, highlighting the need for further research.

https://pubmed.ncbi.nlm.nih.gov/41104887/

--

Measurement of Electromagnetic Fields Exposure to Humans from Electric Vehicles and Their Supply Equipment

Murthy GPP, Varadarajan GS.Measurement of Electromagnetic Fields Exposure to Humans from Electric Vehicles and Their Supply Equipment.Measurement. Volume 258, Part D,2026, doi: 10.1016/j.measurement.2025.119378.

Abstract

Electric Vehicles (EVs) represent a progressive approach to fostering a healthy and sustainable transportation future on Earth. Intended for mitigating global warming by virtue of decarbonizing transportation sector. Besides being environmentally beneficial, electrification in transportation necessitates a study of non-ionizing Electromagnetic Fields (EMF). Despite extensive research, the health effects related to long-term exposure to EMF remain inconclusive. The main aim of this research is to

share and disseminate measurement findings of EMF emission from EVs and their Electric Vehicle Supply Equipment (EVSE). The specification of the measuring instrument, methodology adhered to measurement, and observed results have been discussed. Measurements have been carried out at different EVs and EVSEs in and around Chennai city, up to 400 kHz frequency band. To analyze human exposures, Electric Field Intensity (E) and Magnetic Flux Density (B) have been measured in the vicinity of driver and passenger seats inside EVs during driving condition, furthermore fields near different positions of EVSEs during charging condition. The measured values have been compared with the ICNIRP and IEEE standard recommendations for human external body exposure level. In accordance with the findings, E & B inside EVs fall within the standard limit. The E in the vicinity of EVSE is under standard limit, but at certain places B is concurrently above general public exposure threshold (>200 μT). In addition, a preliminary computational analysis has been done using Finite Element Method (FEM). Simulated fields are then compared with measurement readings of AC charger, where the electric and magnetic field magnitudes are relatively higher. With an ever-increasing electrification of vehicle fleets, knowledge of health impacts of long-term exposure to humans can be enhanced through more observational research.

Highlights

• Measurement of Electric Field Intensity (E) and Magnetic Flux Density (B) from EVs and EVSEs while driving and charging respectively.
•The measured values have been compared with the ICNIRP and IEEE standard recommendations for human external body exposure level.
• E and B inside EVs falls within the standard limit.
• The E in the vicinity of EVSEs is also under standard limits but B is concurrently above the standard exposure threshold at certain positions.
• FEM model of electric vehicle AC charging cable using COMSOL Multiphysics® software for EMF computational analysis has been done.
• The measurement results are compared with simulation outputs.
• From the findings, policy recommendations to mitigate EMF exposure from EVs and EVSEs in the context of global electric vehicle adoption are addressed.

https://www.sciencedirect.com/science/article/pii/S026322412502737X

--

A Cohort Study on Alzheimer's Disease in Relation to Residential Magnetic Fields From Indoor Transformer Stations

Liimatainen A, Roivainen P, Juutilainen J, Höytö A, Naarala J. A Cohort Study on Alzheimer's Disease in Relation to Residential Magnetic Fields From Indoor Transformer Stations. Bioelectromagnetics. 2025 Dec;46(8):e70031. doi: 10.1002/bem.70031.

Abstract

Meta-analyses of epidemiological studies have suggested that Alzheimer's disease (AD) may be linked with exposure to extremely low frequency (ELF) magnetic fields (MF). This is the first study investigating the association of AD with exposure to residential ELF MFs from indoor transformer stations, using a

study design that avoids shortcomings of previous studies. All cohort members had lived in buildings with indoor transformer stations. MF exposure was assessed based on the location of their apartment in relation to the transformer room. AD patients were identified from Drug Purchase Register and Drug Reimbursement Register. Out of the 155,562 individuals, 5652 (111,357 person-years of follow-up) living in apartments next to transformer stations were considered as exposed, while 115,772 (2,289,526 person-years of follow-up) individuals living on higher floors of the same buildings were considered as referents. Associations between MF exposure and AD were examined using Cox proportional hazard models. The hazard ratio (HR) was 1.02 (95% confidence interval: 0.85-1.22), indicating that the risk of AD is not associated with residential ELF MFs present in apartments next to transformer stations. The duration of residence did not essentially change the HR. The risk of AD was slightly but not statistically significantly higher (HR 1.22, 95% confidence interval: 0.94-1.57) for those whose residence started before the age of 50 years. The results did not support positive findings from previous studies that have reported a link between AD and occupational or residential MF exposure.

Summary

- The study included high residential extremely low frequency (ELF) magnetic field (MF) levels from indoor transformer stations and avoided many limitations of previous studies.
- Residential ELF MFs did not affect the risk of Alzheimer's disease (AD).
- Further studies should focus on investigating occupational ELF MF exposure and the potential effect of diurnal timing of MF on the risk of AD.

Conclusions

No evidence for an association between residential ELF MFs and AD was found in this study that involved relatively high exposure levels and avoided many limitations of previous studies addressing residential MF exposure. We recommend that, if any further studies on the possible link between ELF MF exposure and AD are conducted, they should focus on occupational MF exposure, trying to identify occupational groups exposed to particularly strong ELF MFs. Studies (including experimental studies) on the possible differences between night-time and day-time exposure might also be helpful.

Open access: https://onlinelibrary.wiley.com/doi/10.1002/bem.70031

--

Effect of Increased Ionizing Radiation and Near-Null Magnetic Field on Electrical Signals of Plants

Grinberg MA, Il'in NV, Nemtsova YA *et al.* Effect of Increased Ionizing Radiation and Near-Null Magnetic Field on Electrical Signals of Plants. *Radiophys Quantum El* (2025). https://doi.org/10.1007/s11141-025-10418-y.

Abstract

There is currently a significant gap in understanding the effects of astro- and geophysical factors such as ionizing and non-ionizing radiation, including electromagnetic fields, on living organisms. The effects caused by these factors are important for planning scenarios in which significant changes in the intensity of such factors can be expected both on the Earth and during space missions. There is a reason to believe that, of all the indicators of the functioning of living organisms, the signaling systems responsible

for the perception, processing, and amplification of the incoming stimulus are the most sensitive. In this work, we have studied the effect of increased levels of ionizing radiation (β radiation, dose rate of 31.3 μGy/h) and hypomagnetic conditions (induced magnetic field 0–1.5 μT) on electrical signals of plants caused by an additional stimulus. To conduct the experiments, a setup based on triaxial Helmholtz rings was created, which permits one to set and maintain altered conditions of the electromagnetic environment with high accuracy. The work showed that an increased level of ionizing radiation strengthens, while the near-null level of magnetic field weakens electrical signals in plants. It is assumed that the mechanism of action of such factors is their effect on a single component of stress signaling, namely, concentration of reactive oxygen species. The observed effects open up a new way for studying the mechanisms of the influence of ionizing and non-ionizing radiation on the state of living organisms.

https://link.springer.com/article/10.1007/s11141-025-10418-y

--

Electromagnetic Interference in the Modern Era: Concerns, Trends, and Nanomaterial-Based Solutions

Prekodravac Filipovic J, Milenkovic M, Kepic D, Dorontic S, Yasir M, Nardin B, Jovanovic S. Electromagnetic Interference in the Modern Era: Concerns, Trends, and Nanomaterial-Based Solutions. Nanomaterials (Basel). 2025 Oct 13;15(20):1558. doi: 10.3390/nano15201558.

Abstract

Electromagnetic interference (EMI) represents a growing challenge in the modern era, as electronic systems and wireless technologies become increasingly integrated into daily life. This review provides a comprehensive overview of EMI, beginning with its historical evolution over centuries, from early power transmission systems and industrial machinery to today's complex environment shaped by IoT, 5G, smart devices, and autonomous technologies. The diverse sources of EMI and their wide-ranging effects are examined, including disruptions in electrical and medical devices, ecological impacts on wildlife, and potential risks to human health. Beyond its technical and societal implications, the economic dimension of EMI is explored, highlighting the rapid expansion of the global shielding materials market and its forecasted growth driven by telecommunications, automotive, aerospace, and healthcare sectors. Preventative strategies against EMI are discussed, with particular emphasis on the role of advanced materials. Carbon-based nanomaterials-such as graphene, carbon nanotubes, and carbon foams-are presented as promising solutions owing to their exceptional conductivity, mechanical strength, tunable structure, and environmental sustainability. By uniting perspectives on EMI's origins, consequences, market dynamics, and mitigation strategies, this work underscores the urgent need for scalable, high-performance, and eco-friendly shielding approaches. Special attention is given to recent advances in carbon-based nanomaterials, which are poised to play a transformative role in ensuring the safety, reliability, and sustainability of future electronic technologies.

Open access: https://www.mdpi.com/2079-4991/15/20/1558

--

The WHO-commissioned systematic reviews on health effects of radiofrequency radiation provide no assurance of safety

Melnick RL, Moskowitz JM, Héroux P, Mallery-Blythe E, McCredden JE, Herbert M, Hardell L, Phillips A, Belpoggi F, Frank JW, Scarato T, Kelley E, on behalf of the International Commission on the Biological Effects of Electromagnetic Fields (ICBE-EMF). The WHO-commissioned systematic reviews on health effects of radiofrequency radiation provide no assurance of safety. Environ Health 24, 70 (2025). https://doi.org/10.1186/s12940-025-01220-4.

Abstract

The World Health Organization (WHO) commissioned 12 systematic reviews (SR) and meta-analyses (MA) on health effects of exposure to radiofrequency electromagnetic fields (RF-EMF). The health outcomes selected for those reviews (cancer, electromagnetic hypersensitivity, cognitive impairment, birth outcomes, male fertility, oxidative stress, and heat-related effects) were based on a WHO-conducted international survey. The SR of the studies of cancer in laboratory animal studies was the only one that did not include a MA, because those authors considered it inappropriate due to methodological differences among the available studies, including differences in exposure characteristics (carrier frequency, modulation, polarization), experimental parameters (hours/day of exposure, duration of exposure, exposure systems), and different biological models. MAs in all the other SRs suffered from relatively few primary studies available for each MA (sometimes due to excessive subgrouping), exclusion of relevant studies, weaknesses in many of the included primary studies, lack of a framework for analyzing complex processes such as those involved in cognitive functions, and/or high between-study heterogeneity. Due to serious methodological flaws and weaknesses in the conduct of the reviews and MAs on health effects of RF-EMF exposure, the WHO-commissioned SRs cannot be used as proof of safety of cell phones and other wireless communication devices. However, the animal cancer SR, which was rated as “high certainty of evidence” for heart schwannomas and “moderate certainty of evidence” for brain gliomas, provided quantitative information that could be used to set exposure limits based on reducing cancer risk. The multiple and significant dose-related adverse effects found in the SRs on male fertility and pregnancy and birth outcome should also serve as the basis for policy decisions to lower exposure limits and reduce human reproductive risks. The report of harmful effects (e.g., cancer, reproductive toxicity, etc.) at doses below the adverse health effect threshold claimed by ICNIRP demonstrates that current exposure limits to RF-EMF, which were established by applying arbitrary uncertainty factors to their putative adverse threshold dose, lack scientific credibility.

Open access: https://ehjournal.biomedcentral.com/articles/10.1186/s12940-025-01220-4

Supplementary Material 1. "Examples of Working Group Authors’ Ties to Industry"

Supplementary Material 2. "The Meta-Analyses in the WHO RF-EMF Systematic Reviews Yielded Unreliable Results"

https://www.saferemr.com/2022/10/international-commission-on-biological.html

--

Review of the evidence on the influence of Wi-Fi 2.4 GHz radiation on oxidative stress and its possible relationship with Alzheimer’s disease

Laván D, Argüelles N, Lluncor A, Huaman D, Moyano J, Ubillus J, Peña M, Paredes M, Hernández I, Guerra A, De La Cruz-Vargas J, Cruz V. Review of the evidence on the influence of Wi-Fi 2.4 GHz radiation

on oxidative stress and its possible relationship with Alzheimer's disease. Frontiers in Neurology. Volume 16, 2025. doi: 10.3389/fneur.2025.1616435.

Abstract

To date, there is no scientific consensus on whether wireless communication systems, such as 2.4 and 5 GHz Wi-Fi, play a crucial role in the development of Alzheimer's disease through oxidative stress. Although numerous studies have linked oxidative stress to exposure to electromagnetic radiation from wireless systems in various biological contexts, these studies have not established a direct connection to neurodegenerative diseases. Research on Alzheimer's disease and oxidative stress is an active field in neuroscience and medicine, as oxidative stress involves an imbalance between the production of free radicals and the antioxidant system's ability to neutralize them, leading to cellular and neuronal damage. It is essential to recognize that Alzheimer's disease is multifactorial, and its development generally results from a complex interaction of genetic, environmental, and lifestyle factors. The relationship between wireless systems such as Wi-Fi and oxidative stress, as well as its possible link to Alzheimer's disease, continues to be the subject of research and debate in the scientific community. Although some studies have explored this possible association, the results have been mixed and inconclusive. While research on the health effects of wireless systems remains relevant, it is prudent not to consider this association as an established fact until solid and consistent scientific evidence is available. The study we present focuses on indirectly analyzing the relationship between genes that respond to oxidative stress upon exposure to Wi-Fi 2.4 GHz electromagnetic waves and genes associated with the development of Alzheimer's disease. Our results indicate that the modification of key genes involved in neurodegeneration, such as GSK3B and APOE, could be exacerbated by prolonged exposure to this radiation. It is essential for future research to explore this hypothesis to further clarify the potential risks associated with electromagnetic radiation and its impact on neuronal health and the progression of Alzheimer's disease.

Conclusion

In summary, exposure to 2.4 GHz electromagnetic fields emitted by Wi-Fi devices could have an indirect impact on the regulation of genes involved in Alzheimer's disease, particularly those related to oxidative stress and cellular homeostasis. Although a direct relationship has not been demonstrated, current findings suggest that the alteration of genes such as GSK3B and APOE, which are fundamental in neurodegeneration, could be exacerbated by chronic exposure to this radiation. Future research should address this hypothesis to provide a clearer understanding of the potential risks associated with electromagnetic radiation and its impact on neuronal health and the development of Alzheimer's disease.

Open access: https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2025.1616435

--

Microwave and RF Exposure-Induced Molecular and Genetic Alterations

Lin JC. Microwave and RF Exposure-Induced Molecular and Genetic Alterations [Health Matters]. *IEEE Microwave Magazine*, vol. 26, no. 11, pp. 16-17, Nov. 2025, doi: 10.1109/MMM.2025.3594272.

No abstract

"Recent reports of studies on the effects of microwave and RF exposure on molecular and cytogenetic materials have drawn considerable attention. They suggest that RF exposure could induce DNA damage at levels considered safe by promulgated guidelines and regulatory standards.

In particular, a new study described the observation of genetic alterations in rat cancers resulting from lifetime exposure to low-level RF radiation [1]...."

[1] A. M. Brooks et al., “Genetic profiling of rat gliomas and cardiac schwannomas from life-time radiofrequency radiation exposure study using a targeted next-generation sequencing gene panel,” *PLoS One*, vol. 19, no. 1, 2024, Art. no. e0296699, doi: 10.1371/journal.pone.0296699.

Open access: https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=11195922

--

Experience of Polish Physicians on Electromagnetic Hypersensitivity

Undas K, Kanclerz G, Popielak J, Tatoń G. Experience of Polish Physicians on Electromagnetic Hypersensitivity. *2025 Progress in Applied Electrical Engineering (PAEE)*, Koscielisko, Poland, 2025, pp. 1-4, doi: 10.1109/PAEE68231.2025.11155985.

Abstract

The aim of this project was to assess the knowledge of Polish physicians about the impact of electromagnetic fields (EMF) on human health, their knowledge of electromagnetic hypersensitivity (EHS), and their experiences with patients who consider themselves hypersensitive to EMF. A questionnaire survey was conducted among physicians in a traditional paper-based format, as well as using an online form. Responses were obtained from 355 doctors of various specialties and with different levels of experience, both clinical and scientific. A subjective and objective assessment of the respondents' knowledge about the impact of EMF on humans leads to the disturbing conclusion that their knowledge cannot be considered broad. For example, the subjective self-assessment of knowledge in 1/3 of the respondents is at a very low or low level. Physicians do not seem to be familiar with the WHO guidelines for dealing with people who perceive themselves as hypersensitive to EMF. Respondents are aware of the low level of their knowledge about EMF and most of them are ready to improve their qualifications in this field. They also see the need to spread reliable knowledge about the EMF and its impact on society. The low assessment of doctors' knowledge of EMF and EHS is even more concerning as EHS is a phenomenon many encounter in their daily professional practice. Almost 45% of physicians declared that they had to deal with patients who associated their symptoms with the impact of EMF, and some physicians claim that they deal with such patients even several times a month or more often. These findings may lead to significant for public health conclusions. EHS is not a marginal phenomenon, and physicians are striving to properly address this condition. At the same time, their knowledge about EHS and about EMF in general is limited. In such cases, they are unable to adequately help their patients and may become a source of unreliable or unverified information, which could lead to anxiety.

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=11155985&isnumber=11155907

--

Is Cellphone Carrying Below the Waist (Exposure to Non-Ionizing Radiation) Contributing to the Rapid Rise in Early-Onset Colorectal Cancer?

Li DK, Ferber J, Hirst A, Odouli R, Quesenberry C, Levin T. Is Cellphone Carrying Below the Waist (Exposure to Non-Ionizing Radiation) Contributing to the Rapid Rise in Early-Onset Colorectal Cancer? ISEE 2024: 36th Annual Conference of the International Society of Environmental Epidemiology. Vol. 2024, Issue 1. doi: 10.1289/isee.2024.1903.

Abstract

BACKGROUND AND AIM: The incidence of early-onset (EO) (< 50 years of age) colorectal cancer (CRC) has been rising rapidly in recent decades, suggesting the presence of a new environmental risk factor. One emerging, yet under-investigated factor is increased exposure to radiofrequency (RF) non-ionizing radiation from widespread cellphone use. The young population has been documented to carry their cellphones in pants' pockets many hours a day, leading to extensive exposure to RF radiation in the colorectal region. Intestinal epithelial cells are among the tissue cells most sensitive to radiation exposure. The study was designed to investigate whether cellphone carrying below the waist is associated with increased risk of EOCRC.

METHOD: We conducted a case-control study among 50 recently diagnosed EOCRC cases and 50 randomly selected age- and sex-matched controls. Information on carrying habits included specific locations below the waist before EOCRC diagnosis, the average amount of time carried per day and the ages at which they started and stopped carrying at a given location.[¤]RESULTS[|]After controlling for confounders, a longer duration (the highest tertile) in total hours of carrying a cellphone below the waist had more than 4-fold risk of EOCRC than shorter carrying duration (lower tertiles), adjusted odds ratio (aOR)=4.1, 95% confidence interval (CI):1.0-16.4. The association was stronger for rectal cancer (OR=6.4) vs. colon cancer (OR=1.9), given the closer proximity of pants' pockets to the rectum. Most strikingly, the association was much stronger for ipsilateral carrying (cellphone carrying on the same side of EOCRC tumor location) with a OR=12.1 (1.5-97) vs. contralateral carrying (cellphone carrying on the opposite side of tumor location) with OR=1.6.

CONCLUSIONS: A longer duration of cellphone carrying below the waist is associated with an increased risk of EOCRC. Though preliminary, the finding revealed an emerging environmental contributor to the rapid rise in EOCRC, whose reasons remain unknown.

https://ehp.niehs.nih.gov/doi/10.1289/isee.2024.1903

--

Causal relationship between the duration of mobile phone use and risk of stroke: A Mendelian randomization study

Jin R, Jiang X, Lin Q, Huang X, Gao W, Wang B, Zheng F. Causal relationship between the duration of mobile phone use and risk of stroke: A Mendelian randomization study. Medicine (Baltimore). 2025 Sep 19;104(38):e44485. doi: 10.1097/MD.0000000000044485.

Abstract

This study investigates the causal relationship between the duration of mobile phone use (DMPU) and risk of stroke using Mendelian randomization (MR) analysis. Independent single nucleotide polymorphisms from genome-wide association study datasets were employed as instrumental variables to estimate the effects of DMPU on the risk of stroke, intracerebral hemorrhage, ischemic stroke, and its subtypes (cardioembolic infarction, small-vessel disease, large artery atherosclerosis [LAAS]). Inverse-variance weighting was utilized as the primary MR method and sensitivity analyses were performed. Ninety single nucleotide polymorphisms associated with stroke from genome-wide association study datasets were selected as instrumental variables. Inverse-variance weighted analysis showed a significant causality between DMPU and an increased risk of LAAS (odds ratio [OR] = 1.120; 95% confidence interval [CI] = 1.005–1.248; *P* = .040). No genetic association was found for stroke (OR = 1.000; 95% CI = 0.999–1.001, *P* = .677), intracerebral hemorrhage (OR = 1.020; 95% CI = 0.912–1.140, *P* = .734), ischemic stroke (OR = 1.020; 95% CI = 0.979–1.062, *P* = .344), cardioembolic infarction (OR = 1.066; 95% CI = 0.974–1.166, *P* = .166), and small-vessel disease (OR = 1.052; 95% CI = 0.944–1.173, *P* = .356). MR-Egger regression (intercept = 2.75×10^{-3}; *P* = .888) suggested multidimensionality was unlikely to bias the results; Cochran *Q* test and funnel plot showed no heterogeneity or asymmetry, indicating the robustness of present findings. The current investigation confirmed a causal relationship between DMPU and an increased risk of LAAS, suggesting significant implications for public health initiatives and policy development.

Summary

This study used Mendelian randomization to explore if the duration of mobile phone use (DMPU) affects stroke risk. Researchers analyzed genetic data to see if DMPU is linked to different types of stroke. They found a significant link between DMPU and an increased risk of large artery atherosclerosis (LAAS), with an odds ratio of 1.120. However, no genetic association was found for other stroke types, including intracerebral hemorrhage and ischemic stroke. The study suggests that DMPU may specifically increase the risk of LAAS, highlighting potential public health concerns and the need for policy development.

Open access: https://journals.lww.com/md-journal/fulltext/2025/09190/mobile_phone_usage_duration_and_male_fertility__a.4.aspx

--

Mobile phone usage duration and male fertility: A two-sample Mendelian randomization analysis

Xiang Y, Xu L, Sun Y, Hu C, Lv L. Mobile phone usage duration and male fertility: A two-sample Mendelian randomization analysis. Medicine (Baltimore). 2025 Sep 19;104(38):e44668. doi: 10.1097/MD.0000000000044668.

Abstract

Mobile phone use has become ubiquitous in modern life, raising public concern over its potential effects on male reproductive health. While several observational studies have reported associations between prolonged phone use and decreased sperm quality or testosterone levels, these findings remain inconclusive due to residual confounding, reverse causation, and inconsistent exposure measurement. We conducted a two-sample Mendelian randomization (MR) analysis using genetic variants associated with mobile phone usage duration to investigate its causal effects on male fertility. Inverse variance weighting (IVW) was employed as the primary analytical method. Genetic predisposition to longer

mobile phone use was not associated with levels of sex hormone-binding globulin (SHBG), total testosterone, or the risk of abnormal sperm parameters, erectile dysfunction, or testicular dysfunction. These null findings were consistent across all MR methods and sensitivity analyses. Our findings suggest that mobile phone use is unlikely to have a direct causal impact on male reproductive hormones or sexual function. Future research incorporating precise exposure measurements and mechanistic evaluations is warranted.

Open access: https://journals.lww.com/md-journal/fulltext/2025/09190/mobile_phone_usage_duration_and_male_fertility__a.4.aspx

--

Differential metabolic responses of mouse Leydig and spermatogonia cells to radiofrequency electromagnetic field exposure

Miao Xia , Lin Yanyun , Guo Juan , Lin Jiajin , Gao Peng , Zhang Wei , Zeng Lihua , Guo Guozhen , Li Jing. Differential metabolic responses of mouse Leydig and spermatogonia cells to radiofrequency electromagnetic field exposure. Frontiers in Public Health. Volume 13 - 2025. doi: 10.3389/fpubh.2025.1623701.

Abstract

Introduction: Although existing studies have shown that radiofrequency electromagnetic fields (RF-EMFs) have a variety of effects on living organisms, the specific impact of RF-EMFs on the metabolism of reproductive cells and their underlying mechanisms remain unclear.This study aims to explore the effects of RF-EMFs on the metabolism of mouse Leydig cells (TM3) and spermatogonia cells (GC-1) through metabolomics analysis, revealing the potential mechanisms by which RF-EMFs affect reproductive health.

Methods: We employed liquid chromatography-mass spectrometry (LC-MS) to analyze the metabolomic profiles of TM3 and GC-1 cells under two irradiation modalities: continuous and intermittent RF-EMF exposure. The data were further analyzed using KEGG pathway analysis to identify significantly enriched metabolic pathways. The ELISA (Enzyme-Linked Immunosorbent Assay) was used to detect glutathione levels.

Results: Our results showed that continuous irradiation had a more pronounced impact on the metabolism of TM3 cells, primarily affecting amino acid metabolism, the citric acid cycle, ABC transporters, bile secretion, and glutathione metabolism. In contrast, intermittent irradiation mainly altered the levels of fatty acyls and purine nucleosides, with significant enrichment in purine metabolism, biosynthesis of unsaturated fatty acids, and fatty acid metabolism. Compared to TM3 cells, GC-1 cells exhibited lower sensitivity to RF-EMF irradiation. Both irradiation modalities affected purine metabolism and lysine degradation pathways in TM3 cells, suggesting that changes in ADP levels may serve as a key metabolic signature in the cellular response to RF-EMF exposure.

Conclusion: Continuous irradiation significantly impacts TM3 cell metabolism, particularly amino acid and glutathione pathways, while intermittent irradiation mainly affects fatty acyls and purine

metabolism. GC-1 cells show lower sensitivity to RF-EMF. ADP level changes may be a key metabolic signature of RF-EMF exposure.

https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1623701/full

--

Radiofrequency regulates the BET-mediated pathways in radial glia differentiation in human cortical development

Cakir B, Tanaka Y, Choe MS, Kiral FR, Kim J, Micali N, Kang YJ, Dharmadhikari B, Patterson B, Yang WS, Cho Y, Xiang Y, Zhong M, Lee SH, Patra P, Rakic P, Park IH. Radiofrequency regulates the BET-mediated pathways in radial glia differentiation in human cortical development. Cell Rep. 2025 Sep 24:116238. doi: 10.1016/j.celrep.2025.116238.

Abstract

The human brain represents one of the most complex organs in our body, with development regulated by an intricate genetic program. Recently, non-genetic factors, such as prenatal stress, infection, and diet, have been shown to influence brain development. Radiofrequency radiation (RF; 800-2,400 MHz), emitted by natural and artificial sources such as microwaves and cell phones, represents a non-invasive environmental factor. Using human cortical organoids (hCOs) derived from human embryonic stem cells (hESCs), we investigate RF's effects on corticogenesis. We find that RF exposure regulates the differentiation of human and non-human primate radial glia progenitors, maintaining stem cell identity and delaying differentiation. Neurons differentiated under RF treatment show induction of expression of human endogenous retroviruses. Importantly, inhibitors for the BET (bromodomain and extraterminal) protein rescue RF-induced developmental defects in hCOs. Our findings reveal a mechanism by which RF modulates early brain development, offering a non-biological approach to regulate neural stem cell self-renewal.

Highlights

RF exposure alters radial glial differentiation in human cortical organoids

RF induces ASD-related gene and retroelement expression in cortical organoids

BET protein dysregulation mediates RF-induced neurodevelopmental defects

BET inhibitors rescue RF-induced developmental defects in cortical organoids

Open access: https://www.sciencedirect.com/science/article/pii/S2211124725010095

--

Investigation of fetal exposure to electromagnetic waves between 2.45 and 5 GHz during pregnancy

İl N, Ateş K, Özen Ş. Investigation of fetal exposure to electromagnetic waves between 2.45 and 5 GHz during pregnancy. Radiat Prot Dosimetry. 2025 Sep 25:ncaf110. doi: 10.1093/rpd/ncaf110.

Abstract

In this paper, we have assessed the specific absorption rate (SAR) in fetal brain and lungs during the second trimester of pregnancy conditions of body exposure to radiofrequency electromagnetic field (RF-EMF). SAR calculations were performed for frontal and lateral incidences, with both vertical and horizontal polarization of the incident electromagnetic (EM) waves at frequencies ranging from 2.45 to 5 GHz. A realistic pregnant human model with a belly button piercing was implemented for numerical simulation for nonionizing dosimetry. The simulation results reveal that SAR tends to rise in the presence of belly-button piercing. The highest SAR10g recorded on the fetus's lungs was 16 mW/kg at a frequency of 2.45 GHz. Similarly, the maximum SAR10g value on the fetus's brain was measured 14 mW/kg, occurring at a frequency of 2.45 GHz. Results indicate that metal objects can lead to an increase in SAR values. However, obtained values remain below limits set by international organizations like the Institute of Electrical and Electronics Engineers and the International Commission on Non-Ionizing Radiation Protection.

https://pubmed.ncbi.nlm.nih.gov/40995977/

--

3.5GHz radiofrequency electromagnetic fields (RF-EMF) on metabolic disorders in Drosophila melanogaster

Wang Y, Zhang Z, Zhang L, Liao Y, Cai P. 3.5GHz radiofrequency electromagnetic fields (RF-EMF) on metabolic disorders in Drosophila melanogaster. Ecotoxicol Environ Saf. 2025 Oct 6;304:119132. doi: 10.1016/j.ecoenv.2025.119132.

Abstract

The widespread implementation of fifth-generation (5 G) technology has raised concerns regarding its impact on biosafety. This study aims to uncover metabolic alterations in Drosophila melanogaster in response to electromagnetic radiation through metabolomics under 3.5 GHz radiofrequency electromagnetic fields (RF-EMF) conditions. Newborn male flies were continuously exposed to 3.5 GHz RF-EMF at intensities of 0.1 W/m^2, 1 W/m^2, and 10 W/m^2 throughout their entire life cycle under 24-hour constant darkness. Metabolomic analysis was performed on Drosophila samples after long-term exposure. The results indicate that long-term exposure to RF-EMF leads to disruptions in four metabolic pathways - alanine, aspartate and glutamate metabolism, tryptophan metabolism, ascorbate and aldehyde metabolism, and purine metabolism - as well as 34 differential metabolites across various power density in Drosophila melanogaster. Notably, the most significant metabolites (GABA, glucose-6-phosphate, AMP, N-formylglycinamide nucleotide, LPC, and MG) exhibited high sensitivity to RF-EMF showing significant decreases. The suggests that long-term exposure to electromagnetic wave treatment may reduce the overall metabolic levels in fruit fly. These findings provide valuable insights into the metabolic disturbance induced by RF-EMF and enhance our understanding of the potential health effects associated with 5 G radiation.

Highlights

Long-term 3.5 GHz RF-EMF exposure disrupts metabolism in Drosophila melanogaster.
Key pathways affected: energy, neurotransmitter, membrane lipid, purine metabolism.
34 altered metabolites did not exhibit a clear dose-dependent response to RF-EMF exposure.

Six sensitive biomarkers identified (e.g., GABA, glucose-6-phosphate), with significant decreases. Results provide mechanistic insights into 5 G RF-EMF biosafety impacts.

Open access: https://www.sciencedirect.com/science/article/pii/S0147651325014770?via%3Dihub

--

Bacterial Adaptation to Radiofrequency Electromagnetic Fields Based on Experiences from Ionizing Radiation

Said-Salman I, Mortazavi SMJ, Khatib SE, Mortazavi SA, Sihver L. (2025). Bacterial Adaptation to Radiofrequency Electromagnetic Fields Based on Experiences from Ionizing Radiation. *Journal of Biomedical Physics and Engineering*, (), -. doi: 10.31661/jbpe.v0i0.2405-1770.

Abstract

Bacteria, part of the three domains of life (Eukarya, Archaea, and Bacteria), are constantly exposed to man-made electromagnetic fields, which often exceed the intensity of natural electromagnetic sources. In response to this exposure, bacteria have developed various defensive and resistant traits. This article presents an overview of both historical and recent research on how bacteria adapt to common sources of Radiofrequency Electromagnetic Fields (RF-EMF. The widespread use of mobile phones and Wi-Fi, both utilizing Radiofrequency (RF) radiation, raises potential public health concerns, which have been addressed by international organizations like the World Health Organization (WHO). Understanding how bacteria adapt to EMF is important for mitigating the risk of increased pathogenicity of radio-resistant bacteria in the human environment.

Conclusion

The exposure of bacteria to EMF can be appreciated when altering with bacterial survival mechanism and may compromise therapeutic success. However, continuous exposure of microorganisms to common sources of EMF such RF may emerge super-pathogens with high resistance to treatments.

Open access: https://jbpe.sums.ac.ir/article_51139_0eca0c408ce6ec99ffd5908dc60c3449.pdf

--

Associations between Individual and Geospatial Characteristics and Power of 4G Signals Received by Mobile Phones

Laplanche A, Guida F, Moissonnier M, Launay L, Beranger R, Lagroye I, Orlacchio R, Fontaine M, Bories S, Mazloum T, Conil E, Huss A, Wiart J, Danjou A, Schüz J, Dejardin O, Deltour I. Associations between Individual and Geospatial Characteristics and Power of 4G Signals Received by Mobile Phones. Environ Res. 2025 Oct 6:123030. doi: 10.1016/j.envres.2025.123030.

Abstract

Background: The Received Signal Strength Indicator (RSSI) measures downlink signal intensity received by smartphones in 4th Generation LTE networks.

Objective: This study evaluated how individual, technical, and spatial factors influenced LTE-RSSI during daily activities.

Methods: Between November 2022 and October 2023, adults in France used the XMobiSensePlus Android smartphone application to record RSSI and GPS data. Distance to the operator's nearest antenna, obtained from Cartoradio, population and antenna density and urbanicity were analyzed using a geographic information system. Determinants of RSSI were assessed using an autoregressive mixed model incorporating restricted cubic splines for distance. Environmental exposures were estimated at 1800 MHz using conversion factors.

Results: From 1,969,913 records of 187 participants, with measurements taken every 30 seconds over 7.9 days, the average LTE-RSSI was -79.3 dBm. The estimated electric field strength was 0.12 V/m, albeit with large uncertainty. The median distance to the nearest antenna was 536 m. Proximity to antennas increased RSSI. Antenna density positively influenced RSSI (overall β = +0.37 dBm per additional antenna per km2). Lower RSSI was observed in the evening and night, particularly in urban areas. Smartphone's technical parameters (Android version and System-on-a-Chip) influenced RSSI, operators did not. Proximity to antennas had greater impact in rural areas.

Conclusion: Urbanicity, distance to the nearest 4G antenna, antenna density, time of day, and smartphone's technical parameters influenced RSSI levels in 4G networks in France, but not operator.

https://pubmed.ncbi.nlm.nih.gov/41061881/

--

Extremely Low-Frequency Magnetic Fields (ELF-MF) and Radiofrequency: Risk of Childhood CNS Tumors in a City with Elevated ELF-MF Exposure

Correa-Correa V, Núñez-Enríquez JC, Mezei G, Rivera-Luna R, Peñaloza-González JG, Rivas-Carrillo SD, et al. Extremely Low-Frequency Magnetic Fields (ELF-MF) and Radiofrequency: Risk of Childhood CNS Tumors in a City with Elevated ELF-MF Exposure. Environmental Research, 2025, doi: 10.1016/j.envres.2025.122858.

Abstract

Background Central nervous system tumors (CNSTs) are the second most common childhood cancer. While their etiology is unclear, exposure to extremely low-frequency magnetic fields (ELF-MF) and radiofrequency (RF) may be associated with increased risk. This study aims to examine the relationship between ELF-MF and RF exposure and CNST risk in Mexico City's pediatric population.

Methods A case-control study was conducted in Mexico City (2017-2022), to assess ELF-MF and RF exposure in 200 CNST patients and 793 controls under 16 years old. Residential ELF-MF exposure was measured over 24 hours, and RF exposure was assessed through the total duration of cell phone calls and electronic device usage (with and without internet). ELF-MF exposure levels below 0.1 μT and the lowest RF exposure quartile (Q1) served as reference groups for adjusted odds ratio (aOR) analyses. Data analysis was performed using R and SPSS software.

Results Of all participants, 5.1% had ELF-MF exposure ≥0.3 μT. Elevated ELF-MF exposure (≥0.4 μT) was significantly associated with an increased risk of CNST (aOR (95% confidence interval)=2.39 (1.15-5.00). While cell phone use showed no differences between cases and controls, prolonged tablet use (with and

without internet) was significantly associated with increased CNST risk (aOR=2.53 (1.39-4.61), and 3.53 (1.45-8.59), respectively).

Conclusion A high proportion of children in Mexico City are exposed to ELF-MF levels ≥0.3 μT, exceeding levels reported in other populations. Elevated residential ELF-MF exposure, as well as prolonged tablet use, both with and without internet connectivity, were significantly associated with an increased risk of developing CNST in childhood.

Highlights

- Many Mexico City children exposed to ELF-MF ≥0.3 μT, higher than global averages.
- Elevated ELF-MF exposure was associated with an increased risk of CNST.
- Prolonged tablet use increased CNST risk, especially in children aged 6-10.
- Cell phone use showed no association with CNST risk.

https://www.sciencedirect.com/science/article/pii/S0013935125021103

--

Dosimetric Electromagnetic Safety of People With Implants: A Neglected Population?

Kranold L, Xi J, Goren T, Kuster N. Dosimetric Electromagnetic Safety of People With Implants: A Neglected Population? Bioelectromagnetics. 2025 Sep;46(6):e70023. doi: 10.1002/bem.70023. P

Abstract

Electromagnetic (EM) safety guidelines are designed to protect the general public and workers from the risks posed by exposure to EM sources of all types, with the exception of medical EM sources. However, it has never been systematically evaluated whether individuals with conductive implants are also protected by these guidelines or whether the local field enhancement due to presence of the implant may pose an unacceptable risk under certain realistic exposure conditions. To address this important knowledge and regulatory gap, we first evaluated the upper bound of the local enhancement of bare and insulated generic implants of 0.5 λ (approximately equal to resonant) and 0.1 λ lengths, but restricted the maximum length to 2 m, as a function of tissue properties and frequency (10 kHz to 1 GHz). Results for uniform electric field excitation showed local enhancement of $psSAR_{10mg}$ and of locally averaged E-field, respectively, compared to the background in the presence of a generic implant of 10 dB (1 GHz) to over 100 dB at frequencies under 100 MHz. In the next step, we tested the hypothesis that fields induced inside the human body by realistic near-field sources are not sufficiently uniform to generate results in enhancement that could pose unacceptable risk. Common implant trajectories were inserted into the Virtual Population human anatomical model Ella V3.0, and the model was exposed to the following conditions (i) a standard source representing a wireless power transfer source operating at 85 kHz and (ii) a dipole source that operates at 450 MHz within the current exposure limits. Results show that the safety limit is exceeded at the tip of the implant by a factor of > 10 (> 20 dB) or > 115 V/m at 85 kHz, whereas the locally induced specific absorption rate averaged over 10 mg at 450 MHz was 7.9 W/kg, resulting in a temperature increase after 6 min of < 0.4 K. Hence, as the hypothesis was falsified at frequencies < 450 MHz, patients with implants are inadequately protected by current safety and product guidelines. In the discussions, proposals for how to close this regulatory gap are provided.

Summary

- In this study, we evaluated systematically whether people with conductive implants are protected by electromagnetic safety guidelines and standards or whether certain exposure conditions result in an unacceptable risk to the subpopulation.
- A systematic exposure study of generic insulated and non-insulated implants showed local enhancements of the $psSAR_{10mg}$ and locally averaged E-field in a frequency range of 10 kHz to 1 GHz of up to 100 dB.
- The human anatomical model ViP Ella v3 with common implant trajectories was exposed to a realistic 85 kHz wireless power transfer coil and a 450 MHz dipole source, resulting in a local field enhancement at the implant tip of over > 115 V/m and 7.9 W/kg, respectively, showing that people with implants are not adequately protected by current safety guidelines.

Open access: https://pmc.ncbi.nlm.nih.gov/articles/PMC12464515/

--

Numerical analysis of the thermal effects on adult with brain pacemaker implantation exposed to WIFI antennas

Tian R, Zhao Y, Lu M, Deng J. Numerical analysis of the thermal effects on adult with brain pacemaker implantation exposed to WIFI antennas. Case Studies in Thermal Engineering, Volume 74, 2025, https://doi.org/10.1016/j.csite.2025.106941.

Abstract

In the regard of whether it's safe for the Parkinson's Disease (PD) patients with brain pacemaker (BP) implantation to use their cell phones, we investigated the electromagnetic (EM) radiation of cell phone to the patients. In this work, a three-dimensional model of the human brain including the real Basal Ganglia, two dual-frequency planar inverted-F antennas (PIFA) operating at WIFI/5G and two types of implanted BPs were modeled utilizing COMSOL finite element software. By varying the antennas operating frequency, the angles between the antennas and the human brain, the thermal conductivity of the Grey Matter (GM) and the electrode structure, we numerically analyzed the specific absorption rate (SAR), the temperature increase, the thermal strain and the displacement of the brain tissues. The results indicated that regardless of the frequencies and angles, the SAR value and temperature increase are below the limits of 2 W/kg and 2 °C specified by International Commission on Non-Ionizing Radiation Protection (ICNIRP) 2020, but they reached their maximum at the angle of 90°. The change in GM thermal conductivity and electrode geometry affected the trend of the temperature increase in the brain tissues. The thermal strain and displacement of brain tissues during their temperature increase were found to have a potential impact on the postoperative efficacy of PD patients. The displacement of brain tissues was the similar as the displacement that was observed in some medical literatures, and the two displacements approximately had the same order of magnitude. Therefore, we recommended that one should avoid using the cell phones in the gesture of θ= 90°. The PD patients should also maintain the distance as far as possible. Considering that the PD patients may use cell phones frequently, we also suggested that they might visit the hospital once in a while, in order to confirm that the pacemaker is working properly.

Open access: https://www.sciencedirect.com/science/article/pii/S2214157X25012018

--

Assessment of Electromagnetic Exposure Levels for Humans from Electric Vehicle DC Charging Stations

Dong S, Lu M. Assessment of Electromagnetic Exposure Levels for Humans from Electric Vehicle DC Charging Stations. Sensors (Basel). 2025 Sep 14;25(18):5735. doi: 10.3390/s25185735.

Abstract

The potential health risks of DC charging piles to human health were investigated by quantifying the internal electromagnetic exposure level. In this study, the transformer in the DC/DC circuit of a DC charging pile was selected as the radiation source, and two realistic human models (adult and child) were used as exposure subjects. A simulation model, including the vehicle body, charging pile, and transformer, was established using COMSOL(COMSOL Multiphysics 6.2) Multiphysics software to calculate the magnetic induction intensity (**B**-field) and electric field intensity (**E**-field) in various organs at distances of 0.1 m, 0.3 m, and 0.6 m from the charging pile. The results show that at 0.1 m, the peak **B**-field (1.91 μT) and **E**-field (447 mV/m) in the adult body were 1.91% and 2.07% of the ICNIRP occupational exposure limits, respectively, and 7.07% and 4.14% of the public exposure limits. For the child model, the peak electromagnetic exposure levels (2.31 μT and 259 mV/m) were only 8.56% and 2.40% of the public limits. Further evaluation of exposure levels for in-vehicle occupants during charging showed that the peak **B**-field and **E**-field for an adult driver and a child in the front passenger seat were 0.0225×10^{-2} μT, 0.0237×10^{-2} μT, 5.81 mV/m, and 5.82 mV/m, respectively, far below the ICNIRP public limits. Additionally, analyses at multiple frequency bands (85 kHz, 90 kHz, and 95 kHz) under a typical scenario (adult at 0.1 m from the charging pile) revealed that the **B**-field in the human body decreased with increasing frequency, while the **E**-field showed minimal variation due to shielding effects. All electromagnetic exposure levels were below both ICNIRP public and occupational limits, indicating the broad applicability of the results. Under normal operating conditions of DC charging piles, the electromagnetic exposure from the DC/DC transformer fully complies with safety standards and poses no threat to human health. This study provides a scientific basis for alleviating public concerns about the health risks of electromagnetic radiation from DC charging piles for electric vehicles.

Conclusions

This study uses the transformer in the DC/DC circuit of a DC charging station module as the radiation source and employs realistic adult and child human models as exposure subjects. At an 80 kHz switching frequency, the electromagnetic field excited by the AC signal input to the primary winding was calculated, and electromagnetic exposure levels were assessed for individuals in proximity to the charging station and for an adult in the driver's seat and a child in the front passenger seat during EV charging. The main conclusions are as follows:

(1) For an adult standing 0.1 m from the charging pile, the peak **B**-field (1.91 μT) and the peak **E**-field (447 mV/m) in the body are 1.91% and 2.07% of the ICNIRP occupational exposure limits, respectively, and 7.07% and 4.14% of the public exposure limits. For the child, the electromagnetic exposure levels (2.31 μT and 259 mV/m) are 8.56% and 2.40% of the public exposure limits, respectively. These results indicate that electromagnetic exposure levels for occupational and private individuals near the DC charging pile are within safe limits.

(2) For an adult standing at distances of 0.1 m, 0.3 m, and 0.6 m from the DC charging pile, the peak **B**-field in the body corresponds to 7.07%, 3.02%, and 0.80% of the ICNIRP public exposure limits, respectively, while the peak **E**-field corresponds to 2.07%, 1.77%, and 0.44%. For a child at the same distances, the peak **B**-field is 8.56%, 3.45%, and 2.15% of the ICNIRP public exposure limits, and the peak **E**-field is 2.40%, 1.74%, and 1.43%. These findings demonstrate a significant negative correlation between the distance from the charging pile and the electromagnetic exposure level in the body.

(3) During electric vehicle charging, the peak **B**-fields in the body for an adult in the driver's seat and a child in the passenger's seat are 0.0225×10^{-2} μT and 0.0237×10^{-2} μT, respectively. The peak **E**-fields in the body and brain are 5.81 mV/m and 5.59 mV/m for the adult, and 5.82 mV/m and 5.67 mV/m for the child, all significantly below the ICNIRP public exposure limits.

(4) In the typical scenarios (adult at 0.1 m from the charging pile), analyses were conducted at multiple frequency bands, including 85 kHz, 90 kHz, and 95 kHz. The results show that as the frequency increases, the peak **B**-field in the human body gradually decreases, while the **E**-field in the body shows no significant change, possibly influenced by the vehicle body. The electromagnetic exposure levels in all frequency bands are below the ICNIRP safety limits.

This study has certain limitations, and future research will focus on the following areas: (1) expanding the types of human models to include sensitive populations such as pregnant women and adolescent females; (2) evaluating electromagnetic exposure from multiple radiation sources (e.g., inverters and cables) acting concurrently; (3) conducting field measurements and validation of electromagnetic fields from DC charging stations.

This study addresses gaps in the safety assessment of electromagnetic environments associated with DC charging stations, providing a scientific basis for establishing electromagnetic environment standards and industry exposure limits for EV charging facilities, thereby contributing to public health protection.

Open access: https://www.mdpi.com/1424-8220/25/18/5735

--

Bioelectricity in Morphogenesis

Nunes CO, Barriga EH. Bioelectricity in Morphogenesis. Annu Rev Cell Dev Biol. 2025 Oct;41(1):187-208. doi: 10.1146/annurev-cellbio-101323-032747. Epub 2025 Aug 19.

Abstract

Bioelectricity is likely as old as life itself. From the moment the first proto-cell was enclosed in a lipid bilayer, a membrane potential arose. Thus, one can expect that bioelectrical activities influence single-cell and collective cell behaviors in processes such as embryo development, tissue repair, and even disease. Despite the ubiquity of bioelectrical phenomena, most research has focused on bioelectrical control of neural tissues, and as a result, our knowledge of nonneural contexts remains comparatively less understood, scattered, and often misunderstood. Still, there are strong reasons for supporting the idea that bioelectricity contributes to diverse morphogenetic contexts. Thus, in this review we provide an overview of the current knowledge of how cells generate and perceive bioelectrical inputs, and discuss how cells translate these stimuli into responses that influence tissue morphogenesis in physiology and pathology.

Open access: https://www.annualreviews.org/docserver/fulltext/cellbio/41/1/annurev-cellbio-101323-032747.pdf?expires=1759524214&id=id&accname=guest&checksum=4B7FD77ABD1E5E6E4D9239BE689002C4

--

Evaluation of Exposure Assessment Methods and Procedures for Induction Hobs (Stoves)

Xi J, Kühn S, Fortunato C, Ofli E, Kuster N. Evaluation of Exposure Assessment Methods and Procedures for Induction Hobs. Bioelectromagnetics. 2025 Oct;46(7):e70024. doi: 10.1002/bem.70024.

Abstract

Induction hobs generate strong alternating magnetic fields to heat pots by inducing eddy currents. These fields are the strongest close to the bottom of the cookware, but stray fields at large distances can still be substantial. In general, these are higher than the reference levels defined by international electromagnetic exposure safety guidelines (ICNIRP 1998; ICNIRP 2010; IEEE 2019). That the reference levels are exceeded does not imply that the basic restrictions are also violated. In this study, we assess the exposures caused by the latest generation of induction hobs by applying the advanced instrumentation and different methods that include the procedures developed by the International Electrotechnical Commission (IEC) for household appliances (IEC 62233) (IEC International Electrotechnical Commission 2005), the 4-tier approach developed for inductive wireless power transfer systems (IEC 63184) (IEC International Electrotechnical Commission 2021), and their derivatives. First, methods for determining the maximum exposure configuration were assessed. Then, the 3D distribution of the incident magnetic field was sampled with a scanning system and analyzed, and the contact currents assessed. Lastly, numerical dosimetric evaluations were performed in anatomical models to determine the maximum fields induced by the measured incident fields directly or by a representative coil model converted from the measured fields. The study's findings reveal significant variations in exposure across different induction hobs, with differences of up to a factor of > 20 (> 26 dB) as a function of power, coil size, and proximity to the coil. This suggests that low-exposure hobs can be designed without compromising cooking performance. Furthermore, the study strengthens the conclusions of previous studies that IEC 62233 (IEC International Electrotechnical Commission 2005) may underestimate the exposure for persons standing next to the hob by up to a factor of > 30-based on testing according to the exposure limits from (ICNIRP 1998; IEEE 2019)-and thus does not ensure safety. A dosimetric analysis, the most accurate method, would be relatively costly. Alternative approaches derived from (IEC International Electrotechnical Commission, 2021) that are affordable and not overly conservative are discussed.

Summary

- The low-exposure induction hobs can be achieved by employing small heating coils without compromising cooking performance.
- The product standards IEC 62233 (IEC International Electrotechnical Commission 2005) and EN 62233 (CENELEC, European Committee for Electrotechnical Standardization 2008) may significantly underestimate the exposure, and therefore require revision to ensure the safety of induction hobs.

- Conservative and affordable approaches from IEC 63184 (IEC International Electrotechnical Commission 2021) and IEC 60990 (IEC International Electrotechnical Commission 2016) are applied to demonstrate the compliance of induction hobs with the limits of incident/induced fields and contact current, respectively.

Open access: https://onlinelibrary.wiley.com/doi/10.1002/bem.70024

--

Simultaneous 4G and 5G EMF Exposure and Field Uniformity in a Reverberation Chamber for Animal Studies

Kwon D, Jeon S, Lee A-K, Choi H-D, Moon J-I. Simultaneous 4G and 5G EMF Exposure and Field Uniformity in a Reverberation Chamber for Animal Studies. *Applied Sciences*. 2025; 15(18):10286. https://doi.org/10.3390/app151810286.

Abstract

The design and validation of a reverberation chamber (RC) specifically constructed for conducting large-scale experimental animal carcinogenicity studies using RF electromagnetic fields (EMF) relevant to contemporary 4G and 5G mobile communication (900 MHz, 2.12 GHz, and 3.65 GHz) is proposed. The RC's electric field (E-field) uniformity is evaluated under four practical loading conditions: empty, apparatus only, and two apparatus variations with 80 experimental animals (Sprague–Dawley rats) with approximate weights 400 g and 520 g, respectively. Measurement results show E-field uniformity better than 1.36 dB under all test conditions, with frequency-dependent variation becoming negligible once the RC is loaded with cage racks and 80 rats. Additionally, a predictive method is introduced to estimate composite E-field intensities under simultaneous multi-frequency exposures, potentially reducing experimental measurements. These findings confirm that the designed RC is capable of accurately evaluating RF EMF exposure in biological studies.

Open access: https://www.mdpi.com/2076-3417/15/18/10286

--

Occupational exposure to extremely low-frequency magnetic fields (ELF-MF) and postmenopausal breast cancer risk

Moayedi-Nia, S, Almadin C, Labrèche F, Goldberg MS, Richardson L, Cardis E, Ho V. Occupational exposure to extremely low-frequency magnetic fields (ELF-MF) and postmenopausal breast cancer risk. Journal of Occupational and Environmental Medicine ():10.1097/JOM.0000000000003564, October 1, 2025. | DOI: 10.1097/JOM.0000000000003564.

Abstract

Objective Estimate the association between occupational exposures to extremely low-frequency magnetic fields (ELF-MF) and postmenopausal breast cancer.

Methods Lifetime job histories from a population-based case-control study (2008-2011) of histologically-confirmed breast cancer in Montréal, Canada were linked to a job-exposure matrix to assign geometric mean ELF-MF exposure/workday. Logistic regression estimated odds ratios (OR) and 95% confidence

intervals (CI) for cumulative, average, maximum, and duration of maximum exposure to ELF-MF (per interquartile range increase), adjusting for individual-level and ecological covariables.

Results Data from 663 cases and 592 controls revealed no association between occupational ELF-MF exposure and postmenopausal breast cancer, though restricting exposures to 0-10 years before interview and to those during breast development, some positive associations was observed, particularly for ER+/PR+ tumours.

Conclusions Our findings suggest no association between occupational ELF-MF exposure and postmenopausal breast cancer risk.

https://journals.lww.com/joem/abstract/9900/occupational_exposure_to_extremely_low_frequency.993.aspx

--

The effect of alpha-lipoic acid on liver damage induced by extremely low-frequency electromagnetic fields in a rat model

Aydinbelge Dizdar N, Aylin Akbulut, Koca G, Demirel MK, Yumusak N, Umurhan G, Canseven Kursun AG, Korkmaz M. The effect of alpha-lipoic acid on liver damage induced by extremely low-frequency electromagnetic fields in a rat model. Toxicol Ind Health. 2025 Sep 29:7482337251383412. doi: 10.1177/07482337251383412.

Abstract

This study aimed to investigate the effect of alpha-lipoic acid (ALA) on liver damage caused by extremely low-frequency magnetic fields (ELF-MFs) in rats. Thirty-two male rats were arbitrarily assigned into four groups: Group 1 (control group), Group 2 (ELF-MF exposure group), Group 3 (ELF-MF exposure + ALA administration), and Group 4 (sham group). Groups 2 and 3 were exposed to ELF-MF at 2 mT intensity emitted by two Helmholtz coils for 4 hours/day throughout 30 days. Group 3 received ALA intraperitoneally (100 mg/kg/day) 1 hour prior to each exposure for 30 days. Group 4 was placed inside the coils, but the generator was turned off. Histopathological analysis of the liver revealed statistically significant increases in hyperemia, inflammation, fibrosis, vacuolization, multiple nuclei, and biliary proliferation in Group 2 compared to all other groups. Immunohistochemical evaluation showed that Group 2 had statistically significantly higher TUNEL and caspase-3 levels than all other groups. ELF-MF-induced alterations, including hyperemia, inflammation, vacuolization, and multiple nuclei, were significantly reduced in Group 3 compared with Group 2. However, no significant difference was observed between Group 2 and Group 3 regarding bile duct proliferation or fibrosis. No pathological changes were observed in Groups 1 and 4. ALA administration effectively reduced some histopathological changes caused by ELF-MF, particularly hyperemia, inflammation, vacuolization, and multiple nuclei, but did not improve fibrosis or biliary proliferation. Additionally, it reduced TUNEL and caspase-3 expression. Our findings suggest that ALA exhibits anti-apoptotic and anti-inflammatory effects against ELF-MF-induced liver damage.

https://pubmed.ncbi.nlm.nih.gov/41022467/

--

Female Crabs Are More Sensitive to Environmentally Relevant Electromagnetic Fields from Submarine Power Cables

James E, Ghodsi M, Ford AT. Female Crabs Are More Sensitive to Environmentally Relevant Electromagnetic Fields from Submarine Power Cables. Environmental Science and Technology Letters. Sep 15, 2025. https://pubs.acs.org/doi/10.1021/acs.estlett.5c00862.

Abstract

The expansion of offshore wind and marine renewable energy devices (MREDs) is increasing anthropogenic electromagnetic fields (EMFs) from submarine power cables (SPCs). SPC-generated EMFs can exceed 2700 μT, well above the geomagnetic field, and may affect benthic animal behavior. In decapod crustaceans, sex-specific habitat uses and seasonal migrations are well-documented, yet their role in EMF sensitivity remains untested. We exposed juvenile shore crabs (Carcinus maenas) (n = 120; 1:1 sex ratio) to EMFs of 500, 1000, and 3200 μT using a Helmholtz coil system and tracked behavior over 10 min trials. Females exhibited strong attraction across EMF levels, spending up to 131% more time in the EMF-exposed zone and significantly less time in the low-field zone. They also drove differences in distance moved, whereas males showed no consistent spatial preference and indifferent activity at the highest field strength. These sex-specific responses suggest SPC EMFs could disrupt female-driven reproductive behaviors like seasonal migrations and larval release. Attraction may cause disorientation, aggregation, or delays in migration, potentially altering sex ratios and reducing larval export. This study provides the first evidence of sex-specific EMF responses in crustaceans and highlights the importance of incorporating sex as a key variable in ecological risk assessments of offshore infrastructure.

Open access: https://pubs.acs.org/doi/10.1021/acs.estlett.5c00862

--

A Systematic Review of the Impact of Electromagnetic Waves on Living Beings

Danho S, Escobar Huertas J, Schoellhorn WI (August 17, 2025). A Systematic Review of the Impact of Electromagnetic Waves on Living Beings. Cureus 17(8): e90355. doi:10.7759/cureus.90355.

Abstract

The effects of electromagnetic fields (EMFs) have been extensively debated among researchers and the public, with their critical consequences often dismissed or deemed unscientific. In light of this, we conducted this systematic review that extensively focuses on the detrimental effects of EMFs on living organisms.

A comprehensive and systematic literature search was performed on various electronic databases, including PubMed, Scopus, and the Cochrane Library, using Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines. This review concentrates on experimental studies published between 2017 and 2024 that investigated physiological or behavioral responses to EMF exposure, with particular attention given to those reporting harmful or concerning effects. Documented impacts include effects on humans, animals, and plants, targeting various cell types (e.g., blood, cancer,

thyroid, cochlea), genotoxicity, cardiovascular parameters (e.g., heart rate, blood pressure), male fertility (e.g., testes, sperm), neuronal brain activity, and photosynthesis in plants. Methodological quality was assessed using established bias assessment tools, and certainty of evidence was evaluated according to the GRADE framework.

After screening, 24 studies were included in the present review; five studies were non-randomized and involved humans, seven studies were in vitro, and 12 studies were conducted on animals. The findings demonstrated that EMFs negatively affect a wide array of biological systems of living organisms, including mechanisms of oxidative stress, inflammatory responses, and disruptions in cellular, physiological, and ecological processes. Most of the included studies showed a moderate to high risk of bias, which contributed to a lower overall certainty of the evidence.

These findings underscore the significant health and environmental risks associated with rising exposure levels of EMF, highlighting the urgent need for strategies to mitigate the risks. Despite these valuable insights, significant research gaps persist because the long-term effects of EMF exposure, especially on human populations, remain poorly understood and warrant further investigation and targeted mitigation strategies.

Conclusions

The rapid evolution of wireless communication technologies continues to introduce new applications and expand into higher frequency ranges. This review identifies many experimental studies reporting the biological effects of EMFs on humans, animals, and plants. These include changes in fertility parameters, cellular responses associated with oxidative stress, developmental outcomes, behavioral alterations, and cognitive effects. Our findings highlight why it is important to take a closer, structured look at how EMF exposure might affect both human health and the environment. This review focused on reported biological effects, without implying universality or disregarding studies that observed neutral or application-related outcomes. The diversity in reported results highlights the need for cautious interpretation and context-specific analysis.

Significant research gaps remain. Short-term effects have drawn much attention in past studies, but the more pressing question (what happens over time?) remains largely unanswered, especially concerning people and species highly sensitive to their environment. Additionally, a lack of standardized exposure protocols and inconsistent reporting of critical parameters such as field strength and frequency limit reproducibility and hampers the formulation of robust exposure guidelines. Addressing these challenges means committing to research that is not rushed, not vague, and not influenced by unclear funding. Long-term, transparent, and solid work is the only way forward. These should include clearly defined exposure parameters, attention to population-specific vulnerability factors (e.g., age, baseline health), and ecological relevance. As seen historically, public concern and scientific controversy can foster critical inquiry and the development of evidence-based safety standards. A coordinated effort is needed to enhance our understanding of EMF interactions with biological systems and to support informed decision-making in public health initiatives and policy.

Open access: https://www.cureus.com/articles/400923-a-systematic-review-of-the-impact-of-electromagnetic-waves-on-living-beings#!/

--

Effect of electromagnetic radiations from mobile towers on genetic damage and genetic polymorphism in humans: a review on India's perspective

Priya, K., Singh, M., Haokip, L. *et al.* Effect of electromagnetic radiations from mobile towers on genetic damage and genetic polymorphism in humans: a review on India's perspective. *Toxicol. Environ. Health Sci.* (2025). doi: 10.1007/s13530-025-00273-9.

Abstract

There have been concerns about possible health implications of radiofrequency electromagnetic radiation owing to increased cell phone use and proliferation of cell phone towers. The amount of RF-EMR energy absorbed in the human tissues relies on both the exposure characteristics of frequency, intensity, and exposure duration. This effect could be due to the exposure of genetic material by RF, resulting in an increased risk of cancer development and death in cells. Genetic polymorphisms involve variation in the DNA sequence in and among people, populations, or groups. These polymorphisms might contribute to susceptibility to RF effects. These genetic variations-which could be in the form of single nucleotide polymorphisms (SNPs) or copy number variations (CNVs-may occur either by a sheer chance process or by external factors such as viruses or radiation. Knowing the influence of genetic polymorphisms on RF exposure would help level off related health risks of cell phone use and cell towers. This scientific review examines related scientific research data about the effects that cell terminal EMR has on human genetic damage and polymorphism. We critically appraise literature and peer review journals relevant to data in assessing our current knowledge in this area. Our study reveals the fact that some studies indicate exposure to EMR does bear a relation with genetic damage, but such results should be further researched as to which health impacts they bring forward.

https://link.springer.com/article/10.1007/s13530-025-00273-9

--

Systematic reviews and meta-analyses for the WHO assessment of health effects of exposure to radiofrequency electromagnetic fields, an introduction

Verbeek J, Zeeb H, van Deventer E, Ijaz S, Doré J-F, Driessen S, Roth N, Whaley P. Systematic reviews and meta-analyses for the WHO assessment of health effects of exposure to radiofrequency electromagnetic fields, an introduction. Environment International. 2025, doi: 10.1016/j.envint.2025.109751.

No abstract

Introduction

"We, the editors (SI, JFD, SD, NR, PW) and organizers (JV, EvD, HZ) of this special issue of Environment International on the World Health Organization's (WHO) assessment of the health effects of exposure to radiofrequency (RF) electromagnetic fields (EMF), are pleased to present this collection of systematic reviews of the effects of RF EMF. This issue represents the culmination of four years of meticulous, collaborative work by more than 80 scientific experts from around the world. We extend our sincere congratulations to the authors of the reviews and commend their dedication to producing reviews of such high scientific quality.

This Special Issue includes nine protocols and subsequent twelve systematic reviews, all designed to support the human health risk assessment of RF-EMF exposure. In this editorial, we summarize the processes used to commission and develop the systematic reviews, describe the methodological tools employed, and summarise the key findings. We also reflect on the successes and challenges encountered throughout this ambitious undertaking.

Collectively, these systematic reviews offer a robust model for global health risk assessments, contributing to the development of evidence-based norms, standards, and research priorities. This project stands as one of the most comprehensive evaluations of environmental health evidence to date – an effort from which we continue to learn how to best approach this task (Jarosińska et al., 2018, Pega et al., 2021, Pérez Velasco and Jarosińska, 2022)."

Excerpts

"Although no other mechanisms beyond nerve cell excitation and thermal effect are known to pose health risks (Sheppard et al. 2008), the possibility remains that other, as yet unknown, biophysical mechanisms could exist and potentially lead to health effects."

"The last EHC Monograph on RF fields was published in 1993 (World Health Organization 1993) and is currently being updated to reflect new scientific evidence."

"The approach to developing the overall scientific basis for the updated EHC Monograph on RF Field exposure had three sequential parts: a scoping review, a prioritisation of outcomes, and the set of systematic reviews presented in this special issue."

"For one of the review topics − heat-related effects − a systematic review was not completed. The protocol did not pass peer review process in time and proceeding with it would have significantly delayed the overall project. As a result, this systematic review was decommissioned."

"The systematic reviews report evidence for the effects of RF EMF on all important health outcomes but the number of studies finally included for each outcome varied significantly, ranging from five studies on cognitive function in human observational studies (Benke et al. 2024) to 215 studies on fertility in animal experimental studies (Cordelli et al., 2023, Cordelli et al., 2024).

The systematic reviews of human observational studies on cancer found moderate-certainty evidence of no or only a small effect for several important cancer types, such as glioma and lymphoma (Fig. 1), based on 74 studies/databases (Karipidis et al., 2024, Karipidis et al., 2025). Only for thyroid cancer and oral cavity/pharynx cancer the evidence was respectively low and very low. The review of cancer in experimental animals (Mevissen et al. 2025) identified 10 long-term bioassays and dozens of other studies. Based on only one or two of these studies, the authors concluded that there was high- to moderate-certainty evidence of an effect of RF EMF on five cancer types in animals."

"The review of fertility outcomes in experimental animals yielded the most varied results. It found high-certainty evidence of no or only a small effect on litter size, but also high-certainty evidence of a large adverse effect on male fertility (Cordelli et al., 2023, Cordelli et al., 2024).(Fig. 2) Unfortunately, there were only few human observational studies on fertility that produced only very low-certainty evidence

(Johnson et al., 2024, Pw Kenny et al., 2024)."

"For cognition, human experimental studies provided consistent moderate- to high-certainty evidence of no or only a small effect across several domains of cognitive performance (Pophof et al. 2024). (Fig. 3) However, the corresponding human observational review (Benke et al. 2024) included only a few studies with results that were assessed as very low certainty."

"Regarding symptoms, human experimental studies (Bosch-Capblanch et al. 2022) showed moderate-certainty evidence of no or a small effect of RF EMF exposure. Yet again, the limited number of human observational studies resulted in very low certainty evidence (Roosli et al. 2024).(Fig. 4). "

"Finally, the review of effects of RF EMF on oxidative stress (Meyer et al. 2024) found highly variable results, ranging from large increases to large decreases in oxidative markers across all outcomes, even within one homogenous PECO. The certainty of evidence was rated very low largely due to a high risk of bias in the included studies. This was particularly evident in the exposure domain, where many experimental studies still relied on mobile phones to generate exposure – a method prone to bias if not carefully controlled (Kuster and Schönborn 2000)."

"... separate reviews were conducted for human and animal evidence for fertility and for cancer. This led to clear conclusions of the individual reviews but, as is particularly clear in cases where the animal evidence appears to contradict the human evidence, will necessitate another step of integration of the conclusions of the reviews in an overall conclusion “what this evidence means for human health”. This is the next step still to be conducted by a dedicated WHO Task Group."

"One notable exception was the systematic review on the effects of RF EMF on cancer in experimental animals, which stood out due to its use of a different synthesis method. The authors concluded that there was an effect of RF EMF if two studies showed statistically significant results, disregarding null findings from other studies. This approach deviated from the protocol, which had indicated the use of relative risk as the primary synthesis method (Mevissen et al. 2025). Despite extensive discussions with editors and peer reviewers, the review ultimately concluded that there was high-certainty evidence of an effect on cancer. Whether this method proves valid – and whether alternative synthesis approaches would yield the same conclusion—remains to be seen."

"This complex project has led to high-quality summaries of the evidence of effects of RF EMF on the most important health outcomes reported in the WHO prioritization survey in a standardized and transparent way. The results and conclusions from the present set of systematic reviews provide the most solid achievable basis for further evaluation of the effects of EMF exposure on health by the WHO task group, and the upcoming EHC Monograph of RF fields."

"Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: 1. Jos Verbeek received financial support from WHO for preparing this introductory article based on financial support from the Australian Radiation Protection and Nuclear Safety Agency, the Public Health Agency of Canada, the Environmental Protection Agency of Ireland, the Swiss Federal Office of Public Health and the Bill and Melinda Gates Foundation. The funders had no role in the decision to publish or in the preparation of the manuscript. 2. Up until the end of 2022, PW’s

contribution to the Special Issue was financially supported by the honorarium he received as an Associate Editor for Environment International. This contract ended in December 2022. From January 2023, PW's contribution was pro bono as a Guest Editor at the journal. From April 2024 to May 2025, PW provided paid contractual services to WHO on a project to develop systematic review protocols and an evidence-mapping methodology for evaluating health effects of exposure to PFAS chemicals. 3. SI's time is supported by the National Institute for Health and Care Research Applied Research Collaboration West (NIHR ARC West)."

Open access: https://www.sciencedirect.com/science/article/pii/S0160412025005021

--

Carcinogenicity of Radio-Frequency Radiation: Similarities and Differences Between Outcomes of Two Studies

Peleg M, Berry EM, Deitch M, Nativ O, Richter E. Carcinogenicity of Radio-Frequency Radiation: Similarities and Differences Between Outcomes of Two Studies. Med Discoveries. 2025; 4(7): 1267

Abstract

We discuss the similarities and the differences between two papers published on the carcinogenicity of radio frequency radiation in the military setting with some overlap of the patients' groups. The first paper reported statistically significant increases of cancer cases while the second did not. We show that the two papers presented similar and non-conflicting characteristics of the reported cancers, mainly high hematolymphoid cancers percentage frequencies and earlier onset of cancers in the exposed group. We think that the conclusion of not identifying carcinogenicity, which is explained in the second paper by not reaching statistical significance, is due to much smaller size of the patients' group in that paper.

Open access: https://www.meddiscoveries.org/pdf/1267.pdf

--

Altered development in rodent brain cells after 900 MHz radiofrequency exposure

Bodin R, Godin L, Mougin C, Lecomte A, Larrigaldie V, Feat-Vetel J, Méresse S, Montécot-Dubourg C, Marcelo P, Mortaud S, Villegier AS. Altered development in rodent brain cells after 900 MHz radiofrequency exposure. Neurotoxicology. 2025 Sep 2;111:103312. doi: 10.1016/j.neuro.2025.103312.

Abstract

Health risks related to 900 MHz 2 G frequency exposure remain inconclusive under current regulatory standards. Research into potential long-term effects is ongoing, particularly as the use of mobile networks and wireless devices increases. This study investigates the effects of non-thermal exposure levels of mobile phone 900 MHz radiofrequency electromagnetic field (RF-EMF) on rodent neurodevelopment. In vivo, the effects of pre- and post-natal 0.08 and 0.4 W/kg specific absorption rate (SAR) exposure were assessed for their impact on the proteomic profile at postnatal day 0 (PND 0). Brain-derived neurotrophic factor (BDNF), BrdU+ proliferative cells, synaptogenesis, and oxidative stress

in the hippocampus and cortex of rat pups were studied at PND 8 and PND 17. Effects of the lowest SAR (0.08 W/kg) were assessed in vitro to afford mechanistic data regarding neural stem cells (NSCs) differentiation. In vivo results showed a decrease in BDNF level and BrdU+ proliferative cells with a decrease in synapse balance (excitatory synapses/inhibitory synapses). In vitro, at 0.08 W/kg there was an increase in Ki-67 + proliferative cells, apoptosis, and double-strand DNA breaks in NSCs. A lower ratio of B1 cells (primary progenitors of NSCs) among total cerebral cells and a higher ratio of oligodendrocyte progenitor cells and astrocytes were observed in the exposed NSCs. Our findings suggest that key cellular events for brain ontogenesis are likely to undergo changes with RF-EMF 900 MHz exposure during early development. These support the hypothesis that the developing central nervous system is vulnerable to RF-EMF exposures in rodents at regulatory thresholds.

Highlights

• In vivo exposure reduced cortex cell proliferation, BDNF and synapses balance.
• In vivo exposure reduced hippocampus synapses density and balance.
• Proteomic analysis showed dysregulation of proteins involved in synaptic signaling.
• In vitro exposure increased apoptosis, DNA damage and glial differentiation shift.
• 900 MHz exposure affect rodent neurodevelopmental at regulatory threshold levels.

Open access: https://www.sciencedirect.com/science/article/pii/S0161813X2500110X?via%3Dihub

--

Oxidative stress and testicular damage induced by chronic exposure to 35.5 GHz millimeter wave radiation in male Wistar rats

Gautam R, Jha N, Tomar AK, Nirala JP, Arora T, Rajamani P. Oxidative stress and testicular damage induced by chronic exposure to 35.5 GHz millimeter wave radiation in male Wistar rats. Andrology. 2025 Aug 12. doi: 10.1111/andr.70107.

Abstract

Background: The widespread usage of millimeter waves (MMW) in different spheres of the public domain necessitates to have a better understanding of any possible health impacts caused by them. The latest example of this is the global implementation of 5G technology.

Objectives: The present investigation aimed to study the effects of 35.5 GHz frequency on reproductive parameters of male Wistar rats.

Methods: The study employed a randomized controlled design, with animals being assigned to control, sham-exposed, and exposed groups, with six animals in each group (n = 6). The experimental group was exposed to 35.5 GHz frequency for 2 h per day for 60 days. At the end of the exposure period, various sperm parameters such as sperm count, morphology, viability, and sperm mitochondrial activity were evaluated. The evaluation of oxidative stress was done by lipid peroxidation assay in testis homogenate. Superoxide dismutase, total sulfhydryl level, and total antioxidant capacity were assessed in testis homogenate. The DNA damage in the testis was assessed through the comet assay.

Results: The results showed a significant decrease in sperm viability and count, as well as morphometric changes in testis histopathology. Lipid peroxidation testis was significantly elevated in the exposed group, while superoxide dismutase (SOD), total sulfhydryl level, and total antioxidant capacity were significantly decreased in the exposed group. The comet assay revealed a significant increase in DNA damage in the exposed group, indicating genotoxic effects.

Discussion: The observed changes in sperm parameters, testicular morphology, and oxidative stress markers highlight the potential for testicular tissue damage via oxidative mechanisms. These results underscore the need for further research to elucidate the biological implications of chronic MMW exposure.

Conclusion: Chronic exposure to 35.5 GHz frequency may negatively impact male reproductive function and testicular tissue, primarily through mechanisms involving oxidative stress.

Open access: https://onlinelibrary.wiley.com/doi/abs/10.1111/andr.70107

--

Prospective long-term follow-up of patients with idiopathic environmental intolerance attributed to electromagnetic fields after a provocation trial

Li KH, Guo HR. Prospective long-term follow-up of patients with idiopathic environmental intolerance attributed to electromagnetic fields after a provocation trial. Ecotoxicol Environ Saf. 2025 Sep 1;303:118944. doi: 10.1016/j.ecoenv.2025.118944.

Abstract

Previous short-term follow-ups of idiopathic environmental intolerance attributed to electromagnetic fields (IEI-EMF) found no patients recovered after receiving provocation trials, and only a small portion were willing to consider something else caused their symptoms, although levels of their symptoms and concerns decreased. Few long-term follow-ups have been conducted, so we conducted a study to fill the data gap. We recruited participants of the initial provocation trial between 2010 and 2015 to assess changes in symptoms and concerns related to EMFs and the reversibility of IEI-EMF. The same questionnaire used in the trial was administered through telephone interviews. Of the 147 participants recruited, 70 completed the follow-up, including 35 IEI-EMF patients and 35 referents. Of the patients, 60 % were willing to consider something else caused their symptoms, and 62.9 % reported recovery from IEI-EMF after the trial in 1.80 years on average, among whom 86.4 % reported idiopathic recovery. Most of the proportions of symptoms and concerns related to EMFs among patients decreased after the trial although changes in concerns between the study groups did not reach significant differences. In conclusion, different from the previous short-term follow-ups, most patients recovered in our long-term follow-up. Most of recovered cases were idiopathic, indicating IEI-EMF is self-limited in most cases given sufficient time. Most of patients were willing to consider something else caused their symptoms, indicating most cases of IEI-EMF might be attributable to nocebo effects. Our findings suggest a possible therapeutic effect of provocation trials, which needs to be confirmed by further randomized controlled intervention studies.

Highlights

Previous studies found no IEI-EMF patients recovered after provocation trials.
We describe the first long-term follow-up of IEI-EMF after a provocation trial.
Most of the patients reported recovery, with 86.4 % being idiopathic recovery.
Most of the patients were willing to consider something else caused their symptoms.

Conclusions

Most participants with IEI-EMF had recovered in the current long-term follow-up study, while none of the participants with IEI-EMF in the two previous short-term follow-ups reported recovery, and most recovered cases in our study were idiopathic. This indicates that IEI-EMF is self-limited in most cases given sufficient time. In addition, provocation trials might accelerate the natural evolution course of IEI-EMF. A relatively large proportion, up to 60.0 %, of participants were willing to consider the possibility that their symptoms might have been caused by something other than EMFs, while the previous short-term studies observed much lower proportions. In addition, the proportion of feeling concerns that EMFs might be threatening their health and feeling worries about EMFs may affect every aspect of their life decreased in the participants with IEI-EMF. However, these did not relieve them from the worries about visualizing large EMF devices in the environment. In conclusion, our findings supported that IEI-EMF may be related to the nocebo effect in most cases. Our study also suggested a possible therapeutic effect of provocation trials when the results are negative, which needs to be confirmed by further randomized controlled trials. These findings might be applied to other types of IEI and cast some light on its etiology.

https://pubmed.ncbi.nlm.nih.gov/40897052/

--

Looking for Biomarkers Which May Explain Idiopathic Environmental Intolerance Attributed to EMF (IEI-EMF): Does RF-EMF Exposure Influence Salivary Cortisol Response?

My note: "... the current study tested predominantly young, healthy adults, the results cannot be generalized to other populations, including children, older adults, or those with poorer health." Since none of the participants were EHS, why do they consider this an IEI-EMF study?

Verrender A, Manley J, Wallace NK, Loughran SP, Croft RJ. Looking for Biomarkers Which May Explain Idiopathic Environmental Intolerance Attributed to Electromagnetic Fields (IEI-EMF): Does RF-EMF Exposure Influence Salivary Cortisol Response? Bioelectromagnetics. 2025 Sep;46(6):e70021. doi: 10.1002/bem.70021.

Abstract

In order to understand Idiopathic Environmental Intolerance attributed to Electromagnetic Fields (IEI-EMF), it has been argued that it is crucial to test for effects of radiofrequency electromagnetic fields (RF-EMF) on biomarkers, given that they can be more objective than symptom reports. While no clear evidence links RF-EMF exposure to biomarker changes, research remains limited and largely speculative due to the lack of known bioeffect mechanisms. However, there is in vitro evidence that cortisol is affected by heating, which, as RF-EMF causes heating, raises the possibility that RF-EMF exposure may

increase cortisol via thermally mediated processes. If cortisol is affected by RF-EMF exposure, it may form part of a broader aetiology of IEI-EMF, where RF-EMF-induced physiological (cortisol) inputs first generate somatic sensations, which are then fostered by expectancy or learning-based processes to generate symptoms. However, studies investigating whether RF-EMF exposure influences cortisol have reported inconsistent, but mostly null results, and many suffer from methodological issues. The current study was designed with several methodological improvements to determine whether RF-EMF affects cortisol. Seventy-two participants completed a randomized, double-blind, counterbalanced provocation study where they were exposed to both active (2 W/kg peak SAR_{10g} in head) and sham RF-EMF (0 W/kg peak SAR_{10g} in head). Despite implementing several methodological improvements, the current study failed to find an effect of RF-EMF exposure on salivary cortisol concentration. This study provides a valuable direction for future research and stresses the importance of establishing and testing theoretically plausible interactions between low-level RF-EMF exposure, the human body, and IEI-EMF symptoms.

Summary

- The current study incorporated several methodological improvements to test whether exposure to RF-EMF increases salivary cortisol concentration in humans.
- The current study failed to find an effect of RF-EMF exposure on salivary cortisol concentration, and there was no influence of participants' sex.
- Given the novelty of the methodological improvements implemented in this study, this lack of effect on salivary cortisol is very robust.

Open access: https://onlinelibrary.wiley.com/doi/10.1002/bem.70021

--

Navigating Environmental Crossroads: Pesticides, Bee Pollinators, and the Wireless Revolution

Davis D, Malkan S, Ogunseitan OA. (2025). Navigating Environmental Crossroads: Pesticides, Bee Pollinators, and the Wireless Revolution. *Environment: Science and Policy for Sustainable Development*, *67*(5), 3–4. https://doi.org/10.1080/00139157.2025.2518032.

The webinar series hosted by *Environment: Science and Policy for Sustainable Development* is open to the public and welcomes ongoing conversations about strategies for navigating environmental cross-roads. The webinar entitled “Navigating Environmental Crossroads: Pesticides, Bee Pollinators, and the Wireless Revolution” occurred on the Zoom platform on 26 March 2025 with 263 registrants and more than 80 attendees from a broad range of countries. The video recording of the webinar is embedded in the online version of this article, available here: https://doi.org/10.1080/00139157.2025.2518032

Video 1. (1:04:02)

As 2025 unfolds, we find ourselves at a critical environmental crossroads. In early January, wildfires driven by hurricane-force winds swept across Southern California, forcing hundreds of thousands to flee. Nearly 90 square miles burned where urban and wilderness areas meet—another stark signal of a changing climate.Footnote1

The fires were still smoldering on 20 January when the United States announced its withdrawal from the United Nations Paris Agreement.Footnote2 Weeks later, on 12 March, the U.S. Environmental Protection Agency launched what it called the “Biggest Deregulatory Action in U.S. History,” moving to roll back the mercury and air toxics standards, dismantle environmental justice programs, and restructure the agency’s Science Advisory Board.Footnote3

These events framed the launch of a new webinar series by Environment: Science and Policy for Sustainable Development, designed to engage a multidisciplinary audience in seeking solutions to complex challenges at the intersection of science, policy, and sustainability.

Scientific integrity is nonnegotiable. Yet translating research into regulatory policies involves negotiation among stakeholders with competing values. Our best tools for navigating environmental crossroads must maximize transparency and consider trade-offs across One HealthFootnote4—human, animal, plant, and ecosystems—as well as economics, equity, and resource conservation.Footnote5

The first webinar focused on two underrecognized global threats: the impact of neonicotinoid pesticides on bee pollinators and food security, and the growing risks of wireless radiation to all living systems. The session underscored the urgency of turning robust scientific evidence into practical policy—limiting exposures, promoting safer alternatives, and closing critical research gaps.

Neonicotinoids (neonics), the most widely used insecticides in history, have come under intense scrutiny in recent years.Footnote[6] While they may protect some crops from pests, mounting evidence suggests they are also contributing to the alarming decline in bee populations worldwide—particularly honey bees, one of over 2,000 species vital to food production.Footnote[7] The long-term consequences for pollination and ecosystem stability are profound and deeply concerning.Footnote[8]

Research on neonics’ broad-spectrum harms has provoked strong resistance from pesticide manufacturers, who often follow the tobacco-industry playbook:Footnote[9] casting doubt on science and attacking scientists. These industries fund research programs at universities and push back when government-sponsored studies, such as those from the National Toxicology Program, produce inconvenient results.Footnote[10]

Simultaneously, our world is increasingly bathed in electromagnetic radiation from mobile phones, Wi-Fi networks, and other wireless technologies. While these tools have reshaped communication and daily life, evidence continues to build around their potential health impacts, especially with long-term exposure.Footnote[11] Adverse effects have been observed not only in humans, but also in animals and bee populations.Footnote[12,]Footnote[13]

The electronics industry, now the most profitable in history, has shown little appetite for investigating or mitigating these harms. With the rise of generative artificial intelligence and global 5G expansion, exposure will only intensify. Yet industry investment in health risk research remains minimal.

A recent review for the World Health Organization examined 52 experimental studies and found high-certainty evidence linking radiofrequency radiation to tumors of the brain and heart nerves.Footnote[14] These findings, led by independent experts, add urgency to calls for applying the precautionary principle.Footnote[15]

We do not need perfect certainty to take protective action. In many cases, the science is already sufficient to justify precautionary policies. But we must also continue investing in independent research and accelerating the development of safer technologies.

For pesticides, nontoxic alternatives exist—but will not be widely adopted until manufacturers and users recognize the long-term benefits of sustainable practices. The same is true for wireless technology: We can innovate to reduce radiation exposure while maintaining connectivity. Doing so requires incentives, transparency, and public pressure.

When in doubt, we must err on the side of caution. This means restrictingFootnote[16] neonicotinoid use, especially in sensitive environments, and setting stricter guidelines for wireless radiation—particularly in schools and homes. It also means attentiveness to potential synergy between wireless radiation and chemical exposures leading to amplified adverse impacts on the productivity and stability of bee hives. These challenges cross borders, demanding international cooperation in science, regulation, and innovation.

Public education is essential. Informed citizens make better choices and demand better policies. The *Environment* webinar series aims to foster honest, inclusive dialogue about how we face today's environmental crossroads—and how we find a better path forward.

At stake is more than academic debate. The decisions we make now will shape the future of our health, our ecosystems, and our shared planet. We can find our way—but only if we insist on transparency, independent science, democratic participation, and precaution in the face of uncertainty. Without them, we risk losing the trail entirely—at a cost we cannot afford.

Disclosure Statement

Devra Davis is treasurer of the Board of Directors of Environmental Health Trust, a nonprofit organization that voluntarily discloses sources of funding more than $5,000: https://ehtrust.org/about/. Stacy Malkan is managing editor of U.S. Right to Know, a nonprofit newsroom and public health research group that also voluntarily discloses sources of funding more than $5,000 at https://usrtk.org/donors/. Oladele A. Ogunseitan is unpaid co-chair of Apple Inc.'s Green Chemistry Advisory Board.

https://www.tandfonline.com/doi/full/10.1080/00139157.2025.2518032

--

From particulates to pathways: environmental exposures and their impact on Alzheimer's disease

Liu B, Ahmad MA, Abbas G, Ahmed U, Javed R, Ali I, Ao Q, Deng X. From particulates to pathways: environmental exposures and their impact on Alzheimer's disease. Brain Res. 2025 Aug 8:149880. doi: 10.1016/j.brainres.2025.149880.

Abstract

Alzheimer's disease (AD) is a leading cause of cognitive decline and mortality, with its neurobiological mechanisms and etiology still not fully understood. Emerging evidence highlights the significant role of environmental pollutants in AD onset and progression. This review examines the impact of environmental compartments-air, water, soil, and pollutants-on AD pathology. Prolonged exposure to particulate matter ($PM_{2.5}$), heavy metals (lead, cadmium, mercury, arsenic), and engineered

nanomaterials (silver, iron oxide, silica) increases AD risk. Additional factors like obesity, smoking, infections, cardiovascular diseases, diabetes, and electromagnetic fields may exacerbate AD. These exposures potentially drive key pathological features such as amyloid-β plaque deposition and tau protein aggregation. By analyzing recent studies, this review highlights the intersection between environmental exposure and AD progression, emphasizing how such factors can accelerate the disease. It provides practical guidelines to mitigate these risks, aiming to reduce AD incidence while advancing understanding of its environmental contributions.

4.9. Electromagnetic fields

Electromagnetic fields (EMFs) are produced by various electronic devices, including those used in wireless communication, such as cell phones, Wi-Fi routers, and smart meters. EMFs are categorized into two main types: low-frequency (e.g., power lines) and high-frequency (e.g., radiofrequency radiation from wireless devices) (Meenu et al., 2024). The increasing prevalence of EMF exposure in daily life has raised concerns regarding its potential impact on brain health, particularly its role in neurodegenerative diseases such as AD (Jiang et al., 2016).

One of the central hypotheses regarding EMF exposure and AD is based on the calcium hypothesis of neurodegeneration. High-intensity EMFs are thought to interact with voltage-gated calcium channels in brain cells, which are responsible for regulating the influx of calcium ions (Ca^{2+}) into neurons. Research suggests that the increased concentration of Ca^{2+} ions resulting from EMF exposure may contribute to a cascade of pathological processes associated with neurodegeneration, including excessive calcium signaling, oxidative stress, and inflammation (Pall, 2022). The dysregulation of calcium homeostasis is a well-established mechanism in AD, where elevated intracellular calcium levels can lead to neuronal damage, synaptic dysfunction, and ultimately cell death.

Animal studies have provided compelling evidence supporting the role of EMFs exposure in the accumulation of amyloid-β, a hallmark protein involved in AD. For example, a study involving rats exposed to daily pulsed EMFs demonstrated elevated levels of amyloid-β plaques in the brain, suggesting that EMFs exposure could accelerate amyloid pathology and contribute to cognitive decline (Li et al., 2019). The deposition of amyloid-β plaques is a critical event in the pathogenesis of AD, as these plaques disrupt neuronal communication and promote inflammation, thereby exacerbating neurodegeneration. In addition to amyloid-β accumulation, neurodegeneration itself is another key consequence of EMF exposure. Human studies have shown that prolonged exposure to EMFs may lead to widespread loss of dendritic spines and synaptic connections in the brain (Pritchard et al., 2019). Dendritic spines are small protrusions on neurons that play a critical role in synaptic signaling and cognitive processes. Their loss is a key feature of neurodegenerative diseases, including AD (Wyszkowska et al., 2019). Increased calcium ion influx resulting from EMF exposure may disrupt the structural integrity of dendritic spines, leading to synaptic dysfunction and cognitive impairment (Glaser et al., 2019, Popugaeva et al., 2017, Tong et al., 2018).

In a study by Tong et al., (2018), exposure to EMFs was found to increase calcium ion levels in neuronal cells, which led to the activation of signaling pathways associated with neuroinflammation and oxidative stress. These pathways are implicated in the progressive damage to neurons seen in AD and other

neurodegenerative disorders. Additionally, chronic EMFs exposure has been shown to induce neuroinflammatory responses, which play a central role in the pathophysiology of AD. Inflammation in the brain, often triggered by the activation of microglial cells, can lead to the release of cytokines and other inflammatory mediators that further exacerbate neuronal damage and amyloid plaque accumulation (Kim et al., 2021). Another key aspect of the EMF-AD connection is the role of oxidative stress. Exposure to EMFs has been shown to increase the production of reactive oxygen species (ROS) in neurons, which can lead to oxidative damage to cellular components, including lipids, proteins, and DNA. This oxidative damage is thought to be a major contributor to the pathogenesis of Alzheimer's Disease, as it impairs neuronal function and accelerates the formation of amyloid plaques and tau tangles, both of which are characteristic of AD pathology (Bektas and Dasdag, 2025). Some studies have even suggested that individuals living in close proximity to sources of high-intensity EMF radiation, such as cell towers or power lines, may have an increased risk of developing neurodegenerative diseases, including Alzheimer's (Huss et al., 2009).

4.10. Synergistic effects of combined environmental pollutants

While much of the research on environmental risk factors for AD focuses on individual pollutants, growing evidence suggests that the combined effects of multiple exposures may significantly increase the risk of developing the disease (Wang et al., 2025). The synergistic interactions between various pollutants, such as smoking and heavy metals, can lead to more severe neuroinflammation, oxidative stress, and neuronal damage than the effects of any single exposure (Krisanova et al., 2024). For example, smoking and heavy metal exposure are two well-established risk factors for AD. Smoking leads to the generation of ROS and inflammation, which can directly damage neurons and alter amyloid-β processing (Wallin et al., 2017). When combined with exposure to heavy metals like cadmium, lead, or arsenic, the effects of smoking are often amplified. Heavy metals, by disrupting cellular homeostasis and enhancing oxidative stress, can exacerbate the damage caused by smoking. Specifically, cadmium has been shown to increase the permeability of the blood–brain barrier (BBB), allowing for easier access of amyloid-beta peptides into the brain, and promoting tau hyperphosphorylation (Ramírez-Mendoza et al., 2024). The combination of these pollutants leads to a vicious cycle of neuroinflammation, neuronal death, and cognitive decline, significantly increasing the risk of AD development in individuals who are exposed to both factors (Fu et al., 2022). The synergistic effects of combined exposures to environmental pollutants underscore the complexity of AD risk factors and highlight the need for more comprehensive studies that examine multifactorial exposures (Ahmed et al., 2025). Current research often focuses on individual risk factors in isolation, but the combined impact of pollutants is likely to provide a more accurate representation of real-world exposures (Aderinto et al., 2025b).

7. Conclusions

This monograph has underscored the critical impact of environmental risk factors stemming from industrialization, modern agriculture, and other pollutants on mental health, particularly in the context of AD. As AD research reaches a pivotal moment, the multifaceted role of environmental influences in the onset and progression of this disorder is becoming increasingly clear. Despite significant advances in understanding the pathobiology of AD, many aspects remain unresolved, particularly regarding the early causative events triggered by environmental pollutants, which continue to be elusive.

Biochemical analyses and both *in vitro* and *in vivo* studies have confirmed the role of amyloid-beta (Aβ) aggregation in initiating AD, but therapeutic strategies targeting amyloid have thus far proven ineffective. This highlights the urgent need for a deeper exploration of the mechanisms that link environmental exposures to AD onset. Additionally, the rise of NMs as emerging contributors to AD progression further emphasizes the complexity of environmental impacts on neurological health.

By addressing the limitations of current research and pursuing the outlined future directions, we can gain valuable insights into how environmental factors contribute to AD and how these risks can be mitigated. This research holds the potential to enhance our understanding of AD and guide the development of public health policies and interventions aimed at reducing the burden of this devastating disease.

https://pubmed.ncbi.nlm.nih.gov/40784621/

--

Acoustic Pressures in the Head from Pulsed Microwaves: Can They Explain the Havana Syndrome?

Foster KR, Diao Y, Chou C-K. Acoustic Pressures in the Head from Pulsed Microwaves: Can They Explain the Havana Syndrome?. *TechRxiv.* August 12, 2025. doi: 10.36227/techrxiv.175502739.93949559/v1

Abstract

The microwave auditory effect-referring to "clicks" or "buzzing" sensations that a person experiences when their head is exposed to pulsed microwave energy such as emitted by radar transmitters, has been known since the 1960s. Formerly a threshold hearing phenomenon, the effect has gained renewed interest due to the advent of high powered microwave sources in recent decades. Does it represent a potential danger to people exposed to high powered microwave pulses?

Open access: https://www.techrxiv.org/users/952786/articles/1322094-acoustic-pressures-in-the-head-from-pulsed-microwaves-can-they-explain-the-havana-syndrome

--

***Panax ginseng* Ameliorates Pituitary-ovarian Dysfunction Induced by Radiofrequency Electromagnetic Radiation from Cell Phones via Upregulation of the CREM Signaling Pathway**

Oyewopo OA, Badejogbin OC, Ajadi IO, Enye LA, Ajadi MB, Ebuwa IV, Owolabi OV, Areloegbe SE, Olaniyi KS. *Panax ginseng* Ameliorates Pituitary-ovarian Dysfunction Induced by Radiofrequency Electromagnetic Radiation from Cell Phones via Upregulation of the CREM Signaling Pathway. Curr Drug Discov Technol. 2025;22(2):e300424229527. doi: 10.2174/0115701638279386240425050818.

Abstract

Background: Panax ginseng (PG) is a plant that contains ginsenosides, which are considered adaptogens that confer cellular protection. However, the impact of PG on pituitary-ovarian dysfunction and subsequent infertility is unknown. This study investigated the hypothesis that PG would attenuate pituitary-ovarian dysfunction associated with mobile phone's Radiofrequency Electromagnetic Radiation (RF-EMR) in experimental rat models and the possible involvement of a cAMP Response Element Modulator (CREM)-dependent pathway.

Methods: Twenty adult female Wistar rats were divided randomly into four groups, each consisting of five rats. The control group was administered a vehicle (distilled water) orally, while the P. ginseng group received 200 mg/kg of P. ginseng extract orally. The RF-EMR group was exposed to 900MHz radiation, and the RF-EMR + PG group was exposed to the same radiation while also being treated with 200 mg/kg of P. ginseng orally. These treatments were administered daily for a period of 56 days.

Results: The RF-EMR group exhibited significant reductions in serum levels of LH, FSH, estradiol, and progesterone compared to the control group. Moreover, levels of superoxide dismutase (SOD) and glutathione peroxidase (GPx) were significantly lower in the RF-EMR group compared to the control. Additionally, there was a notable decrease in the expression of the CREM gene, accompanied by disrupted pituitary/ovarian morphology in the RF-EMR group compared to the control. However, the administration of PG mitigated these changes.

Conclusion: The findings of this study indicate that P. ginseng extract shields against pituitary-ovarian impairment linked to RF-EMR exposure from cell phones by boosting antioxidant capacity and promoting the CREM-dependent pathway.

https://pubmed.ncbi.nlm.nih.gov/38693729/

--

Impact of in vitro exposure to 5G-modulated 3.5 GHz fields on oxidative stress and DNA repair in skin cells

Haidar J, Nabos P, Orlacchio R*, et al.* Impact of in vitro exposure to 5G-modulated 3.5 GHz fields on oxidative stress and DNA repair in skin cells. *Sci Rep* **15**, 31214 (2025). https://doi.org/10.1038/s41598-025-15090-w.

Abstract

The rapid deployment of fifth-generation (5G) wireless networks has raised societal concerns regarding potential biological effects, particularly on human skin, due to the use of higher carrier frequencies that penetrate tissue less deeply. Consequently, whether 5G-modulated radiofrequency (RF) electromagnetic fields (EMFs) at 3.5 GHz affect oxidative stress and DNA repair in skin cells remains an open question. Using genetically encoded Bioluminescence Resonance Energy Transfer (BRET)-based biosensors targeted to the cytoplasm and mitochondria, we assessed whether exposure of human fibroblasts to 5G RF-EMF at specific absorption rates (SAR) of 0.08 and 4 W/kg for 24 h could alter basal reactive oxygen species (ROS) levels or potentiate the effects of known ROS inducers, including H_2O_2, Kp372-1, and Antimycin A. We also evaluated whether pre-exposure to 5G RF-EMF could induce an adaptive response (AR), by modulating ROS production following a subsequent challenge with arsenic trioxide (As_2O_3). Additionally, we investigated the impact of combined RF-EMF and ultraviolet-B (UV-B) exposure on the formation and repair of cyclobutane pyrimidine dimer (CPD) lesions in HaCaT keratinocytes. Our results showed no significant effect of 5G RF-EMF exposure, either alone or in combination with chemical ROS inducers, on oxidative stress markers in either compartment. Likewise, RF-EMF exposure did not induce an adaptive response to oxidative challenge, nor did it alter the kinetics or the efficiency of CPD repair by the nucleotide excision repair (NER) pathway. These findings support the conclusion that the

exposure to 5G RF-EMF at 3.5 GHz up to 4 W/kg does not induce oxidative stress or impair DNA repair efficiency in human skin cells, within the experimental conditions tested.

Excerpt

In summary, our investigation—combining novel biosensing technology with robust experimental design—demonstrates that 5G-modulated 3.5 GHz RF-EMF does not affect oxidative stress levels, trigger adaptive response, or interfere with DNA repair processes in human skin cells at steady temperature. These results are consistent with current assessments by international experts and provide meaningful data for an evidence-based risk assessment of RF-EMF exposure in the context of 5G deployment. Nevertheless, it is important to note that our findings are based on acute in vitro exposure protocols lasting up to 24–48 h. While such durations are commonly used to assess direct cellular responses, they do not fully recapitulate the chronic or long-term exposures experienced in real-life settings. Therefore, future studies will be necessary to confirm these findings in more physiologically relevant models, including long-term or repeated exposures in 3D reconstructed skin or in vivo systems

Open access: https://www.nature.com/articles/s41598-025-15090-w

--

Dual Evaluation and Spatial Analysis of RF-EMF Exposure in 5G: Theoretical Extrapolations and Direct Measurements

Najera A, Villaescusa-Tebar A, Gonzalez-Rubio J, Garcia-Pardo C. Najera A, Villaescusa-Tebar A, Gonzalez-Rubio J, Garcia-Pardo C. Dual Evaluation and Spatial Analysis of RF-EMF Exposure in 5G: Theoretical Extrapolations and Direct Measurements. Bioelectromagnetics. 2025 Sep;46(6):e70020. doi: 10.1002/bem.70020.

Abstract

This study evaluates radiofrequency electromagnetic field (RF-EMF) exposure in 5G networks using a dual approach that combines theoretical extrapolations and direct measurements in diverse semiurban and urban environments, specifically on the campus of the Polytechnic University of Valencia. Measurements were conducted using personal exposimeters under active traffic conditions on the 5G network, complemented by a code-selective measurement system based on an R&S TSME6 scanner. This approach enabled the calculation of maximum theoretical exposure by detailed analysis of 5G signals and the capture of key parameters such as cell ID and beam indices at 16 representative points across the campus. For precise spatial visualization, Kriging interpolation techniques in ArcGIS were employed to generate continuous exposure maps illustrating the spatial distribution of RF-EMF in the study area. The results indicate that both the extrapolated theoretical values and measured levels align with the limits recommended by the ICNIRP, even under high data demand scenarios, supporting current safety assessments of 5G infrastructure regarding electromagnetic exposure. The correlation between theoretical and instantaneous exposures validates the applied methodology and its effectiveness in assessing exposure in diverse environments. This study provides a robust framework for future research and highlights the importance of continuous monitoring to ensure public safety during the deployment of new telecommunications infrastructure in urban areas.

Open access: https://onlinelibrary.wiley.com/doi/10.1002/bem.70020

--

Assessment of 5G RF-EMF Exposure during Large-Scale Public Events via Field Measurements

Villaescusa-Tebar A, Garcia-Pardo C. Assessment of 5G RF-EMF Exposure during Large-Scale Public Events via Field Measurements. IEEE Access, 2025. doi: 10.1109/ACCESS.2025.3602263.

Abstract

The development of Fifth-Generation (5G) mobile networks has raised public concern regarding human exposure to radiofrequency electromagnetic fields (RF-EMF). Key 5G features such as higher operating frequencies, broader bandwidths, beamforming, and traffic-dependent signal variability significantly influence RF-EMF exposure dynamics. Although many studies have assessed EMF levels induced by specific base stations under controlled user traffic, little empirical data are available on population-wide 5G environmental exposure across diverse scenarios. This study addresses this gap through field measurements conducted during a large public festival in Valencia, Spain and a baseline day prior to the event. Downlink signals from the 700 MHz and 3500 MHz 5G bands were measured across Spain's three main network operators. Five locations surrounding the event venue were selected to capture the influence of the measurement location, and two measurement durations (6 and 30 min) were compared to assess temporal dynamics. The results showed that during the event, the power density in the 3500 MHz band increased by up to a factor of eight compared to the baseline conditions. However, even the highest recorded exposure levels remained well below the international safety limits. Additionally, a 30-minute continuous measurement revealed an approximately 30% increase in average power density compared to a 6-minute snapshot. These findings confirm that 5G environmental RF-EMF exposure remains within safe limits, even under high-demand conditions, and underscore the relevance of considering measurement location and temporal variability in exposure assessments.

Open access:

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=11135488&isnumber=6514899

--

5G RF EMF Spectral Exposure Assessment in Four European Countries

Deprez K, Stroobandt B, Veludo AF, Vecsei Z, Necz PP, Politański P, Verloock L, Polanska K, Thuróczy G, Röösli M, Plets D, Joseph W. 5G RF EMF Spectral Exposure Assessment in Four European Countries. Bioelectromagnetics. 2025 Sep;46(6):e70019. doi: 10.1002/bem.70019.

Abstract

This study assesses the exposure to 5G radio frequency electromagnetic fields (RF EMF) across four European countries. Spot measurements were conducted indoor and outdoor in both public spaces and educational institutions, encompassing urban and rural environments. In total, 146 measurements were performed in 2023, divided over Belgium (47), Switzerland (38), Hungary (30) and Poland (31). At 34.9% of all measurement locations a 5G connection to 3.6 GHz was established. The average cumulative incident power density (S_{avg}) and maximum cumulative incident power density (S_{max}) were determined, for both "background" exposure (no 5G user equipment; No UE) and worst-case exposure (maximum downlink with 5G user equipment; Max DL). Furthermore, 3.6 GHz 5G-specific average $S_{avg,5G}$ and

maximum $S_{max,5G}$ incident power density are considered as well. For the No UE scenario, the highest S_{max} is 17.6 mW/m^2, while for the Max DL, the highest S_{max} is 23.3 mW/m^2. Both values are well within the ICNIRP guidelines. The highest $S_{max,5G}$ measured over all countries and scenarios was 10.4 mW/m^2, which is 3.2% of the frequency-specific ICNIRP guidelines. Additionally, a comparison was made between big cities, secondary cities, and villages for all four countries. The ratio of power density measured in rural areas was significantly lower than in urban areas (−4.8 to −10.4 dB). Under LOS conditions, the average incident power density was 2.3 mW/m^2, whereas under NLOS conditions, the average incident power density decreases to 0.9 mW/m^2. Furthermore, the relative variation increases under NLOS scenarios. Lastly, an analysis was performed regarding the power density in educational institutions compared to all other measurement locations, both indoors and outdoors for the different city types. The measured incident power density is not extensively lower in or around schools compared to public places, neither in the big cities, secondary cities, or the villages.

Summary

- Radio frequency electromagnetic fields exposure assessment measurements across four European countries are conducted using a standardized measurement method.
- All measurements were well within the ICNIRP guidelines.
- The incident power density measured in rural areas was significantly lower than in urban areas.

Conclusion

The findings of this study provide a comparison of downlink RF EMF exposure from base stations for mobile telecommunications, presented in terms of spectral power density, across Belgium, Switzerland, Hungary, and Poland. First, a comparison was made of the RF EMF exposure regulations of all countries. While all countries follow similar RF EMF exposure regulations, Belgium and Switzerland have additional precautionary limits. In total, 146 measurement locations were assessed in 2023, divided over Belgium (47), Switzerland (38), Hungary (30) and Poland (31). 34.9% of all measurement locations had 5G coverage at 3.6 GHz, although there was a distinction between urban and rural areas in available 3.6 GHz 5G base stations. The study included both average and maximum cumulative incident power densities for two scenarios: “background” (No UE) and “worst-case” (Max DL) exposure. Furthermore, 3.6 GHz 5G-specific average and maximum incident power density ($S_{avg,5G}$ and $S_{max,5G}$) were included as well. The highest maximum cumulative power density for the No UE scenario was 17.6 mW/m^2. For the Max DL scenario, the highest maximum cumulative power density was 23.3 mW/m^2. These cumulative power densities, and by extension all measurements, remain well below 1, indicating compliance with ICNIRP guidelines. The highest 3.6 GHz 5G power density measured over all countries and scenarios was 10.4 mW/m^2, which is 0.1% of the frequency-specific ICNIRP guidelines. The ratio of power density measured in rural areas was significantly lower than in urban areas (−4.8 to −10.4 dB), for all countries over the two scenarios, as expected due to less nearby base stations and generally less users.

A comparison between LOS and NLOS measurement locations is made. Under LOS conditions, the average incident power density was 2.3 mW/m^2, whereas under NLOS conditions, the average incident power density decreases to 0.9 mW/m^2. Furthermore, the relative variation increases under NLOS scenarios. Lastly, an analysis was performed regarding the power density in educational institutions compared to all other measurement locations, both indoors and outdoors for the different city types.

The measured incident power density is not extensively lower in or around schools compared to public places, in neither the big cities, secondary cities or the villages. The RF EMF exposure levels in all measured environments were well below the ICNIRP guidelines.

For future work, increasing the number of measurements and regularly reassessing exposure levels will be essential to monitor the continued rollout of 5G networks and related exposure trends. Therefore, these measurements will be repeated in 2025. Furthermore, the different measurement methods within GOLIAT will be combined to achieve an even better understanding regarding RF EMF exposure.

Open access: https://onlinelibrary.wiley.com/doi/10.1002/bem.70019

--

Assessment of spatial-average absorbed power density and peak temperature rise in skin model under localized electromagnetic exposure

Zheng J, Zhang Y, Diao Y, Shi D. Assessment of spatial-average absorbed power density and peak temperature rise in skin model under localized electromagnetic exposure. Radiat Prot Dosimetry. 2025 Aug 16:ncaf096. doi: 10.1093/rpd/ncaf096.

Abstract

Numerical dosimetry for assessments of the absorbed power density (APD) and temperature rise has been conducted using multi-layer skin models, incorporating skin, fat, muscle, and other components, providing a scientific foundation for setting exposure limits. However, the influence of the vasculature on dosimetry outcomes remains underexplored. In this study, we developed a synthetic blood vessel model and integrated it into multi-layer skin models. Electromagnetic computations were performed, followed by steady-state temperature rise evaluations using the Pennes bioheat transfer equation across a frequency range of 3 to 30 GHz. To quantify the effect of vascular modeling on dosimetry results, simulations incorporating vasculature with varying endpoint diameters were compared to those without vasculature. Results showed that the effect of vascular modeling on peak spatial-averaged APD was negligible, and its influence on peak temperature rise was ~8% at 3 GHz, decreasing to less than <3% above 6 GHz. And the effect of the endpoint diameter is marginal. These variations are smaller than those previously reported due to changes in tissue thickness and dielectric or thermal properties. While the effect on peak temperature rise is modest, including vasculature helps refine localized thermal distributions and may inform future improvements in anatomical modeling.

https://pubmed.ncbi.nlm.nih.gov/40817877/

--

Traceable Assessment of the Absorbed Power Density of Body Mounted Devices at Frequencies Above 10 GHz

Chitnis N, Karimi F, Kühn S, Fallahi A, Christ A, Kuster N. Traceable Assessment of the Absorbed Power Density of Body Mounted Devices at Frequencies Above 10 GHz. Bioelectromagnetics. 2025 Sep;46(6):e70018. doi: 10.1002/bem.70018.

Abstract

In this study, a comprehensive approach for the experimental assessment of the absorbed power density (APD) is developed. The method includes several novel components: (i) a specialized probe, (ii) a composite phantom, (iii) a reconstruction technique, (iv) a calibration method, and (v) a validation process. The described solution has been developed for the frequency range from 24 to 30 GHz, but can be extended to all frequency bands between 10 and 45 GHz. A novel composite phantom emulates the reflection and transmission coefficients of human skin for propagating and evanescent modes, while its increased penetration depth, in comparison to dermis tissue, enables the measurement of the induced electromagnetic fields (EMFs) with a new miniaturized dosimetric broadband probe. The implementation has a wide dynamic range and sufficient spatial resolution to use it for type approval of mobile devices. Its probe is calibrated with low uncertainty in a novel, traceable setup. A set of reference antennas with known numerical target values for the APD has been compiled to validate the measurement system. The validation demonstrates that the deviation is within the expanded uncertainty of 1.6 dB for pAPD and 1.5 dB for psAPD.

Summary

- First traceable, broadband absorbed power density (APD) assessment method and procedures covering 10–45 GHz with a validated implementation for 24–30 GHz, having low uncertainty of < 1.6 dB for peak APD and < 1.5 dB for peak spatial-averaged APD (psAPD), enabling type approval of wireless devices.
- Composite phantom design that emulates reflection and APD of skin across propagating and evanescent modes.
- Reference-antenna-based validation that is instrumentation agnostic and covers realistic spatial gradients, spectral domain (*k*-space), and modulated signals in conformance with standard requirements for product test system qualification.

Open access: https://onlinelibrary.wiley.com/doi/10.1002/bem.70018

--

Assessing exposure from different vehicular antennas in military applications: a computational study

Micol C, Biscarini M, Ferrante D, Pellegrino G, De Meis M, Mei L, Cavagnaro M, Apollonio F, Liberti M. Assessing exposure from different vehicular antennas in military applications: a computational study. Frontiers in Public Health. Volume 13 - 2025. doi: 10.3389/fpubh.2025.1620240.

Abstract

Introduction Military personnel frequently operate in close proximity to electromagnetic (EM) sources such as vehicular communication antennas. Despite this occurrence, detailed evaluations of exposure scenarios remain limited. This study bridges this gap by examining EM exposure from military vehicular antennas, covering a broad spectrum of frequencies (high frequency—HF, very high frequency—VHF, ultrahigh frequency—UHF), power levels, and positions.

Methods The study used computational modeling to simulate realistic military scenarios, including personnel partially outside armored vehicles and equipped with personal protective equipment.

Simulations accounted for a broad spectrum of frequencies (HF, VHF, and UHF) as well as different power levels, antenna types and locations.

Results The analysis revealed substantial variability in EM exposure levels depending on the configuration and conditions. While all simulated scenarios complied with the ICNIRP Basic Restrictions (BR), certain cases exceeded the Reference Levels (RL), particularly under specific positioning and frequency combinations.

Conclusions These findings, based on a detailed case-specific analysis, suggest that personnel safety in military contexts is generally maintained, even in the presence of variable exposure conditions and elevated levels of radiated E-field. Considering the basic structure of the radiating source (i.e., monopole) and the nature of near-field interactions, this paper suggests that safe exposure conditions could be expected to persist across a range of antenna-operator positioning configurations, and building on previous preliminary research on this topic, it provides relevant insights for operational instructions and improving safety regulations in the military field.

https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1620240

--

In situ electric field dosimetry analysis for powerline frequency peripheral nerve magnetic stimulation

Fresnel E, Bouisset N, Soyka F, *et al.* In situ electric field dosimetry analysis for powerline frequency peripheral nerve magnetic stimulation. *Sci Rep* **15**, 29887 (2025). https://doi.org/10.1038/s41598-025-12284-0.

Abstract

Humans are exposed to environmental 60 Hz magnetic fields (MFs), inducing in our body electric fields (EFs) and currents, potentially stimulating the peripheral nervous system (PNS). Uncertainties exist regarding the 60 Hz MF PNS stimulation threshold. The spatially extended nonlinear node model (SENN) is used to help define international MF exposure guidelines and standards protecting workers and the general public. However, other models exist, particularly the McIntyre–Richardson–Grill (MRG) model, the new gold standard for electrostimulation. This study aims (1) to model a new extremely low frequency MF exposure system for the human leg and (2) to investigate the in situ EFs generated by the system at 60 Hz at the skin level and in the nerves of the leg using a realistic human body model with both the SENN and the MRG models. A Helmholtz like-coil system was designed to generate in situ EFs sufficient for nerve stimulation, modeled using Biot–Savart and Faraday laws. Sim4Life simulations assessed the induced EFs at skin and nerve levels using a detailed human body model and two nerve excitation frameworks: the SENN and MRG models. High EF intensities were observed in four sensory and sensory-motor nerves, with MRG-derived thresholds lower than SENN-derived thresholds. Results also highlight the significance of nerve orientation in EF induction. This study emphasizes the critical role of comprehensive modeling for the design and validation of MF exposure systems and underscores the need for experimental data to refine models, standards, and guidelines.

Open access: https://www.nature.com/articles/s41598-025-12284-0

--

Instruments and Measurement Techniques to Assess Extremely Low-Frequency Electromagnetic Fields

Rathebe PC, Kholopo M. Instruments and Measurement Techniques to Assess Extremely Low-Frequency Electromagnetic Fields. Sensors (Basel). 2025 Aug 7;25(15):4866. doi: 10.3390/s25154866.

Abstract

This study presents a comprehensive evaluation and selection framework for extremely low-frequency electromagnetic field (ELF-EMF) measurement instruments. Recognizing the diversity of application environments and technical constraints, the framework addresses the challenges of selecting appropriate tools for specific scenarios. It integrates a structured, quantitative approach through a weighted scoring matrix that evaluates instrumentation across six criteria: monitoring duration, sensitivity, environmental adaptability, biological/regulatory relevance, usability, and cost. Complementing this is a logic-based flowchart that visually guides decision-making based on user-defined operational needs. The framework is applied to a realistic occupational case study, demonstrating its effectiveness in producing evidence-based, scenario-sensitive instrument recommendations. This method provides stakeholders with a transparent and adaptable tool for ELF-EMF device selection.

Open access: https://www.mdpi.com/1424-8220/25/15/4866

--

Weak anthropogenic electric fields affect honeybee foraging

Mallinson VJ. Woodburn FA, O'Reilly KJ. Weak anthropogenic electric fields affect honeybee foraging. Cell Press. 2025. https://doi.org/10.1016/j.isci.2025.112550.

Highlights

• Weak anthropogenic-like electric fields reduce honeybee floral landing

• AC and positive DC fields deter honeybee landing, but negative DC does not

• Electric fields near power lines match experimental levels, spanning tens of meters

• These findings highlight electric pollution as a potential disruptor of pollination

Summary

Aerial electroreception, the detection of airborne electric fields (E-fields), is an emerging sensory system in arthropods, including bees, which can use floral E-fields as foraging cues. However, the influence of anthropogenic E-fields on these interactions remains underexplored. Through field experiments in urban meadows, we demonstrate that weak anthropogenic E-fields, including alternating current (AC) and direct current (DC) fields, significantly alter honeybee floral landing behaviors. AC and positive DC fields reduced landings by 71% and 53%, respectively, whereas negative DC fields had no statistically significant impact. Measurements of E-fields near high-voltage transmission lines revealed persistent field strengths comparable to those used experimentally, spanning tens of meters at heights relevant for bee foraging. These findings underscore the potential for anthropogenic E-fields to affect and potentially disrupt plant-pollinator interactions, threatening pollination efficiency, a cornerstone of agriculture and

biodiversity. Our study highlights the need for advancing research on the ecological impacts of electric pollution.

Open access paper: https://www.cell.com/iscience/fulltext/S2589-0042(25)00811-9

--

Combined effects of constant temperature and radio frequency exposure on Aedes mosquito development

Dom NC, Dapari R, Halim NMHNA, Rahman ATA. Combined effects of constant temperature and radio frequency exposure on Aedes mosquito development. Sci Rep. 2025 Aug 20;15(1):30571. doi: 10.1038/s41598-025-09383-3.

Abstract

Mosquito-borne diseases, such as dengue, Zika, and chikungunya, pose significant public health threats, particularly in tropical regions like Malaysia. Aedes aegypti and Aedes albopictus are primary vectors of these diseases, with their developmental stages being highly sensitive to environmental factors. While temperature is a well-known driver of mosquito biology, the potential influence of anthropogenic factors such as radio frequency (RF) exposure remains underexplored. This study investigates the combined effects of temperature and RF exposure on the developmental stages of these mosquito species to provide insights into their population dynamics and inform vector control strategies. A factorial experimental design was employed, incorporating four temperature conditions (20 °C, 25 °C, 30 °C, and 35 °C) and three RF exposure levels (900 MHz, 18 GHz, and a control group with no RF exposure). The developmental durations for hatching, larval, pupation, and adult emergence stages were monitored daily under controlled laboratory conditions. Data were analyzed using a quadratic response surface model to evaluate the main effects and interactions between temperature and RF exposure. Temperature emerged as the dominant factor influencing developmental durations, with optimal conditions observed at 30-32 °C. RF exposure, particularly at 18 GHz, acted as a secondary modulating factor, accelerating developmental stages under certain temperature conditions. Ae. aegypti exhibited greater sensitivity to temperature changes compared to Ae. albopictus, which displayed higher adaptability and resilience to environmental variations. Interaction effects were most evident at intermediate temperatures (25-30 °C), where RF exposure synergistically reduced developmental durations. However, extreme RF exposure levels and suboptimal temperatures prolonged developmental periods. This study highlights the critical role of temperature in mosquito development while identifying RF exposure as a potential modulator under specific conditions. The findings underscore the importance of considering both environmental and anthropogenic factors in vector management strategies. Future research should explore the molecular mechanisms underlying these interactions to refine predictive models and enhance vector control efforts in rapidly urbanizing regions.

Conclusion

In summary, temperature is the primary determinant of mosquito developmental durations, with RF exposure exerting secondary modulating effects under specific conditions. *Ae. aegypti* was more sensitive to environmental variations, while *Ae. albopictus* displayed greater resilience and adaptability. These findings provide a foundation for incorporating environmental variables, including anthropogenic factors such as RF exposure, into predictive models for mosquito population dynamics and vector

management. Future research should explore the molecular mechanisms underlying these interactions and assess their implications for disease transmission and control in different ecological settings.

Open access: https://www.nature.com/articles/s41598-025-09383-3

--

Effect of Static Electromagnetic Field on Growth Parameters, Survival Rate, Sex Distribution, Ratio, and Liver and Gonadal Health of Zebrafish

Çek Ş, Ünal E, Doğan EA, Yaraş KU, Dede K, Karaaslan M. Effect of Electromagnetic Field on Growth Parameters, Survival Rate, Sex Distribution, Ratio, and Liver and Gonadal Health of Zebrafish (Danio rerio). Bull Environ Contam Toxicol. 2025 Aug 25;115(3):33. doi: 10.1007/s00128-025-04103-z. PMID: 40855214.

Abstract

This study examines the impact of electromagnetic fields (EMF) on zebrafish (Danio rerio) sex ratio, survival, and organ health. Embryos were exposed to static EMFs for 63 days post-hatching across five aquariums, with varying intensities based on distance (30-99 cm). The control group remained EMF-free. The strongest exposure (30 cm) led to 100% feminization ($p < 0.001$), with all exposed groups showing significant deviations from the expected 1:1 sex ratio. Survival rates dropped below 35% in EMF-exposed groups, compared to 82.5% in controls. Histological analysis revealed gonadal abnormalities, necrosis, and liver degeneration in exposed fish. These findings provide the first documented evidence that EMF exposure disrupts sex ratios and health in fish, potentially affecting population stability in aquatic ecosystems. The predominance of females suggests long-term ecological consequences that could impact biodiversity in both aquatic and terrestrial environments.

https://pubmed.ncbi.nlm.nih.gov/40855214/

--

Exposure of human lymphocytes to sweeping-frequency extremely low frequency magnetic field

Zastko L, Makinistian L, Petrovičová P, Tvarožná A, Belyaev I. Exposure of human lymphocytes to sweeping-frequency ELF MF. Int J Radiat Biol. 2025 Aug 7:1-10. doi: 10.1080/09553002.2025.2542322.

Abstract

Purpose Whereas most of the research on possible bioeffects of extremely low frequency magnetic fields (ELF MF) on blood cells has been performed at 50/60 Hz regardless of background fields, here, we exposed human lymphocytes to sweeping-frequency ELF MF in a different range, defined by the DC background field of our incubator and the corresponding ion-cyclotron frequencies.

Materials and methods Umbilical cord blood lymphocytes (UCBL) were isolated and exposed for 48 h to an ELF MF (sinusoidal, frequency sweeping 3–26 Hz) with 6 different amplitudes between 6 µT and 24 µT, utilizing an oblong coil. DNA double-strand breaks (DSB) were assessed by enumeration of γH2AX, 53BP1 and γH2AX/53BP1 co-localized DNA repair foci. Percentage of viable, early apoptotic (EA), and late apoptotic/necrotic (LAN) cells were determined.

Results No statistically significant effects were seen for DNA repair foci or apoptosis induction after a 48-h exposure of UCBL with frequency-sweeping ELF MF. More than 2-fold decrease (p = .064) of γH2AX foci level was detected for 8 μT amplitude when compared to controls. For the same field intensity, a decrease in viable cells was suggested by the data.

Conclusions Our findings suggest that at least part of the exposure could have had a preventive/blocking effect on DNA DSB formation suggesting the possibility of using sweeping-frequency ELF MF as a protective measure against genotoxic agents. It is desirable to continue testing variations of the exposure utilized in this work, in order to search for the most biologically effective frequencies/patterns of exposure.

https://pubmed.ncbi.nlm.nih.gov/40772784/

Magnetic effects in biology: Crucial role of quantum coherence in the radical pair mechanism

Binhi V. Magnetic effects in biology: Crucial role of quantum coherence in the radical pair mechanism. Phys. Rev. E. **112**, 014409. 25 July 2025**.** doi: 1103/n3fs-fsnv.

Abstract

The spin-chemical radical pair mechanism (RPM) has emerged as a leading theory for explaining the biological effects of low-intensity magnetic fields. These intriguing effects occur when the quantum system of radicals is well isolated from the disturbing influence of the environment. In other words, these effects are closely related to the spin coherence relaxation time τ, but an explicit relationship has not yet been established. In our study, we found an analytical solution to the Liouville-Neumann equation for an open system made up of two electrons and one nucleus, considering minimal interactions while concentrating on spin relaxation and chemical kinetics. This solution, supported by numerical integration, highlights the crucial role of quantum coherence. A straightforward expression is proposed that describes the RPM effect as a function of τ, within the ranges of magnetic field strength H and rate κ of chemical kinetics relevant to magnetobiology. Our findings reveal that RPM effects become significant only when fundamental relation $\gamma H\tau>1+\kappa\tau$ holds: it controls the magnitude of the effects, and it is consistent with the principles of spin chemistry. Additionally, by comparing our results with existing experimental data, we estimate that the plausible spin decoherence times in magnetosensitive radical pairs within cryptochromelike proteins range from units to tens of nanoseconds. The effects of radio-frequency magnetic fields at the nT level were also examined, taking into account decoherence. These effects turned out to be negligible and incapable of disrupting the RPM patterns. The role of the quantum Zeno effect in magnetobiology is inspected from the perspective of the τ dependence of the RPM effect.

https://journals.aps.org/pre/abstract/10.1103/n3fs-fsnv

--

Radiofrequency Electromagnetic Field Emissions and Neurodevelopmental Outcomes in Infants: A Prospective Cohort Study

Setia M S, Natesan R, Samant P, et al. (July 10, 2025) Radiofrequency Electromagnetic Field Emissions and Neurodevelopmental Outcomes in Infants: A Prospective Cohort Study. Cureus 17(7): e87671. doi:10.7759/cureus.87671.

Abstract

Purpose It has been argued that children are particularly at risk of developing health effects due to the emitted radiofrequency electromagnetic fields (RF-EMF). We designed this cohort to measure the association between exposure to RF-EMF radiation and neurodevelopmental changes in neonates and infants.

Methods We present an analysis of 261 observations from a cohort of 105 neonates. The cohort was formed of pregnant women, and the neonates born to these women were followed for a period of one year. We assessed the level of radiation in the house using the Selective Radiation Meter 3006 (Narda Worldwide, Germany) and neurodevelopment using the Ages and Stages Questionnaire® (ASQ®)-3. We used random effects models for multiple observations in the same individual, and the main explanatory variable was household radiation levels (divided into tertiles as low/medium/high).

Results The median (interquartile range (IQR)) range of radiation in all the households was 8.66 (IQR: 1.58, 23.11) mW/m^2. It was 0.62 (IQR: 0.43, 1.58) mW/m^2 in the lowest tertile, 8.66 (IQR: 5.00, 10.78) mW/m^2 in the middle tertile, and 32.36 (IQR: 23.11, 45.60) mW/m^2 in the highest tertile (p=0.0001). The mean scores were significantly lower in the middle and higher tertiles of LOR for the gross motor, fine motor, and problem-solving domains. The odds of children classified as 'monitor/refer' was significantly higher in the 'high radiation group' compared with 'low radiation group' for the fine motor (adjusted odds ratio (aOR): 2.74, 95% CI: 1.10, 6.78; p=0.03) and problem-solving domains (aOR: 3.67, 95% CI: 1.41, 9.55; p=0.008). We also found that low birth weight babies were significantly more likely to be classified as 'monitor/refer' for fine motor (aOR: 4.19, 95% CI: 1.73, 10.14; p=0.001), and problem solving (aOR: 2.57, 95% CI: 1.08, 6.13; p=0.033) domains.

Conclusions Even after adjusting for low birth weight, we found that higher levels of radiation were associated with poorer outcomes for cognitive domains of development such as problem solving, and personal-social areas. Thus, there is a need to monitor the neuro-development of children in whom the RF-EMF radiations are expected to be higher (such as very close to cell phone towers, too many gadgets in the house).

Excerpts

"In this cohort study, we found that mean ASQ scores were, in general, lower in the highest radiation tertile for all five domains; and specifically, significantly lower for the gross motor, fine motor, and problem-solving domains. Poor development outcomes (such as monitor/refer) for the fine motor, problem-solving, and personal-social domains were associated with higher levels of radiation in the house. In addition, low birth weight was also significantly associated with poor development outcomes for fine motor and problem-solving. Finally, even though mean scores for the social-emotional domain were high in the higher radiation groups, the difference was not statistically significant."

"There are potential limitations in the present analysis. In this study, the focus was on the association between the levels of electromagnetic field radiation and neurodevelopmental outcomes in neonates and infants. Other studies have included prenatal cell phone use and exposure as the exposure variable; we did not include these in the present analysis. We also did not assess the parent-child interaction as another potential confounder. The EMF radiation in the house may not only be due to that from cell phone towers but also due to cordless phones, WiFi devices, and Bluetooth gadgets. Since we measured the radiation in the house, this would have included the EMF from all these. It is quite likely that a home far away from the tower may still have higher EMF radiation due to these gadgets. However, the measurement was only at baseline, and this may have led to some misclassification. As described earlier, some authors have not found an association between cell phone use, while others have found inconsistencies in the relationship. Most other authors have used other scales, such as the Bayley Scales of Infant Development or scales that were developed by the researcher. We have used the Ages and Stages Questionnaire, both for the developmental domain and the social-emotional domain. ASQ is useful for screening and not a diagnostic instrument. We also do not claim causality in these findings. These are interim results of the cohort, and we are still following the cohort. We intend to publish future results on the neurodevelopmental outcomes. These outcomes will include the Bayley Scales of Infant Development and the Stanford-Binet test for intelligence. We did not use any imputation methods for missing observations and just used the random effects models with available observations. Though we had enough power to detect the difference in scores, some of the events were few (particularly social-emotional and communication domains). Thus, we may be underpowered at this point of analysis for these outcomes. Nonetheless, actual measurement of RF-EMF radiation in the house (which represents the actual exposure levels), a cohort design, and the use of random effects logistic regression models are the potential strengths of the study. These models are useful for longitudinal data, and one additional advantage of random effects models is their use in longitudinal data when observations are present at different time points [42,43].

Conclusions

These preliminary findings are an important contribution to the literature on the association between RF-EMF radiation and neurodevelopment in neonates and infants. The results have to be interpreted taking into account the limitations that have been mentioned. We used random effects models, which account for both within-subject and between-subject correlation, and are useful for time-varying variables. These models are useful for longitudinal data where the outcomes may vary with each observation. Even after adjusting for low birth weight, we found that higher levels of radiation were associated with poorer outcomes for cognitive domains of development such as the problem-solving and personal-social areas. Low birth weight was associated with poorer outcomes for the motor development domains (gross motor and fine motor). It is quite likely that obvious gross motor delays may be identified by parents and caregivers; however, cognitive development domains require specialist monitoring. Thus, there may be a need to consider monitoring of neurodevelopmental outcomes in children in whom RF-EMF radiations are expected to be higher (such as very close to cell phone towers and too many gadgets in the house)."

Open access: https://www.cureus.com/articles/381425-radiofrequency-electromagnetic-field-emissions-and-neurodevelopmental-outcomes-in-infants-a-prospective-cohort-study#!/

--

A scoping review and evidence map of radiofrequency field exposure and genotoxicity: assessing in vivo, in vitro, and epidemiological data

Weller SG, McCredden JE, Leach V, Chu C, Lam AK-Y (2025) A scoping review and evidence map of radiofrequency field exposure and genotoxicity: assessing in vivo, in vitro, and epidemiological data. Front. Public Health 13:1613353. doi: 10.3389/fpubh.2025.1613353

Abstract

Background Studies investigating genotoxic effects of radiofrequency electromagnetic field (RF-EMF) exposure (3 kHz−300 GHz) have used a wide variety of parameters, and results have been inconsistent. A systematic mapping of existing research is necessary to identify emerging patterns and to inform future research and policy.

Methods Evidence mapping was conducted using guidance from the Preferred Reporting Items for Systematic reviews and Meta-Analyses for Scoping Reviews (PRISMA-ScR). A comprehensive search strategy was applied across multiple research databases, using specific inclusion and exclusion criteria within each knowledge domain. Quantitative aggregation using tables, graphs and heat maps was used to synthesize data according to study type, organism type, exposure level and duration, biological markers (genotoxicity, cellular stress, apoptosis), RF-EMF signal characteristics, as well as funding source to further contextualize the evidence landscape. Quality criteria were applied as part of a focused analysis to explore potential biases and their effects on outcomes.

Results Over 500 pertinent studies were identified, categorized as in vitro (53%), in vivo (37%), and epidemiological (10%), and grouped according to type of DNA damage, organism, intensity, duration, signal characteristics, biological markers and funding source. In vitro studies predominantly showed proportionally fewer significant effects, while in vivo and epidemiological studies showed more. DNA base damage studies showed the highest proportion of effects, as did studies using GSM talk-mode, pulsed signals and real-world devices. A complex relationship was identified between exposure intensity and duration, with duration emerging as a critical determinant of outcomes. A complex U-shaped dose-response relationship was evident, suggesting adaptive cellular responses, with increased free radical production as a plausible mechanism. Higher-quality studies showed fewer significant effects; however, the funding source had a stronger influence on outcomes than study quality. Over half (58%) of studies observing DNA damage used exposures below the International Commission of Non-Ionizing Radiation Protection (ICNIRP) limits.

Conclusion The collective evidence reveals that RF-EMF exposures may be genotoxic and could pose a cancer risk. Exposure duration and real-world signals are the most important factors influencing genotoxicity, warranting further focused research. To address potential genotoxic risks, these findings support the adoption of precautionary measures alongside existing thermal-based exposure guidelines.

Implications for policy and practice

The evidence from the evidence map indicates that medium to long-term RF-EMF exposures, particularly at low intensities, can induce genetic damage through non-thermal mechanisms such as increased free radical production and oxidative stress. Genetic damage can have far-reaching, long-term, and potentially irreversible consequences for individual organisms and broader ecological and planetary health (112, 113).

Both In vivo and epidemiological RF-EMF studies provide credible evidence of genotoxicity, suggesting potential risks such as increased cancer susceptibility and reproductive harm. Studies on brain cells frequently reported positive findings for DNA damage, suggesting that brain cells may be particularly sensitive to RF-EMF, indicating a risk for neurological diseases and brain tumors, as observed in animal models (114–116).

Current RF-EMF exposure guidelines established by ICNIRP (15), prioritize the prevention of thermal effects by incorporating substantial safety margins (e.g., a 50-fold reduction from effect thresholds, setting a local SAR limit of 2 W/kg for the head and torso for the general public, averaged over 10 grams of tissue). However, the evidence mapping process found statistically significant DNA damage at extremely low intensities, with the lowest recorded effects occurring at a SAR of 0.000000319 W/kg in an epidemiological study (117) and at 0.000003 W/kg in several in vivo experiments (118, 119). These levels are substantially (>600,000 times) below the ICNIRP public exposure limits (15). This pattern suggests non-thermal genotoxic effects, because temperature changes at these intensities would be negligible and not measurable.

ICNIRP (2020) guidelines (15) set RF-EMF exposure limits to protect against thermal effects from acute exposures, with averaging times of 6 min for local exposure (head and torso) and 30 min for whole-body exposure. However, the above analysis revealed that medium (1 day−3 months) and long-term RF-EMF exposures (>3 months or 1,000 h) were most strongly linked to genotoxic effects, even at very low exposure intensities. ICNIRP (2020) guidelines (15) do not set specific limits for chronic, low-level RF-EMF exposures, particularly for non-thermal effects like genotoxicity, citing “no substantiated evidence of health-relevant effects” [(15), p. 522].

The mapping process also revealed that RF exposures are associated with genetic damage in a wide range of organisms, with an observed sensitivity of non-mammalian organisms, such as plants, insects, and possibly amphibians. Current guidelines neglect potential effects on wildlife or ecosystems (78, 96). Notably, a recent WHO-commissioned systematic review of animal studies suggested carcinogenic effects from RF-EMF exposures (116). Other studies suggest biological effects on non-human species (120, 121). Together, these results suggest that the environmental implications of RF-EMF exposure merit greater scrutiny (122), even though the current evidence remains limited and debated (96).

While these findings do not yet establish causation or a clear No Observed Adverse Effect Level (NOAEL), they indicate risks that ICNIRP's current framework discounts by prioritizing only effects with confirmed harm [(15), p. 487]. ICNIRP's review process and position is best described as a hazard-based assessment focused only on confirmed effects. This approach is overly restrictive, as it delays updating guidelines until absolute certainty is achieved (123), which may not align with the precautionary needs of public health or environmental protection.

Currently, there is a widespread (6) and often non-consensual nature to RF-EMF exposure (92) from mobile phones, base stations, and other wireless technologies. While acknowledging the permanence of this technology in modern society, policy adjustments are required that prioritize health and environmental protection over economic interests. This can be achieved by adopting a precautionary

approach to RF-EMF (123) and addressing potential risks from non-thermal RF-EMF effects, despite scientific uncertainty. Strategies such as justification (assessing net benefits of RF-EMF applications), optimization (balancing protection with societal needs), and As low as Reasonably Achievable or As Low as Technically Achievable - ALARA/ALATA (avoiding deterministic effects and minimizing stochastic effects) per International Commission on Radiological Protection (ICRP) recommendations - ICRP103 (124) could be considered. Further development and deployment of wireless technologies should incorporate improved safety measures in their design (125), such as creating devices that emit lower levels of RF-EMF or using materials and antenna designs to direct emissions away from the body.

Additionally, public information regarding potential health risks and personal protective measures could be disseminated through public health campaigns, making use of existing advice such as the EUROPAEM EMF Guideline 2016 (126); e.g. minimizing the use of wireless devices, prioritizing wired connections, maintaining distance between RF-EMF sources and the body, use of air-tube headsets or handsfree calls, turning off wireless when not in use, and mitigation of oxidative stress by incorporating antioxidants into the diet.

While individual actions are valuable, they are not a substitute for robust regulatory standards and industry accountability. Ensuring the safety of wireless technologies requires a collective effort from manufacturers, policymakers, and consumers to develop comprehensive RF safety guidelines. Future regulatory guidelines could encompass workplace protection measures, including substitution, engineering, and administrative controls (127), integration of building biology standards (128), mandatory detailed product labeling to inform users of potential risks, and standardized safety hygiene practices.

Recommended actions

To address these concerns and bridge existing gaps, the following actions are recommended:

- Standardization of Research Protocols: Harmonizing methodologies across studies, particularly comet assay protocols, is critical for reducing heterogeneity and enabling robust meta-analyses.
- Focus on Long-Term and Low-Intensity Exposures: Future research should prioritize investigating the cumulative effects of prolonged and low-intensity RF-EMF exposures, which are most relevant to real-world scenarios and devices.
- Inclusion of Emerging Frequencies: Given the rapid deployment of 5G and other novel technologies, research focused on higher frequencies and new modulation schemes is urgently needed.
- Targeted Environmental and Health Studies: Targeted research in both human health and ecological systems needs to be conducted independently of vested interest influences, ensuring methodological rigor in each domain.
- Independent Funding and Research Oversight: To mitigate biases associated with industry funding, greater support for independent research is essential. Transparent disclosure of ALL funding sources and researcher affiliations should be mandatory.

- Re-evaluation of RF Standards: Regulatory bodies must update exposure guidelines to reflect non-thermal mechanisms and the potential health effects from long-term chronic exposure settings by incorporating findings from independent, high-quality studies.

Conclusions

The evidence map presented here reveals statistically significant DNA damage in humans and animals resulting from man-made RF-EMF exposures, particularly DNA base damage and DNA strand breaks. The evidence also suggests plausible mechanistic pathways for DNA damage, most notably through increased free radical production and oxidative stress. Sensitivity to damage varied by cell type, with reproductive cells (testicular, sperm and ovarian) along with brain cells appearing particularly vulnerable. A complex U-shaped dose-response relationship was observed for both exposure duration and intensity, with more DNA damage occurring in specific frequency and intensity combination windows. DNA damage was more likely to be found using *in vivo* studies, very weak or very strong signal intensities, very short or very long exposure durations, 900, 1,800 and 2,450 MHz frequencies, GSM-talk mode and pulsed modulations, particularly when using real-world devices. On the other hand, research funded by vested interests has tended to use different experimental design parameters, with a high proportion of studies using *in vitro*, short-term exposures, medium-high intensity signals and using signal generators. Funding source is also a stronger determinant of experimental outcomes than study quality.

Overall, there is a strong evidence base showing DNA damage and potential biological mechanisms operating at intensity levels much lower than the ICNIRP recommended exposure limits. Public policy could benefit from the implementation of precautionary measures such as ALARA or ALATA, along with public information campaigns to better safeguard human and environmental health and wellbeing.

Open access: https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1613353

--

Building the gulf of opinions on the health and biological effects of electromagnetic radiation

Paul Héroux. Building the gulf of opinions on the health and biological effects of electromagnetic radiation. Front. Public Health, 22 July 2025. Volume 13 – 2025. xoi: 10.3389/fpubh.2025.1589021.

Using events that the author was personally involved with over many years, the article attempts to explain how different views solidified over time on the health effects of electromagnetic radiation, some believing they are negligible, while others believe they are substantial.

Introduction

The health impacts of Non-Thermal Electromagnetic Radiation, both in the ELF and RF domains, have been controversial since the early 1980s (1). The report of a link between childhood leukemia and ELF magnetic fields fueled discussion for more than a decade, ushering together two very different areas, electrical engineering and biology.

The arguments oppose the officers of industry and their followers to health environmentalists. Given

that both should have access to the same scientific literature, the opposing positions rely on the selection of different experts, subsets of the literature, and on their interpretation.

This article attempts to explain how different views on the health effects of technological electromagnetic radiation solidified over time, some believing they are negligible, while others propose they are substantial.

We skirt around classically reported events such as the development of the ANSI (21), IEEE (22) and ICNIRP (23) recommendations as well as the Bioinitiative report (24), the ORSAA (25) database, Henry Lai's literature compilation (26), and the NTP (27) and Ramazzini (28) experiments to discuss other incidents that may seem minor but shed some light on the more human aspects of the formation of opinions about EMR health impacts.

To resolve the complex problem of EMR health impacts, several scientific meetings were held worldwide, seemingly to advance science and develop opinions.

We choose to report events where, over the years, the author was *physically* present; such direct experiences easily solidify into valuable memories. We think that assembling such recollections can be used to explain the development of diverging opinions on EMR health impacts.

Section headings

Adair, 1991

Armstrong, 1994

Lai, 1997

Phillips, 2009

Epilogue

The four examples above illustrate different aspects of the divergence of opinions. How are the basic terms of reference in the debate interpreted…are AC and DC fields in some ways equivalent (Adair)? Does the Specific Absorption Rate, a heat variable, provide more clarity than electric and magnetic fields? Should the engineering view of the problem smother its biological aspects? Can humans truly be simulated as a salt and sugar solution, as is done in SAR measurements? What of the issues of not pursuing challenging leads (Armstrong), pressuring the messengers of unwelcome observations (Lai) and undermining or underrating the techniques of biology (Phillips)?

In the recounts above, a leitmotiv is a divergent interpretation of the same facts according to personal education and employment. But control of the collective agenda in the public sphere is equally important, participation at scientific meetings by one group versus another can be strongly biased.

The issue of EMR health impacts was viewed as critical to industry interests at certain times, while for the bio-medical community it may have appeared less urgent.

So, was the industry justified in taking control of the debate and limiting it to heat, given that they had acute interests in the outcomes?

Was industry justified in publicizing its thermalist agenda through the IEEE with so many governments worldwide?

Was industry justified in controlling the evolution of EMR research, using its expertise in electromagnetism to dominate the area, while the most critical elements resided in biology and medicine?

The focus on ionization and thermal limits guaranteed peaceful outcomes for industry. The success of microwave ovens apparently crystallized a view of the human body as a dipole rather than a conductor, but this view may not be everlasting (20).

Open access paper: https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1589021/full

--

Impact of Radiofrequency Electromagnetic Fields on Cardiac Activity at Rest: A Systematic Review of Healthy Human Studies

Michelant L, Selmaoui B. Impact of Radiofrequency Electromagnetic Fields on Cardiac Activity at Rest: A Systematic Review of Healthy Human Studies. Bioelectromagnetics. 2025 Jul;46(5):e70014. doi: 10.1002/bem.70014.

Abstract

Radiofrequency electromagnetic fields (RF-EMF) exposure is increasingly prevalent, raising concerns about potential non-thermal health effects. This systematic review synthesizes current evidence regarding RF exposure effects on cardiac activity, focusing on heart rate (HR) and heart rate variability (HRV). Studies on healthy individuals were selected based on strict methodological criteria, including experimental design, control for confounding variables, and sufficient details on exposure parameters. Articles were included if they compared healthy subjects with and without exposure and provided cardiac measurements, specific absorption rate, or exposure measurement. A total of 28 articles were analyzed. This review included studies with RF exposure ranging from 100 to 110,000 MHz and exposure durations varying from short periods to 7 nights, with most studies lasting between 5 and 50 min. Most studies demonstrated no significant effects of RF exposure on HR, regardless of the exposure system, frequency, duration, age, sex, distance, or subject position. Findings for HRV were more nuanced, with most studies indicating no significant impact on key HRV parameters. However, some position-dependent variations emerged, particularly in antenna-based studies. Additionally, our analysis suggests that RF exposure may particularly interfere with cardiac regulatory mechanisms when the cardiovascular system is challenged and required to adapt, such as during postural changes or physiological maneuvers, although there are insufficient comparable studies to validate this hypothesis. Importantly, all included studies were conducted under resting or non-stressful conditions and involved only healthy participants. Therefore, our conclusions cannot be generalized to stressed states or clinical populations. Moreover, methodological harmonization is needed to improve comparability across future studies. The main

limitation of the current evidence being the heterogeneity of experimental protocols, highlighting the need for methodological standardization in future studies. To address current heterogeneity, we propose specific methodological recommendations, including systematic blinding, accurate exposure measurement and detailed exposure, to improve comparability and reproducibility in future studies.

Summary

- Radiofrequency exposure under ICNIRP guidelines has no significant effect on heart rate across various exposure conditions at rest and for healthy subjects.
- Findings related to heart rate variability are more nuanced; while most studies show no significant impact in calm conditions and for healthy subjects, some position-dependent variations have been observed, particularly in antenna-based studies.
- RF exposure may modulate cardiac response during its physiological regulation, such as in the Valsalva maneuver or orthostatic test, and affect heart rate and variability in these cases, although limited studies prevent definitive conclusions.
- Methodological standardization is necessary for future research.

Open access: https://onlinelibrary.wiley.com/doi/10.1002/bem.70014

--

Men with genetic predisposition face greater fertility challenges when exposed to electromagnetic radiation

Pal S, Paladhi P, Dutta S, Ghosh P, Chattopadhyay R, Ghosh S. Men with genetic predisposition face greater fertility challenges when exposed to electromagnetic radiation. Mol Biol Rep. 2025 Jul 31;52(1):773. doi: 10.1007/s11033-025-10882-9.

Abstract

Purpose: This study investigates the synergistic effects of genetic variants in core meiotic regulator genes-SPO11, RNF212, and SYCP3-and co-occurring exposure to electronic radiation as risk factors for azoospermia among Bengali-speaking men from West Bengal, India.

Materials and methods: A total of 708 azoospermic individuals, who tested negative for Y chromosome microdeletions, and 640 healthy controls were genotyped using Sanger's dideoxy sequencing. Genetic variants identified included SPO11 rs3764674T > G, MN650122delA, MN708963insA, MN708964insA, MN720359T > A; RNF212 MN737491delA, rs4045481C > T; and SYCP3 rs10860779C > A, MN901901delA, and MN995822delA. Binary logistic regression was used to assess the interaction between genetic variants, electronic radiation exposure, and age, with fertility status as the outcome variable.

Results: The analysis revealed a significant association between genetic variations in meiotic regulators and increased risk of azoospermia, particularly among men aged 30 years or older who were exposed to electronic radiation. The findings suggest that effect of electronic radiation may exacerbate meiotic errors, impair germ cell development, and further reduce fertility in affected individuals.

Conclusions: Men carrying genetic variations in SPO11, RNF212, and SYCP3 and exposed to electronic radiation are at an elevated risk of azoospermia, particularly with advancing age. This information may be considered in routine screening of assisted reproductive technology (ART) practices, which will enable clinicians to tailor management strategies for male infertility.

https://pubmed.ncbi.nlm.nih.gov/40742570/

--

The modeling of the interaction of pulsed 5G/6G signals and the fine structure of human skin

Betzalel N, Feldman Y, Ben Ishai P. The modeling of the interaction of pulsed 5G/6G signals and the fine structure of human skin. *Sci Rep* **15**, 28651 (2025). https://doi.org/10.1038/s41598-025-13777-8.

Abstract

Current regulations concerning the allowed levels of human exposure to electromagnetic radiation emanating from wireless technologies are governed by the Specific Absorption Rate standard (SAR). This allows the absorption by tissue of up to 2 W/kg averaged for 6 minutes in a 10 g cube of homogenized tissue. Much criticized, the SAR standard relates only to thermal effects. However, with the advent of 5G and 6G technologies, exploiting frequencies above 4 GHz, the traditional methods of SAR measurement are proving to be inadequate. This is made more poignant as carrier wavelengths approach the dimensions of tissue structures. We present a detailed electromagnetic simulation of human skin that not only accounts for the multi-layered structure of skin, but also sweat ducts, capillaric and arterial blood vessels. The results show an inhomogeneous absorption pattern that reflects the vessels and sweat glands involved. As human skin hosts an array of sensing structures, from nociceptors to thermoreceptors, as well as millions of innervated sweat glands, we point out that the current methods for gauging the SAR rating of devices are wholly inadequate and may lead to a gross underestimation skin electromagnetic absorption, and that nerve excitation should be accounted for in risk assessment.

External electromagnetic source

To emulate realistic source signals as used in real wireless devices nowadays, we have created a sinusoidal signal with carrier frequency of each frequency of our interest (3.5 GHz, 27 GHz, 77 GHz and 300 GHz). We have multiplied the sinusoidal wave with *rect* function with 30% duty cycle, and created a periodical train pulsed signal as described in 6.

$$y(f,t) = sin(2\pi f_c t) rect(2\pi f_i t) \quad (6)$$

Where f_c is the carrier frequency, f_i is the information frequency and t is the time vector.

Figure 12 shows the 77 GHz rectangular pulsed signal. The information frequency for that specific band in 1 GHz.

Conclusions

In this research we have demonstrated the EM vulnerability of skin components such as sweat ducts (its coiled section and sweat glands, in particular) and dermal blood vessels, which are ignored by standard regulation approval methods of wireless devices. We conclude that wireless and remote EM devices and applications, such as cellular phones, laptops, autonomous vehicles, smart cities etc. Working in the 5G bands and planned to be work in 6G bands, are not being treated correctly in terms of EM affects they might generate on humans. In our study, we see the vulnerability of the sweat gland and the coiled section of the gland, possessing almost an EM filter component, which filters in 5G and 6G frequencies (3.5, 27, 77 and 300 GHz) into the blood stream. Moreover, the peripheral area of the sweat gland also absorbs more electromagnetic energy (higher dosimetry) than its surrounding. 5G and 6G might pose unexpected risks to humans, due to the effect of energy dissipation to regulators, such as thermoreceptors or nociceptors, in the immediate vicinity of the sweat duct, with long- or short-term effects. We conclude that since human skin is a relatively complex organ, it cannot be treated as a homogeneous component when it interacts with millimeter waves and sub-THz frequencies.

Open access: https://www.nature.com/articles/s41598-025-13777-8

--

Exploring research trends in health effects of 5G antennas: a bibliometric analysis

Turgut A, Korunur Engiz B. (2025). Exploring research trends in health effects of 5G antennas: a bibliometric analysis. *Electromagnetic Biology and Medicine*, 1–20. doi: 10.1080/15368378.2025.2539147.

Abstract

To better understand the developments and trends of studies on the health impacts of 5th Generation (5 G) antennas, a bibliometric analysis of the literature published in the Web of Science database from 2012 to 2025 was conducted. Before the analysis, the dosimetric quantities “specific absorption rate (SAR)” and “power density (Sab)” used in the examination of the health impacts of antennas were thoroughly defined, and their required limits were provided. Subsequently, information visualization technology was utilized to investigate the yearly distribution of literature, author contributions and collaboration, productive and influential institutions and countries/territories, co-citation analysis, and keyword co-occurrence. However, anticipated publication and citation numbers for the coming years (2025–2029) were predicted regarding the potential health impacts of 5 G antennas using artificial intelligence-assisted forecasting methods. There has been a considerable increase in studies related to health impacts measured by SAR and Sab dosimetric quantities over the last five years, in line with the number of articles published on 5 G antennas. Despite this, no bibliometric analysis has been done so far on the health impact of 5 G antennas below and above 6 GHz. The data presented in this bibliometric study to fill the gap on this subject will provide a better understanding of the health impacts and unique insights to offer good research guidance on 5 G antennas and their attributed adverse health impacts.

Plain Language Summary

This study explores research conducted on potential health effects related to 5G antennas. 5G is the latest wireless communication technology designed to increase internet speeds and connect more devices simultaneously. As 5G technology expands, antennas operating at different frequencies are being widely installed, raising public concerns about their safety and possible health impacts.

Our analysis covered studies from 2012 to 2025, examining how researchers measure human exposure to electromagnetic fields from these antennas, specifically focusing on two measures: Specific Absorption Rate (SAR) for lower frequency antennas and Power Density (Sab) for higher frequency antennas. SAR indicates the rate at which electromagnetic energy is absorbed by body tissues per unit mass, while Sab represents the absorbed power density at the body's surface.

Using advanced methods, we analyzed the growing body of research, highlighting key trends and major contributors in the field. We found a rapid increase in studies, driven by growing public interest and technological advances. Most research currently emphasizes smartphone antennas, but interest in wearable devices and health-monitoring technologies is rising.

Our findings emphasize the need for ongoing research into 5G's health implications, providing guidance for future studies. The study helps policymakers, researchers, and the public understand the safety considerations surrounding 5G technology and encourages continued attention to health-related issues as the use of 5G continues to grow.

Table 4. Top 7 institutions with the most publications on the health impacts of 5 g antennas.

Rank	Institution	Country	TP	P	TLS	CPP	H
1	Ericsson	Sweden	17	4.381%	138	34.35	10
2	Sony Corporation	Japan	14	3.608%	98	31.14	8
3	University Teknikal Malaysia Melaka	Malaysia	13	3.351%	24	7.62	7
4	National Institute Of Technology (NIT) System	India	12	3.093%	2	4.58	4
5	Polytechnic University Of Milan	Italy	12	3.093%	28	7.08	6
6	Zhejiang University	China	12	3.093%	99	18.17	6
7	Consiglio Nazionale Delle Ricerche (CNR)	Italy	11	2.835%	13	7.18	5

TP: Total Publications; P: Percentage (%); TLS: Total Link Strength; CPP: Citations Per Paper; H: h-index.

Conclusion

This study presents a comprehensive bibliometric analysis of research on 5 G antennas and their health impacts, mapping knowledge domains through visualization techniques. The findings indicate that as the deployment of 5 G networks expands, research on SAR (Specific Absorption Rate) and Power Density (Sab) will continue to grow, particularly with the increasing demand for human exposure assessments and safety regulations.

The AI-assisted prediction models applied in this study project a steady increase in research output and citations over the next three years, highlighting the importance of bioelectromagnetic safety and multidisciplinary collaborations in fields such as medicine, chemistry, and physics. This trend suggests a growing scientific and regulatory focus on the potential biological effects of 5 G radiation.

While bibliometric approaches provide valuable insights, they also present certain limitations. The accuracy and completeness of the dataset depend on database indexing and data retrieval methods. Additionally, preprocessing techniques for author and institutional name variations require further refinement to minimize errors. The study also highlights challenges in data visualization, particularly in overlapping clusters and network mapping, which may require advanced machine learning techniques for optimization.

Despite these limitations, the study offers a solid foundation for understanding the evolution of 5 G antenna research, particularly in the context of health risk assessments and SAR compliance studies. The findings emphasize the need for continued interdisciplinary collaboration to ensure that future wireless communication technologies are both efficient and safe for public health.

These findings are further supported by AI-assisted forecasting, which projects stable research output through 2029, driven by ongoing interest in SAR and Sab-based exposure analysis. Current developments – such as increased focus on wearable antennas, improved EMF simulation methods, and regulatory initiatives – are expected to shape future research directions. The results also highlight the need for continued interdisciplinary collaboration and refinement of data analysis techniques to better assess potential health effects of 5 G technologies.

https://www.tandfonline.com/doi/full/10.1080/15368378.2025.2539147

--

Comparative Analysis of Beamforming Techniques and Beam Management in 5G Communication Systems

Andras CM, Barb G, Otesteanu M. Comparative Analysis of Beamforming Techniques and Beam Management in 5G Communication Systems. *Sensors*. 2025; 25(15):4619. doi: 10.3390/s25154619.

Abstract

The advance of 5G technology marks a significant evolution in wireless communications, characterized by ultra-high data rates, low latency, and massive connectivity across varied areas. A fundamental enabler of these capabilities is represented by beamforming, an advanced signal processing technique that focuses radio energy to a specific user equipment (UE), thereby enhancing signal quality—crucial for maximizing spectral efficiency. The work presents a classification of beamforming techniques, categorized according to the implementation within 5G New Radio (NR) architectures. Furthermore, the paper investigates beam management (BM) procedures, which are essential Layer 1 and Layer 2 mechanisms responsible for the dynamic configuration, monitoring, and maintenance of optimal beam pair links between gNodeBs and UEs. The article emphasizes the spectral spectrogram of Synchronization Signal Blocks (SSBs) generated under various deployment scenarios, illustrating how parameters such as subcarrier spacing (SCS), frequency band, and the number of SSBs influence the spectral occupancy and synchronization performance. These insights provide a technical foundation for optimizing initial access and beam tracking in high-frequency 5G deployments, particularly within Frequency Range (FR2). Additionally, the versatility of 5G's time-frequency structure is demonstrated by the spectrogram analysis of SSBs in a variety of deployment scenarios. These results provide insight into how different configurations affect the synchronization signals' temporal and spectral occupancy, which directly affects initial access, cell identification, and energy efficiency.

Open access: https://www.mdpi.com/1424-8220/25/15/4619

--

Numerical Analysis of Human Head Exposure to Electromagnetic Radiation Due to 5G Mobile Phones

Chietera FP, Usmani WU, Caggiano D, Lamacchia C, Chimenti G, Mescia L. Numerical Analysis of Human Head Exposure to Electromagnetic Radiation Due to 5G Mobile Phones. 2025 10th International Conference on Smart and Sustainable Technologies (SpliTech), Bol and Split, Croatia, 2025, pp. 1-5, doi: 10.23919/SpliTech65624.2025.11091791.

Abstract

This paper investigates human exposure to electromagnetic fields from 5G mobile phones operating in the 26 GHz frequency band, recently designated for 5G mm-wave communications. Numerical simulations assess near-field exposure generated by a realistic mobile phone model and the thermal effects on a detailed head model, providing valuable insights into the potential impact of this emerging technology on human health.

Conclusion

A preliminary analysis conducted using a realistic head model with medium-resolution voxels suggests that 5G mm-wave communication in the 26 GHz band results in only moderate, localized thermal increases in superficial tissues. However, further investigations are necessary to better characterize the impact of prolonged exposure and higher spatial detail. Future simulations should consider higher-resolution voxel models, more refined tissue segmentation, and extended exposure durations. In addition, variable phone placements should be explored, as they can lead to higher incidence angles and higher peak irradiation levels in biological tissues.

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=11091791&isnumber=11091382

--

Pilot questionnaire survey shows the lack of diagnostic criteria for electromagnetic hypersensitivity: a viewpoint

Leszczynski D. Pilot questionnaire survey shows the lack of diagnostic criteria for electromagnetic hypersensitivity: a viewpoint. MHealth. vol 11. Jul 14, 2025. doi: 10.21037/mhealth-25-4.

Abstract

Wireless communication devices and networks are currently prevalent in human environment. Some persons claim to be sensitive to emitted by them microwave radiation. Commonly, this sensitivity is called electromagnetic hypersensitivity (EHS) or microwave disease. However, because of the yet scientifically unproven link between radiation exposures and EHS symptoms, this sensitivity is also called idiopathic environmental intolerance attributed to electromagnetic fields (IEI-EMF). The sensitivity is not recognized by the World Health Organization as a disease or as being caused by exposures to wireless radiation. There are no medical tests for detecting sensitivity to wireless radiation. Physicians are not being educated to deal with persons who claim to be sensitive to wireless radiation. However, some persons who consider themselves to be sensitive to wireless radiation exposures claim to have medical

diagnoses made by physicians or other health professionals. This project looked at the contradiction of the lack of diagnostic criteria for sensitivity to wireless radiation with the medical diagnoses claimed by some of the self-declared sensitive persons. Analysis of questionnaire responses of 142 self-declared sensitive persons suggests that, currently, it is not possible to diagnose sensitivity to wireless radiation exposures. The claimed medical diagnoses appear to be based on the anecdotal evidence presented by the self-declared sensitive persons. In some cases, medical tests were used but these tests lacked scientific proof of their ability to detect the sensitivity of a person to wireless radiation exposure. The proof of the existence of sensitivity to wireless radiation remains inadequate. However, logically and by analogy to other environmental stressors, it is likely that individual sensitivity to wireless radiation exists. Because provocation studies in wireless radiation-exposed volunteers alone seem unable to provide definite answers, further research using both, provocation and biochemical methods with controlled wireless radiation exposures in volunteers is necessary to discover diagnostic biomarkers of EHS.

Table 1: Symptoms claimed to be caused by exposures to wireless radiation

• Skin problems: itching; eczema; getting acne; bad skin rash on the face; face gets hot; burning feeling on neck and thyroid area; creeping skin, like something is walking on it; feeling a tinkling on the forehead; red stripes on the skin of hands; skin of hands becoming very dry, splits and bleeds
• Nervousness feelings: exhaustion and fatigue; insomnia; feeling agitated; cardiac arrhythmia; pulsations in chest and heart; anxiousness; over activity; overstimulation; shaking or trembling; panic attacks; fear feeling; psychosis; suicidal thoughts; vertigo—the sensation that the environment around is moving or spinning; buzzing from the bottom of the feet; nervous tension throughout the body; pulsations in arms; restlessness; restless leg syndrome; tingling in fingers and toes; vibration feeling in the head, body, and pelvic area
• Digestive tract problems: belching; diarrhea; digestion problems; increased food sensitivities painful constipation; food intolerance to e.g., cow's milk, eggs yeast, wheat, or gluten
• Brain functioning problems: brain fog; memory problems; problems with concentration; difficulty speaking/stuttering; disorientation; distorted sense of time and space; dizziness; drowsiness; nausea
• Effects on moving, seeing, hearing, breathing: difficulty walking; fuzzy vision; blurred vision; shimmering visual effects; easily lost balance; uncoordinated movements; tinnitus; hearing loss; hearing persistent hum; respiratory depression; shortness of breath; coughOpen access: https://mhealth.amegroups.org/article/view/140469/html

Table 3: Some values of the blood tests of the self-declared EHS persons were elevated and some lower than the average levels

• Elevated: Cu, glucose, IgE, histamine, orotic acid, 3-methyl-glutaric, bacterial metabolites, fungal metabolites, glutaric acid, hippuric acid, oxalic acid, succinic acid, citric acid, homovanillic acid indicating elevated dopamine, methylmalonic acid (vitamin B12 deficiency), quinolinic acid, erythrocyte glutathione-S-transferase, serum glutathione-S-transferase, and blood sedimentation rate, high lactic acid, chronic acidosis
• Low: malic acid, 2-oxoglutaric acid, aconitic acid, serotonin, ratio of adrenalin/noradrenalin, blood sugar

EHS, electromagnetic hypersensitivity; IgE, immunoglobulin E.

Conclusions

Individual sensitivity of people to all kinds of environmental factors, natural and man-made, is common and, therefore, it is logical to consider that also individual sensitivity to man-made EMF exists, even if we still rely on anecdotal evidence from self-declared EHS persons and do not have sufficiently robust scientific proof of it.

Trust in each other is of paramount importance in such a type of survey study. Volunteers need to trust that the information they provide will be kept private and analyzed without bias. Scientists need to trust that volunteers provide honest and not exaggerated responses to all of the questions. Without such trust execution and analysis of such survey studies is very difficult.

The observations collected in this pilot questionnaire survey study show that currently, it is not possible to medically diagnose EHS as being a result of low-level wireless radiation exposures. It is possible that further development of the questionnaire might increase the likelihood of correct identification of EHS persons but, for the final confirmation of the EHS status, some physiological tests might be necessary. Currently, the medical diagnoses of EHS, claimed by some of the EHS persons, are based solely on the anecdotal evidence presented by the EHS person and, in rare cases, on tests that lack scientific proof of their validity as EHS diagnostic tools.

Further research on the causes of EHS and on the development of methods to diagnose it is necessary. However, it is necessary to keep in mind that the biomarkers of EHS might, or will likely, overlap with markers for other IEI-sensitivities or allergies. Future research studies should concomitantly examine both, physiological (12) and cognitive (30,31) responses of the human body to exposures to electromagnetic fields.

Open access: https://mhealth.amegroups.org/article/view/140469/html

--

Smartphone Usage Patterns and Sleep Behavior in Demographic Groups: Retrospective Observational Study

Wang T, Seiger A, Markowetz A, Andone I, Błaszkiewicz K, Penzel T. Smartphone Usage Patterns and Sleep Behavior in Demographic Groups: Retrospective Observational Study. J Med Internet Res 2025;27:e60423. doi: 10.2196/60423.

Abstract

Background: Although previous studies have examined the relationship between smartphone usage and sleep disorders, research on demographic differences in smartphone usage and nocturnal smartphone inactivity patterns remains limited. This study introduces “nocturnal smartphone inactivity duration” as a proxy indicator to address the limitation of lacking direct sleep data and to further investigate the association between smartphone usage patterns and sleep characteristics.

Objective: This study aimed to investigate demographic differences and relationships between daily smartphone usage and nocturnal smartphone inactivity patterns.

Methods: We conducted a retrospective analysis of data collected from the Murmuras app from January 1, 2022, to December 31, 2022. A total of 1074 participants were included, categorized by gender, age, highest degree, employment status, and smartphone usage purpose. All participants consented to participate in the study through the app. To explore the relationship between smartphone usage and nocturnal smartphone inactivity, we first calculated each participant's daily smartphone usage duration (including app usage) and duration of nocturnal smartphone inactivity; then, we assessed the normality and homogeneity of variance tests within each demographic category. Based on the results, the Kruskal-Wallis tests were applied to potentially identify differences between groups. Finally, correlation and regression analyses were conducted to explore associations between smartphone usage and nocturnal smartphone inactivity.

Results: The findings revealed distinct patterns of smartphone use across demographics. Participants predominantly used smartphones for social contact (average daily usage duration=1.52 h) and recreational activities (average daily usage duration=1.08 h) through apps like Facebook and YouTube. Frequent users, especially of social media and entertainment, often increased their phone usage at night. Female participants used their phones more frequently, mainly for digital shopping and social interactions, whereas male participants used phones more at nighttime (P<.001). Both younger users and non–full-time employees engaged more in activities such as gaming and chatting (P<.01 for those comparisons). Higher education was correlated with lower use (P<.001). Those using smartphones for work-related purposes generally decreased their phone usage after work (P<.05 for those comparisons). Correlation and regression analyses of smartphone usage duration and nighttime inactivity across groups indicated that only a small subset of groups exhibited significant positive correlations, a moderate number displayed significant negative correlations, and the majority showed no significant correlation.

Conclusions: This study underscores the significant association between demographic factors and smartphone usage patterns, including nocturnal inactivity patterns. Female individuals, young people, individuals with lower educational qualifications, and those who were unemployed demonstrated higher smartphone usage. Frequent engagement with social media and leisure apps was particularly pronounced during nighttime hours, a behavior that may contribute to disruptions in sleep patterns. These findings underscore the need for targeted interventions addressing excessive smartphone use, particularly at night, to mitigate its potential adverse effects on sleep.

Open access: https://www.jmir.org/2025/1/e60423

--

Temporal change of outdoor RF-EMF levels in four European countries: a microenvironmental measurement study

Beláčková L, Veludo AF, Aminzadeh R, Van Bladel H, Griffon V, Cardis E, Dongus S, Eeftens M, Guxens M, Joseph W, de Llobet P, Mazet P, Van Torre P, Thielens A, Vermeulen R, Wiart J, Röösli M, Huss A. Temporal change of outdoor RF-EMF levels in four European countries: a microenvironmental measurement study. Environ Res. 2025 Jul 11;285(Pt 1):122315. doi: 10.1016/j.envres.2025.122315.

Abstract

Introduction: Over the past two decades, the amount of transmitted mobile data has increased rapidly. It is unknown whether the implementation of the new technologies enabling this has resulted in changes of outdoor radio-frequency electromagnetic fields (RF-EMF) exposure. Therefore, microenvironmental measurements were used to investigate temporal trends in RF-EMF exposure between 2016 and 2023, in the Netherlands, Switzerland, Belgium and Spain, following a similar protocol across campaigns. Microenvironmental measurements refer to exposure measurements performed at predefined small areas that have been differentiated with a specific function in that particular area. This allowed us to compare exposure trends between countries and years.

Methods: The data was collected as part of the ACCEDERA (2016-2018), ETAIN (2023), and GOLIAT (2023) projects, walking repeatedly the same routes with RF-EMF exposimeters. Identical microenvironments were identified in each country and measurements of the exposure from mobile base stations, mobile phones and the total exposure were compared across years.

Results: Comparing measurements between 6 and 14 unique microenvironments in each country, our data did not suggest significant changes in the exposure from the mobile base station origin (total downlink exposure) between baseline measurements in 2016 to follow up and 2023 for the four countries. Across all countries and years the median values of the mobile base station exposure ranged from 0.11 mW/m2 (Switzerland, 2023) to 0.62 mW/m2 (Netherlands, 2018). There was no consistent trend in the individual microenvironments across the countries.

Conclusions: Our measurements of RF-EMF outdoor exposure levels across included microenvironment groups do not indicate change in exposure levels between 2016 and 2023 despite an increase in mobile data traffic by a factor of 8 in Western Europe[1].

Open access: https://www.sciencedirect.com/science/article/pii/S001393512501566X?via%3Dihub

--

Assessment of Electromagnetic Exposure to a Child and a Pregnant Woman Inside an Elevator in Mobile Frequencies

Karatsi I, Bakogianni S, Koulouridis S. Assessment of Electromagnetic Exposure to a Child and a Pregnant Woman Inside an Elevator in Mobile Frequencies. *Telecom*. 2025; 6(3):52. https://doi.org/10.3390/telecom6030052.

Abstract

This study presents an in-depth dosimetry analysis of energy assimilation from EM waves and increase in the temperature during mobile phone usage within an elevator cabin. The cellphone operates at two different frequencies (1000 MHz and 1800 MHz) and is simulated at three different talk positions vertical, tilt, and cheek. Realistic numerical models of a woman in the third trimester of pregnancy and a girl at the age of 5 years are employed. The analysis highlights the necessity of a comprehensive approach to fully grasp the complexities of EM exposure.

Conclusions

A comprehensive investigation into electromagnetic (EM) exposure and thermal safety has been conducted, focusing on three distinct subjects: a girl of five years named Roberta, a gestating female at seven months gestation known as Pregnant II, and her fetus. This study employed anatomically precise computational virtual humans along with a cellular device functioning at two different frequency bands, 1000 MHz and 1800 MHz. A total of 48 separate arrangements involving these numerical phantoms in various configurations and with different phone orientations were meticulously analyzed. The study presented detailed results of SAR10g, SAR-wb, and Tmax for every anatomical virtual human, including the fetus within the confines of an elevator cabin. The results here add to the findings in our prior work [55]. Here, contrary to [55], we examined how exactly, and in which scale, the positioning of the humans affects the quantities measured for all the utilized models, including the seven-month fetus.

Notably, it becomes evident that an integrated approach is essential to fully understand the complexities of EM exposure. Each configuration offers unique insights into how specific absorption rate values and temperature rises differ across subjects and settings. The variations observed underscore the necessity of examining every possible configuration to draw comprehensive conclusions about EM safety.

Moreover, the investigation highlights the importance of further exploring the effects of EM exposure in elevator cabins under various occupancy conditions and at different portable device frequencies. Such studies are crucial for developing targeted safety guidelines and protective measures, particularly in environments where RF exposure is compounded by enclosed spaces like elevators. This research not only enhances our understanding of EM field interactions with human tissues but also aids in the formulation of better health and safety standards for vulnerable populations such as children and pregnant women.

Open access: https://www.mdpi.com/2673-4001/6/3/52

--

Comparison Between Broadband and Personal Exposimeter Measurements for EMF Exposure Map Development Using Evolutionary Programming

Nájera A, Sánchez-Montero R, González-Rubio J, Guillén-Pina J, Chocano-del-Cerro R, López-Espí P-L. Comparison Between Broadband and Personal Exposimeter Measurements for EMF Exposure Map Development Using Evolutionary Programming. *Applied Sciences*. 2025; 15(13):7471. https://doi.org/10.3390/app15137471

Abstract

In this study, we provide a comparison of radiofrequency electromagnetic field exposure level maps as determined using two approaches: a broadband meter (NARDA EMR-300) equipped with an isotropic probe in the range of 100 kHz to 3 GHz, and a Personal Exposimeter (Satimo EME Spy 140) in the range of 88 MHz to 5.8 GHz. The aim of this research was to determine the necessary adjustments to the measurements made with personal exposimeters to obtain RF-EMF exposure maps equivalent to those made with broadband meters. We evaluated different possibilities to obtain the best equivalence of measurements between both devices. For this purpose, the datasets obtained in both cases were analyzed, as well as the possible correction factors. First, the possibility of establishing a single or double correction factor depending on the existence (or lack thereof) of a line of sight with respect to the base

stations was analyzed by minimizing the average value of the error between the values of the broadband meter and the corrected values of the personal exposure meter. Due to the differences observed in the exposure maps, a second procedure was carried out, in which a genetic algorithm was used to determine the ratio between the measurements from both methods (the broadband meter and personal exposure meter), depending on the existence (or lack thereof) of a line of sight, and we compared the exposure maps generated using kriging interpolation.

Conclusions

We conclude that exposure assessments based on spot measurements can serve as a practical proxy to evaluate personal RF-EMF exposure, whether using a static probe or a portable PEM. The developed methodology offers a reliable and efficient approach to characterizing personal exposure levels, enabling continuous monitoring and the detection of potential changes over time. The result of the different proportionality factors obtained is shown in Table 4.

The application of correction factors, particularly the differentiation between LOS and NLOS conditions, significantly improves the accuracy of PEM measurements, bringing them closer to those obtained with BBMs. This highlights the importance of accounting for the line of sight to antennas when estimating exposure levels, as direct exposure to base stations considerably affects measurement accuracy due to body shielding effects. By applying appropriate correction factors, the inherent limitations of PEMs, such as underestimating exposure levels in LOS conditions, can be mitigated.

Moreover, using genetic algorithms to optimize these corrections further enhances the precision of the exposure maps, demonstrating that this method can be a valuable tool in urban RF-EMF monitoring. The ability to generate accurate spatial exposure distributions through PEMs makes this methodology more accessible and cost effective, especially in large-scale studies where deploying multiple BBMs might not be feasible.

Nevertheless, further validation is needed in various environments to refine the correction models and ensure broader applicability. Future studies should focus on refining these algorithms and exploring other factors that could influence exposure, such as linearity, urban infrastructure, and signal interference. Despite these challenges, the methodology developed in this study represents a significant advancement in RF-EMF exposure assessment, offering a flexible and scalable approach to creating consistent EMF exposure maps using PEMs.

Open access: https://www.mdpi.com/2076-3417/15/13/7471

--

Machine Learning Approach for Ground-Level Estimation of Electromagnetic Radiation in the Near Field of 5G Base Stations

Famoriji OJ, Shongwe T. Machine Learning Approach for Ground-Level Estimation of Electromagnetic Radiation in the Near Field of 5G Base Stations. *Applied Sciences*. 2025; 15(13):7302. https://doi.org/10.3390/app15137302

Abstract

Electromagnetic radiation measurement and management emerge as crucial factors in the economical deployment of fifth-generation (5G) infrastructure, as the new 5G network emerges as a network of services. By installing many base stations in strategic locations that operate in the millimeter-wave range, 5G services are able to meet serious demands for bandwidth. To evaluate the ground-plane radiation level of electromagnetics close to 5G base stations, we propose a unique machine-learning-based approach. Because a machine learning algorithm is trained by utilizing data obtained from numerous 5G base stations, it exhibits the capability to estimate the strength of the electric field effectively at every point of arbitrary radiation, while the base station generates a network and serves various numbers of 5G terminals running in different modes of service. The model requires different numbers of inputs, including the antenna's transmit power, antenna gain, terminal service modes, number of 5G terminals, distance between the 5G terminals and 5G base station, and environmental complexity. Based on experimental data, the estimation method is both feasible and effective; the machine learning model's mean absolute percentage error is about 5.89%. The degree of correctness shows how dependable the developed technique is. In addition, the developed approach is less expensive when compared to measurements taken on-site. The results of the estimates can be used to save test costs and offer useful guidelines for choosing the best location, which will make 5G base station electromagnetic radiation management or radio wave coverage optimization easier.

Open access: https://www.mdpi.com/2076-3417/15/13/7302

--

Cluster Analysis of RF-EMF Exposure to Detect Time Patterns in Urban Environment: A Model-Based Approach

Pasquino N, Solmonte N, Djuric N, Kljajic D, Djuric S. Cluster Analysis of RF-EMF Exposure to Detect Time Patterns in Urban Environment: A Model-Based Approach. IEEE Access, vol. 13, pp. 118724-118732, 2025, doi: 10.1109/ACCESS.2025.3586905.

Abstract

The increase in human exposure to electromagnetic fields (EMFs), driven by advancements in telecommunication systems like the 5G mobile system, highlights the need for continuous EMF monitoring. Advanced techniques for data analysis, based on machine learning like clustering, can decompose daily variations in EMF exposure into distinct patterns, providing a clearer understanding of how exposure fluctuates over time. Although several exposure monitoring systems exist in Europe, only a few studies have thoroughly examined the time variability. This study addresses the gap by applying model-based clustering techniques to analyze the temporal patterns. Specifically, the study focuses on characterizing fluctuations in field strength during workdays and holidays, thereby contributing to a deeper understanding of time-distributed exposure. Continuous monitoring data, collected through the Serbian EMF RATEL network's sensors installed in Novi Sad, were processed and analyzed using the Log-Normal Mixture Model (LNMM), a model-based clustering algorithm resorting to mixture distributions. The analyses reveal that the LNMM can separate night and day exposure values and identify periods when values persist longer over the day. This suggests that model-based clustering can be useful for understanding the temporal patterns of local EMF exposure.

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=11072665&isnumber=10820123

--

Impact of Anthropomorphic Shape and Skin Stratification on Absorbed Power Density in mmWaves Exposure Scenarios

Gallucci S, Benini M, Bonato M, Galletta V, Chiaramello E, Fiocchi S, Tognola G, Parazzini M. Impact of Anthropomorphic Shape and Skin Stratification on Absorbed Power Density in mmWaves Exposure Scenarios. Sensors (Basel). 2025 Jul 17;25(14):4461. doi: 10.3390/s25144461.

Abstract

As data exchange demands increase also in widespread wearable technologies, transitioning to higher bandwidths and mmWave frequencies (30-300 GHz) is essential. This shift raises concerns about RF exposure. At such high frequencies, the most crucial human tissue for RF power absorption is the skin, since EMF penetration is superficial. It becomes thus very important to assess how the model used to represent the skin in numerical dosimetry studies affects the estimated level of absorbed power. The present study, for the first time, assesses the absorbed power density (APD) using FDTD simulations on two realistic human models in which: (i) the skin has a two-layer structure made of the stratum corneum and the viable epidermis and dermis layers, and (ii) the skin is modelled as a homogeneous dermis stratum. These results were compared with ones using flat phantom models, with and without the stratified skin. The exposure assessment study was performed with two sources (a wearable patch antenna and a plane wave) tuned to 28 GHz. For the wearable antenna, the results evidence that the exposure levels obtained when using the homogeneous version of the models are always lower than the levels in the stratified skin version with percentage differences from 16% to 30%. This trend is more noticeable with the female model. In the case of plane wave exposure, these differences were less pronounced and lower than 11%.

Open access: https://www.mdpi.com/1424-8220/25/14/4461

--

Weak Radiofrequency Field Effects on Biological Systems Mediated through the Radical Pair Mechanism

Gerhards L, Deser A, Kattnig DR, Jörg Matysik J, Solov'yov IA. Weak Radiofrequency Field Effects on Biological Systems Mediated through the Radical Pair Mechanism. Chemical Reviews. July 14, 2025. doi: 10.1021/acs.chemrev.5c00178.

Abstract

The widespread use of radiofrequency (RF) communication has increased the exposure of organisms to electromagnetic fields, sparking a debate over the potential health effects of weak RF electromagnetic fields. While some experimental studies suggest that low-amplitude RF radiation may influence cellular metabolism or sleep patterns or even promote cancer, these claims remain controversial due to limited theoretical plausibility. Central to this debate is the radical pair mechanism (RPM), a quantum-mechanical framework proposed to mediate RF effects. Despite its role in magnetoreception and various magnetic field effects on chemical reactions, the RPM often fails to align with observations at low, nonthermal RF field strengths. This review examines these contrasting perspectives by discussing

experimental findings and theoretical models that aim to explain putative biological effects of RF magnetic fields. Emphasis is placed on the challenges of reconciling theoretical predictions with empirical data, particularly in the context of weak RF exposure. Additionally, an overview of the theoretical framework used in current modeling efforts highlights the complexity of applying the RPM to biological systems and underscores the importance of critical interpretation. The goal is to clarify the state of understanding and inform future research on RPM-mediated biological effects under weak RF exposure.

Concluding Remarks

This review provides a detailed overview of the current state of research on weak RF-MF-induced biological effects in the context of the RPM. While several experimental studies suggest observable effects, their reproducibility and dosimetry remain significantly challenging. The RPM is frequently proposed as a mechanistic explanation, yet current theoretical models based on spin dynamics often struggle to fully account for experimental findings.

As demonstrated, accurately modeling the RPM is nontrivial, requiring consideration of multiple interactions and features. Essential interactions, such as spin–orbit coupling or dipolar coupling, are often overlooked in studies of potential MFEs. Additionally, decoherence due to spin relaxation and the short lifetimes of SCRPs impose further limitations on the detectability of weak RF-MF effects.

Despite these challenges, this review highlights a clear need for more systematic and interdisciplinary investigations. Experimental studies require robust statistical designs, standardized

methodologies, and improved dosimetry to enhance reproducibility. Meanwhile, theoretical approaches must incorporate the full complexity of spin dynamics beyond simplified models.

Advances such as the SSE method offer promising directions to overcome current limitations.

Ultimately, bridging the gap between experiment and theory will be key to unraveling the role of the RPM in weak RF-MF effects. Future research integrating rigorous theoretical frameworks with high-quality experimental data will be essential to clarify whether and how weak RF fields influence biological systems.

Open access: https://pubs.acs.org/doi/10.1021/acs.chemrev.5c00178

--

Magnetic Field Measurement of Various Types of Vehicles, Including Electric Vehicles

Fukui H, Minami N, Tanezaki M, Muroya S, Ohkubo C. Magnetic Field Measurement of Various Types of Vehicles, Including Electric Vehicles. *Electronics*. 2025; 14(15):2936. doi: 10.3390/electronics14152936

Abstract

Since around the year 2000, following the introduction of electric vehicles (EVs) to the market, some people have expressed concerns about the level of magnetic flux density (MFD) inside vehicles. In 2013, we reported the results of MFD measurements for electric vehicles (EVs), hybrid electric vehicles (HEVs), and internal combustion engine vehicles (ICEVs). However, those 2013 measurements were conducted using a chassis dynamometer, and no measurements were taken during actual driving. In recent years,

with the rapid global spread of EVs and plug-in hybrid electric vehicles (PHEVs), the international standard IEC 62764-1:2022, which defines methods for measuring magnetic fields (MF) in vehicles, has been issued. In response, and for the first time, we conducted new MF measurements on current Japanese vehicle models in accordance with the international standard IEC 62764-1:2022, identifying the MFD levels and their sources at various positions within EVs, PHEVs, and ICEVs. The measured MFD values in all vehicle types were below the reference levels recommended by the International Commission on Non-Ionizing Radiation Protection (ICNIRP) for public exposure. Furthermore, we performed comparative measurements with the MF data obtained in 2013 and confirmed that the MF levels remained similar. These findings are expected to provide valuable insights for risk communication with the public regarding electromagnetic fields, particularly for those concerned about MF exposure inside electrified vehicles.

Conclusions

As the adoption of EVs and PHEVs increases, public concern about MFs from these vehicles has also grown. Narrowing the perception gap between scientific risk assessments and public anxiety is essential for effective risk communication. In this study, we conducted measurements of MFs emitted by domestic vehicles in Japan using IEC 62764-1:2022 methods and were able to provide reliable and easily accessible data to the public as a basis for risk communication.

The results confirmed that the measured MFs comply with ICNIRP guidelines and remain at levels that do not cause known acute adverse effects, such as nerve stimulation. The locations and characteristics of the maximum MF strengths for EVs, PHEVs, and ICEVs were identified.

For EVs and PHEVs, the highest MF values were observed on the rear seat at a 6.5 cm measurement distance, while for ICEVs, the maximum was recorded at the driver's side dashboard at 20 cm. Among the vehicle types, PHEVs showed the highest MF levels, followed by ICEVs, which had higher values than EVs.

As a result of the frequency analysis, both speed-dependent and speed-independent MF components were detected across all vehicle types. Speed-dependent peaks included 6–29 Hz frequency components near the front seats, attributed to magnetized tires. Fixed-frequency peaks included a 294 Hz component observed during air conditioner operation. Low-frequency peaks around 1 Hz were observed near wiring routes around the rear seat area and were attributed to current flow during vehicle operation, with notably higher values in the PHEV during deceleration. In ICE vehicles, low-frequency components in the 1–5 Hz range were associated with wiper motor operation, while higher-frequency peaks at 847 Hz and 1268 Hz were detected but could not be clearly linked to specific sources.

A comparison with a previous study conducted in 2013 revealed that the MF levels are comparable to those of current vehicles as of 2025.

A comparison with the MF measurement results from Seibersdorf Laboratory in Austria revealed that their reported values were significantly higher. This difference is attributed to their capturing of rapid increases in MFD due to transient phenomena occurring on timescales below 200 ms, which were not considered in this study, and reflects differences in measurement conditions.

These findings are expected to provide valuable insights for risk communication with the general public regarding EMFs, particularly for those concerned about MF exposure inside electrified vehicles.

As a future perspective, as the performance of EVs continues to improve each year, with increases in motor output and battery capacity, exposure to MFs is expected to rise. Therefore, it is important to continue publishing measurement results regularly in the future.

This study focused on the vehicle itself and did not include measurements of the charging equipment. Wireless charging systems are being developed as an emerging technology [27,28,29,30,31], and demonstration experiments are currently underway in Japan. As their adoption is expected to increase in the future, opportunities for risk communication with the general public are likely to grow. It is therefore important to conduct MF exposure measurements in line with public road deployment and to advance risk communication accordingly.

In addition to passenger cars, the electrification of other vehicles such as buses is advancing, and the introduction of new mobility technologies, including the construction of linear motor cars, is expected to progress in the future. As such technologies are introduced, public concern about EMF exposure is likely to increase. Therefore, it is important to consider conducting measurements in line with the pace of their deployment.

Open access: https://www.mdpi.com/2079-9292/14/15/2936

--

Analysis of Human Exposure to Electric and Magnetic Fields While Charging and Driving an Electric Vehicle

Gliga, M. *et al.* (2025). Analysis of Human Exposure to Electric and Magnetic Fields While Charging and Driving an Electric Vehicle. In: Vlad, S., Roman, N.M. (eds) 9th International Conference on Advancements of Medicine and Health Care Through Technology. MEDITECH 2024. IFMBE Proceedings, vol 130. Springer, Cham. https://doi.org/10.1007/978-3-031-95671-3_25

Abstract

The transition to electric vehicles is a promising strategy to reduce or eliminate harmful emissions, including carbon dioxide, nitrogen oxides, and particulate matter, from internal combustion vehicles. While electric vehicles offer environmental benefits, their operation involves the generation of electromagnetic fields (EMFs) in their immediate vicinity. Although the biological effects of EMFs have been extensively studied, the specific health risks to humans remain uncertain.

In electric vehicles, occupants are exposed to artificial EMFs due to their close proximity to high-capacity electrical systems operating for extended periods. Recognizing these potential concerns, electric vehicle manufacturers have incorporated various technological solutions during the design and manufacturing processes to mitigate EMF exposure. This study will conduct experimental measurements of the electromagnetic field generated by an electric vehicle under a range of operational conditions.

https://link.springer.com/chapter/10.1007/978-3-031-95671-3_25

--

Standards: Exposure Limits for Brief High Intensity Pulses of Radiofrequency Energy Between 6 and 300 GHz

K. R. Foster, I. Laakso and Q. Balzano, "Standards: Exposure Limits for Brief High Intensity Pulses of Radiofrequency Energy Between 6 and 300 GHz," in *IEEE Access*, vol. 13, pp. 134648-134654, 2025, doi: 10.1109/ACCESS.2025.3592532.

Abstract

This Standards paper considers limits of the International Commission on Nonionizing Radiation (ICNIRP) and Institute of Electrical and Electronics Engineers (IEEE C95.1-2019) on “brief exposures” to radiofrequency energy (RF-EMF), that vary significantly within the 6 minute averaging time specified in the limits. This situation arises in two important contexts: exposures from cellular base stations employing Multiple Input Multiple Output (MIMO) antennas with beamforming that sweep beams around the service area, and exposure to pulsed RF-EMF with high peak power but low duty cycle. The ICNIRP and IEEE limits on brief exposures between 6 and 300 GHz are assessed using numerical and analytical solutions to a one-dimensional thermal model of tissue. Analysis is done both for rectangular pulses of RF-EMF (i.e., the step response to a suddenly imposed RF-EMF field), and to impulses of RF-EMF energy. The analysis shows that both sets of guidelines are uneven in the level of protection they offer against excessive transient heating of skin, with the IEEE limits being far more conservative than the ICNIRP limits. The thermal model suggests a simple modification to the pulse fluence limits that provides a more consistent level of protection against excessive temperature increases for pulses of varying duration and frequency. **The Standards Paper does not consider possible nonthermal hazards or possible thermoacoustic effects produced by brief but very high amplitude RF-EMF pulses that produce high rates of change of tissue temperature but small temperature increases**. The conclusion points to the need for further research and development to establish and refine exposure limits for pulsed RF-EMF.

Open access: https://ieeexplore.ieee.org/document/11095721

--

Weak Radiofrequency Field Effects on Biological Systems Mediated through the Radical Pair Mechanism

Luca Gerhards, Andreas Deser, Daniel R. Kattnig, Jörg Matysik, and Ilia A. Solov’yov. Weak Radiofrequency Field Effects on Biological Systems Mediated through the Radical Pair Mechanism. *Chemical Reviews.* doi: 10.1021/acs.chemrev.5c00178

Abstract

The widespread use of radiofrequency (RF) communication has increased the exposure of organisms to electromagnetic fields, sparking a debate over the potential health effects of weak RF electromagnetic fields. While some experimental studies suggest that low-amplitude RF radiation may influence cellular metabolism or sleep patterns or even promote cancer, these claims remain controversial due to limited theoretical plausibility. Central to this debate is the radical pair mechanism (RPM), a quantum-mechanical framework proposed to mediate RF effects. Despite its role in magnetoreception and various magnetic field effects on chemical reactions, the RPM often fails to align with observations at low, nonthermal RF field strengths. This review examines these contrasting perspectives by discussing experimental findings and theoretical models that aim to explain putative biological effects of RF magnetic fields. Emphasis is placed on the challenges of reconciling theoretical predictions with

empirical data, particularly in the context of weak RF exposure. Additionally, an overview of the theoretical framework used in current modeling efforts highlights the complexity of applying the RPM to biological systems and underscores the importance of critical interpretation. The goal is to clarify the state of understanding and inform future research on RPM-mediated biological effects under weak RF exposure.

Open access: https://pubs.acs.org/doi/10.1021/acs.chemrev.5c00178

--

Effect of the radiation emitted from a cell phone on T lymphocytes in mice

Pei, Y., Gao, H., Li, J., Li, J., Hou, Z., & Han, X. (2025). Effect of the radiation emitted from a cell phone on T lymphocytes in mice. *Radiation Effects and Defects in Solids*, 1–13. https://doi.org/10.1080/10420150.2025.2526402

Abstract

This present study aims to explore the potential impact of cell phone radiation on T lymphocytes in mice. One hundred twenty-eight healthy male BALB/C mice (2–3 weeks old) were randomly distributed into four groups: blank control, control, TD-SCDMA and LTE-Advanced groups, respectively, with 32 mice in each group. Mice were exposed to phone radiation for 2–8 weeks. 8 mice in each group were taken out for measurement, and given exposure periods were 2, 4, 6 and 8 weeks, respectively. Flow cytometry was performed for the examination of T lymphocyte subsets, molecular and cell biological techniques were carried out to detect the apoptosis and transformation of T lymphocytes, respectively, while the enzyme-labelled immune technique was conducted to measure the cytokines secreted by T lymphocytes. The results showed that the percentage of CD4 T lymphocyte subsets was similar among the 4 groups during the entire exposure period, and so were the CD8 T lymphocyte subsets. Compared to the 2 sham groups, the apoptosis ratios of T lymphocytes in the TD-SCDMA and LTE-Advanced groups increased significantly after more than six weeks of exposure, while the transformation rates decreased significantly. The impacts of radiation on apoptosis and transformation are time-dependent. The concentrations of IL-10 and IL-12 remained unchanged in the 2 control groups during the entire experimental period. Compared to the 2 control groups, the concentrations of IL-10 in the 2 exposure groups decreased significantly, while the concentrations of IL-12 increased statistically. Furthermore, the decrease in IL-10 lagged behind the increase in IL-12. Therefore, it could be deduced that radiation from mobile phones resulted in T lymphocyte injury, and this adverse effect was seemingly more severe as the radiation exposure continued. Additionally, cell phone radiation was seen to suppress the secretion of IL-10 while enhancing the production of IL-12.

https://www.tandfonline.com/doi/full/10.1080/10420150.2025.2526402

--

Genotoxic and histopathological effects of 6 GHz radiofrequency electromagnetic radiation on rat liver tissue

Ilgaz NS, Karamazı Y, Emre M, Toyran T, Karaoğlan Ö, Emre T, Dönmez Kutlu M, Öksüz Üçkayabaşı H, Aydın Ç, Yılmaz MB. Genotoxic and histopathological effects of 6 GHz radiofrequency electromagnetic

radiation on rat liver tissue. Electromagn Biol Med. 2025 Jul 22:1-12. doi: 10.1080/15368378.2025.2534381.

Abstract

In this study, the genotoxic and histopathological effects of 6 GHz (0.065 W/kg) Radiofrequency-Electromagnetic Radiation (RF-EMR) on rat liver tissue were investigated. Sham (control) and Radiofrequency Radiation (RFR) groups were formed with 10 adult male rats in each group. Rats in the sham group received no treatment. Rats in the RFR group were exposed to 6 GHz RF-EMR for 4 h/day for 42 days. Immediately after the completion of the exposure, the rats in both groups were sacrificed and liver tissues were removed. Comet Test was performed to determine the genotoxic effect in the samples. Masson Trichrome and Hematoxylin Eosin staining methods were applied histopathologically. According to the Comet Analysis results, the genetic damage index (GDI) and damaged cell percentage (DCP) of the RFR group were higher than the sham group, but this difference was not statistically significant ($p > 0.05$). In histopathologic examinations, portal inflammation, single cell necrosis, vascularity and congestion were more prominent in the RFR group compared to the sham group. In our study, it was shown that 6 GHz RF-EMR can cause histopathologic and DNA level changes in rat liver tissue. As a result of the literature review, no prior studies have specifically examined the genotoxic and histopathological effects of 6 GHz RF-EMR. This makes our study important as it addresses the biological impacts of the 6 GHz frequency band.

https://www.tandfonline.com/doi/full/10.1080/15368378.2025.2534381

--

Single-cell analysis reveals the spatiotemporal effects of long-term electromagnetic field exposure on the liver

Zhang M, Lv Z, Zhao L, Zeng Q, Wu Y, Zhou J, Xi J, Pei X, Wang H, Li C, Yue W. Single-cell analysis reveals the spatiotemporal effects of long-term electromagnetic field exposure on the liver. Front Cell Dev Biol. 2025 Jun 27;13:1579121. doi: 10.3389/fcell.2025.1579121.

Abstract

Introduction: Artificial electromagnetic fields (EMFs) can impair the functions of several organs. The impact of long-term artificial EMF on the liver, the synthetic and metabolic center of the body, has become concerning. The aim of this study was to systematically evaluate the effect of long-term EMF exposure on the liver.

Methods: Mice were exposed to 2.45 GHz EMF daily for up to 5 months, and serum liver function test, lipidomic analysis, and histological analysis were performed to detect the general impact of EMF on the liver. Furthermore, EMF-induced liver transcriptome variations were investigated using single-cell RNA sequencing and a spatiotemporally resolved analysis.

Results: Different hepatic cells exhibited diverse sensitivities and response patterns. Notably, hepatocytes, endothelial cells, and monocytes showed higher sensitivity to electromagnetic radiation, with their lipid metabolic functions, immune regulation functions, and intrinsic functions disturbed,

respectively. Moreover, transcriptomic alterations were predominantly observed in the hepatocytes and endothelial cells in peri-portal regions, suggesting a zonation-related sensitivity to EMF within the liver.

Conclusion: Our study provided a spatiotemporal visualization of EMF-induced alterations in hepatic cells, which ultimately elucidated the biological effects of EMF exposure.

Open access: https://www.frontiersin.org/journals/cell-and-developmental-biology/articles/10.3389/fcell.2025.1579121/full

--

Effects of coenzyme Q10 on sperm parameters and pathological changes induced by Wi-Fi waves in the testicular tissue of rats

Safaeinezhad, Ali et al. Effects of coenzyme Q10 on sperm parameters and pathological changes induced by Wi-Fi waves in the testicular tissue of rats. Annals of Medicine & Surgery 87(7):p 4073-4079, July 2025. | doi: 10.1097/MS9.0000000000003340

Abstract

Background: Despite safety boundaries set by authorities, there is public concern about the health effects of wireless communication tools using radiofrequency. The World Health Organization has advised further research in this area.

Objective: The aim of this study was to investigate the effect of coenzyme Q10 (CoQ10) on sperm parameters and pathological changes in the testicular tissue in rats.

Methods: Twenty healthy rats were divided into four groups (n = 5). The first group was kept in a Wi-Fi-free environment for 2 months. The second group was exposed to Wi-Fi waves for 7 h a day for 2 months. The third group was exposed to Wi-Fi waves for 7 h a day for 2 months and received 150 mg/kg of CoQ10 daily. The fourth group was not exposed to Wi-Fi waves but received 150 mg/kg of CoQ10 daily. Serum testosterone levels, sperm parameters, and pathological findings of testis and epididymis were investigated.

Results: Treatment with CoQ10 (150 mg/day) resulted in a significant increase in sperm concentration from baseline, viability, and motility compared with the Wi-Fi group. CoQ10 therapy also increased testosterone levels compared with the Wi-Fi group. The pathology findings showed that in the treatment with CoQ10, fewer tubes had stopped the process of spermatogenesis due to Wi-Fi.

Conclusion: The study results indicated that the intake of CoQ10 can reduce the complications caused by Wi-Fi on sperm parameters, testosterone levels, and pathological findings.

Open access: https://journals.lww.com/annals-of-medicine-and-surgery/fulltext/2025/07000/effects_of_coenzyme_q10_on_sperm_parameters_and.16.aspx

--

A Prolonged exposure to Wi-Fi Radiation Induces Neurobehavioral Changes and Oxidative Stress in Adult Zebrafish

Swati Fumakiya, Jahnavi Mehta, Aditi Punekar, Payal Sharma, Gowri Kumari Uggini. A Prolonged exposure to Wi-Fi Radiation Induces Neurobehavioral Changes and Oxidative Stress in Adult Zebrafish. International Journal For Multidisciplinary Research. 7(4): July-August 2025. doi: 10.36948/ijfmr.2025.v07i04.52286.

Abstract

The rapid development and use of wireless technologies have raised concerns about the possible harmful effects of long-term exposure to non-ionizing electromagnetic radiation (EMR). Wi-Fi routers are the most commonly used devices at households, offices, hotels, hospitals, airports, train stations etc., to provide wireless internet access to multiple devices. Therefore, prolonged exposure to EMR warrants thorough scientific investigation and risk assessment. Herein, we evaluated the impact of 2.45GHz Wi-Fi radiation exposure for 4 hours daily over a consecutive 30-day period on neurobehavioral and brain oxidative stress parameters in adult zebrafish. Post-exposure assessments revealed significant neurobehavioral impairments, altered locomotion, decreased neurotransmitter levels (AChE), and increased oxidative stress. Further studies are needed to better understand the molecular mechanisms involved and to assess the broader implications for vertebrate neurobiology and public health.

Open access: https://www.ijfmr.com/research-paper.php?id=52286

--

Investigation of the Effects of 2.45 GHz Near-Field EMF on Yeast

Angelova B, Paunov M, Kitanova M, Atanasova G, Atanasov N. Investigation of the Effects of 2.45 GHz Near-Field EMF on Yeast. Antioxidants (Basel). 2025 Jul 3;14(7):820. doi: 10.3390/antiox14070820.

Abstract

The study of the effects of 2.45 GHz electromagnetic fields on the health and safety of people and organisms as a whole is essential due to their widespread use in everyday life. It is known that they can cause thermal and non-thermal effects-at the molecular, cellular and organismal level. Yeast suspensions were treated with 2.45 GHz microwave radiation in the near-field of antenna at two distances (2 and 4 cm) and two time periods (20 and 60 min)-setups resembling the use of mobile devices. The release of UV-absorbing substances from the cells was studied as an indicator of membrane permeabilization, total intracellular antioxidant activity and reduced glutathione were determined, and a comet assay for damage to the DNA was performed. A correlation between reduced antioxidants and increased membrane permeability during EMF treatment was observed at a distance of 2 cm for 20 min, suggesting the presence of oxidative stress, while a similar effect was not observed with conventional heating. Slightly increased membrane permeability was observed after irradiation for 60 min at a distance of 4 cm, but this was not related to the antioxidant status of the cells. A trend towards increased DNA damage was observed under both conditions.

Open access: https://www.mdpi.com/2076-3921/14/7/820

--

Low frequency magnetic field exposure and neurodegenerative disease: systematic review of animal studies

Stam R. Low frequency magnetic field exposure and neurodegenerative disease: systematic review of animal studies. Electromagn Biol Med. 2025 Aug 3:1-15. doi: 10.1080/15368378.2025.2540435.

Abstract

Epidemiological studies have found an association between occupational exposure to low frequency magnetic fields and the occurrence of motor neuron disease and Alzheimer's disease. No association has been found for Parkinson's disease and the evidence for multiple sclerosis is insufficient. Animal models studying the effects of low frequency magnetic fields on neurodegenerative disease induction or progression could provide more evidence on causation and the underlying mechanisms. A systematic search and review was conducted of peer-reviewed research articles involving animal experiments on the effects of low frequency magnetic field exposure on behavioural and neuroanatomical outcomes relevant for neurodegenerative diseases in humans. Firstly, experimental studies in *naive animals* do not support a causal relationship between exposure to low frequency magnetic fields and the induction of neuropathology relevant for Alzheimer's disease, but the number of studies relevant for motor neuron disease, multiple sclerosis and Parkinson's disease is too limited to draw conclusions. Secondly, experimental studies in *existing animal models for neurodegenerative disease* support a therapeutic (beneficial) effect of low frequency magnetic field treatment on behavioural and neuroanatomical abnormalities relevant for dementia (including Alzheimer's disease), multiple sclerosis and Parkinson's disease and no effect on disease progression in models relevant for motor neuron disease.

https://pubmed.ncbi.nlm.nih.gov/40754996/

--

Proteomic Characterization of Human Peripheral Blood Mononuclear Cells Exposed to a 50 Hz Magnetic Field

Bracci M, Lazzarini R, Piva F, Giulietti M, Marinelli Busilacchi E, Rossi E, Di Criscio F, Santarelli L, Poloni A. Proteomic Characterization of Human Peripheral Blood Mononuclear Cells Exposed to a 50 Hz Magnetic Field. Int J Mol Sci. 2025 Jun 24;26(13):6035. doi: 10.3390/ijms26136035.

Abstract

Exposure to extremely low-frequency magnetic fields (ELF-MF) can induce biological alterations in human cells, including peripheral blood mononuclear cells (PBMCs). However, the molecular mechanisms and key regulatory factors underlying this cellular response remain largely unknown. In this study, we analyzed the proteomic profiles of PBMCs isolated from three human subjects. PBMCs were exposed to 50 Hz, 1 mT of ELF-MF for 24 h and compared to unexposed PBMCs from the same individuals. ELF-MF exposure altered the expression levels of several PBMC proteins without affecting cell proliferation, cell viability, or cell cycle progression. A total of 51 proteins were upregulated, 36 of which were intercorrelated and associated with the *Cellular Metabolic Process* (GO:0044237) and *Metabolic Process* (GO:0008152). Among them, solute carrier family 25 member 4 (SLC25A4), which catalyzes the exchange of cytoplasmic ADP for mitochondrial ATP across the inner mitochondrial membrane, was consistently upregulated in all ELF-MF-exposed samples. Additionally, 67 proteins were

downregulated, many of which are linked to *T cell costimulation* (GO:0031295), *Cell activation* (GO:0001775), and *Immune system processes* (GO:0002376) included ASPSCR1, PCYT1A, PCYT2, QRAS, and REPS1. In conclusion, ELF-MF exposure induces metabolic reprogramming in human PBMCs, characterized by the upregulation of mitochondrial proteins and downregulation of immune-activation-related proteins, without compromising cell viability or proliferation.

Open access: https://www.mdpi.com/1422-0067/26/13/6035

--

Microleakage of Amalgam Restorations after Exposure to Electromagnetic Fields of a Commercial Hair Dryer: An Ex-Vivo Study

Paknahad M, Dehghani A, Khaleghi I, Eghildespour M, Mortazavi G, Mortazavi SMJ. Microleakage of Amalgam Restorations after Exposure to Electromagnetic Fields of a Commercial Hair Dryer: An *Ex-Vivo* Study. J Biomed Phys Eng. 2025 Jun 1;15(3):263-270. doi: 10.31661/jbpe.v0i0.2210-1551.

Abstract

Background: Dental amalgam is a popular restorative material used in posterior teeth. Hair dryers can emit electromagnetic fields (EMFs) that may affect the microleakage of the amalgam-tooth interface.

Objective: The aim of this experimental study was to investigate whether the EMFs produced by commercial hair dryers could cause microleakage in amalgam restorations.

Material and methods: In this experimental study, a total of 100 human extracted teeth without cavities were selected and prepared for class V preparations on their buccal aspects. The teeth were divided into five groups (G1-G5), each containing 20 teeth. Group 1 served as the control group and was not subjected to any treatment. Groups 2 to 5 were exposed to EMFs of a hair dryer (2000 W, 220 V, and 50 Hz). Groups 2 and 3 were exposed to "EMFs +Hot Air" for 20 min at 10 cm and 30 min at 5 cm, respectively. Groups 4 and 5 were exposed to "EMFs +Cool Air" for 20 min at 10 cm and 30 min at 5cm, respectively. After preparation, the sectioned teeth were evaluated for microleakage using dye penetration measurement.

Results: The microleakage scores showed a significant difference among the three exposure groups (G2, G3, and G5) and the control group (P=0.001, 0.002, and 0.01, respectively). However, there was no significant difference between G4 and G1. The microleakage score in G2 was higher than that in G4.

Conclusion: This study suggests that the common use of hair dryers can lead to damage in amalgam restorations.

Open access: https://jbpe.sums.ac.ir/article_49466.html

--

Radiofrequency electromagnetic fields reduce bumble bee visitation to flowers

Treder M, Glück M, England SJ, Traynor KS. Radiofrequency electromagnetic fields reduce bumble bee visitation to flowers. Environ Pollut. 2025 Jul 14:126836. doi: 10.1016/j.envpol.2025.126836.

Abstract

Radiofrequency electromagnetic fields (RF-EMF) are an integral part of our daily lives and we may struggle to imagine life without them, using them to transmit and stream data for television, the internet, or cell phones. Debates circulate on whether this anthropogenic radiation may be an additional stress factor for pollinators, potentially disrupting important habitats and refugia, especially in highly developed areas. Could these anthropogenic fields interfere with foraging pollinators, influence their foraging behavior or reduce the attractiveness of entire habitats? Current data on sublethal impacts is sparse. We tested the effects of realistic 2.4 and 5.8 GHz radiation on flower visitation rates of different pollinators over two years in a blinded study using a standardized radiation source and a novel study design. Documenting 2,876 pollinators, we found no effect of this radiation on honey bee visitation rates on Salvia and Lavandula plants. However, we found a significant reduction in the number of bumble bee visits per observation when the plants were subjected to RF-EMF exposure, highlighting the need for in-depth and long-term studies to elucidate the underlying mechanisms of this effect.

https://www.sciencedirect.com/science/article/abs/pii/S0269749125012096?via%3Dihub

--

Perspectives on terahertz honeybee sensing

Prokscha A, Sheikh F, Jalali M, De Boose P, De Borre E, Jeladze V, Ribas FO, Carvajal DT, Svejda JT, Kubiczek T, Aqlan B, Alibeigloo P, Mutlu E, Watermann J, Abts J, Kress R, Preuss C, Clochiatti S, Wiedau L, Weimann NG, Balzer JC, Thielens A, Kaiser T, Erni D. Perspectives on terahertz honey bee sensing. Sci Rep. 2025 Mar 27;15(1):10638. doi: 10.1038/s41598-025-91630-8.

Abstract

Terahertz (THz) technology provides precise monitoring capabilities in dynamic environments, offering unique insights into insect habitats. Our study focuses on environmental monitoring of European honey bees (Apis mellifera) through a combination of measurements and simulations. Initially, the dielectric material properties of honey bee body parts are characterized across the spectral range of 1-500 GHz to collect heterogeneous empirical data. To extend the study, honey bee mockups made from polyamide 12 (PA12) and epoxy resin are employed and validated as effective substitutes for real bees through comparative scattering analyses. The research further explores radar cross-section (RCS), imaging, and spectral properties using advanced THz technologies, including resonant tunneling diodes (RTDs) operating at 250 GHz and THz time-domain spectroscopy (THz-TDS) for frequencies exceeding 250 GHz. High-resolution imaging, utilizing a 450 GHz bandwidth, captures intricate anatomical features of both real and 3D-printed bees, showcasing the potential of THz technology for detailed environmental monitoring. Finally, simulations at 300 GHz assess the dosimetry and feasibility of non-invasive, continuous monitoring approaches based on the heterogeneous honey bee model.

Open access: https://www.nature.com/articles/s41598-025-91630-8

--

13.56 MHz RFID Module – From Application to Process Modelling and Effects on Human Health

Constantinescu, C. *et al.* (2025). 13.56 MHz RFID Module – From Application to Process Modelling and Effects on Human Health. In: Vlad, S., Roman, N.M. (eds) 9th International Conference on Advancements of Medicine and Health Care Through Technology. MEDITECH 2024. IFMBE Proceedings, vol 130. Springer, Cham. doi: 10.1007/978-3-031-95671-3_22

Abstract

RFID (Radio Frequency Identification) is a technology very used in the last few years due to its reliability and ease of use. The applications are present in a large range of frequencies, but this paper is focused on a13.56 MHz module which is usually combined with an Arduino board. First, a simple application for student monitoring capable of storing information about each student and the time he/she checked in and out of class is presented. The advantages of such a system are highlighted and the components used to construct this system are described. The next step of this paper is to determine how the RFID module functions by numerically modelling it with and without the tags and cards near it, attempting also to show how the positioning of the tags influences the characteristics of the active tag's antenna. The last step of the study is centered on determining the influence of such a tag near human tissues and concludes by determining if it is safe or not to use an RFID for long periods of time.

https://link.springer.com/chapter/10.1007/978-3-031-95671-3_22

--

Greater prevalence of symptoms associated with higher exposures to mobile phone base stations in a hilly, densely populated city in Mizoram, India

Sailo L, Laldinpuii, Zosangzuali M, Weller S, Varte CL, Tochhawng L, McCredden JE, Zothansiama. Greater prevalence of symptoms associated with higher exposures to mobile phone base stations in a hilly, densely populated city in Mizoram, India. Electromagn Biol Med. 2025 Jun 8:1-20. doi: 10.1080/15368378.2025.2513900.

Abstract

Members of the scientific community and the general public are raising concerns about the potential health and environmental effects of radio-frequency electromagnetic fields (RF-EMF) for those living nearby mobile phone base stations (MPBS). This study examined the impact of RF-EMF (900-1900 MHz) on symptoms spanning four health categories: mood-energy, cognitive-sensory, inflammatory, and anatomical issues. A questionnaire identifying health symptoms within these categories, was given to 183 highly exposed and 126 reference residents, matched on demographics. While years of residing near the MPBS influenced the prevalence of some symptoms, proximity to the base station and higher levels of exposure (measured using power density) influenced the prevalence of many of the symptoms. A higher proportion of symptoms was found in residents who were either living within 50 meters of a MPBS or who were exposed to power densities of 5-8 mW/m^2, for all four health categories. This relationship between exposure level and symptom prevalence was further influenced by age, daily mobile phone use (over 5 h per day), and lifestyle factors, for certain symptoms. Hierarchical regression analysis revealed that level of exposure (power density) was the only factor contributing to the number of symptoms experienced by residents, for all four health categories. An unexpected finding was that among the more highly exposed residents, the younger individuals (under 40 years) reported more inflammation related issues than older individuals. These results underscore the need to inform

policymakers regarding the benefits of adopting a precautionary approach to potential risks associated with RF-EMF exposures from MPBS.

Plain language summary

Investigating the health effects of man-made electromagnetic fields (RF-EMF) created by telecommunications signals from mobile phone base stations is relevant to people living in cities across the world today. The study was conducted in a hilly, highly populated city in Mizoram, India, where many people live close to and in line of sight of the masts on telecommunications towers. A survey was given to residents in their homes, asking about what health symptoms they were experiencing across a range of health categories (mood-energy, cognitive-sensory, inflammatory, and anatomical). At the same time, the level of RF-EMF in their lounge room was measured. The symptoms reported by people living closer to mobile phone base stations (less than 300 m) were compared with those from people living further away (more than 400 m). More people who lived closer to base stations reported health symptoms in all of the health categories investigated. Relatively fewer people who lived further away reported symptoms. Other factors such as age, high mobile phone use (more than 5 h/day) and smoking and drinking also influenced this outcome, for some of the symptoms. The most significant contributor to the number of symptoms reported by residents was the strength of RF-EMF to which they were exposed in their home. A surprising result was that younger people up to 40 years old showed more inflammatory conditions that were related to higher exposures than older people (such as headache, allergy and chest pain). These health effects of RF-EMF should be heeded by those responsible for the installation of mobile phone base stations in cities.

Study implications

"The maximum recorded power density measured at any dwelling was 7.2 mW/m^2. Most of the measurements taken in dwellings close to the base stations () exceeded the safety limits suggested by the Bioinitiative Report 2012. However, all measured values were significantly lower than both the current ICNIRP whole body exposure public limits of 4500 mW/m^2 (900 MHz) to 10,000 mW/m^2 (2000 MHz and higher frequencies) as well as the present Indian Standard of 450 mW/m^2 (Saravanamuttu et al. 2015). These standards for public RF-EMF exposures are predominantly focused on the mitigation of “known” harmful thermal effects, but do not comprehensively cover a range of biological consequences that are intrinsically linked to non-thermal biological responses and health, especially for those who are continuously being exposed to this man-made radiation without choice or informed consent. In this context, the ALARA principle (As Low As Reasonably Achievable), which has widely been used with ionising radiation, has been recommended (Leach and Bromwich 2018).

Table 3 reveals that symptom prevalence was associated with duration of exposure. In toxicology research, the dose of a toxicant is understood to incorporate both intensity and duration of exposure (Tsatsakis et al. 2018).

The ICNIRP guidelines set safety limits based on exposure intensity, averaged over 6 or 30 minutes. Therefore, they do not factor in cumulative doses occurring over time in the real world. Laboratory studies are mostly restricted to timescales of minutes to weeks. Within studies using the longer of these timescales, biphasic effects have been observed (where effects are positive in the short term but then return to baseline as exposure duration increases and become negative with even longer exposure times) suggesting very short -term protective effects such as immune system priming, but detrimental

effects after longer exposures (e.g., Fesenko et al. 1999). While adaptive responses have been suggested for long term exposures (Vijayalaxmi et al. 2014), the longer timescales of laboratory experiments show more negative effects. However, there are very few long-term laboratory studies.

Epidemiological data, such as that presented in this study, measures exposures over years and thus has been more able to reveal detrimental, cumulative effects that occur over time. Epidemiological research and intensity x time calculations need to be factored in when setting safety limits for populations who are continuously exposed over a lifetime.

While this study sampled only a moderate sized population from the city of Aizawl, the results can be used to inform public policy on this matter. Our results add to the understanding of health effects related to chronic exposure to RF-Fields emitted from MPBS. Risk management does not require definitive conclusions. Early warnings such as the results of this study provide adequate evidence for policy makers to act, including requiring industry to seek solutions (Gee 2009; Leach et al. 2024). The study results indicate that exposure to RF-EMF at levels lower than the Government recommended general public “safety” threshold can still lead to significant health challenges. The study revealed that the level of MPBS RF-EMF exposure was a primary contributor to the elevated prevalence of a range of health symptoms observed among residents in close proximity to MPBS. Concerning results that have revealed more inflammatory symptoms in resident younger than 40 years living close to MPBS require follow up.

Following these important findings, it is crucial to critically reassess current public RF exposure safety limits in order to include the growing evidence of non-thermal biological impacts of RF-EMF emitted by MPBS over years. The study results emphasize the importance of establishing exposure limits that are based on biological factors, other than heating, to minimize exposure and address the long-term potential for serious health consequences; i.e., “An independent re-evaluation of RFR exposure limits based on the scientific knowledge gained over the past 25 years is needed and is long overdue” (Belyaev et al. 2022)."

https://www.tandfonline.com/doi/10.1080/15368378.2025.2513900

--

Trends in Malignant and Benign Brain Tumor Incidence and Mobile Phone Use in the U.S. (2000–2021): A SEER-Based Study

My note: The age-standardized incidence rate for benign (i.e., non-malignant) brain and other nervous system tumors was 9.56 per 100,000 in 2004 and 18.48 in 2021, a 93% increase. Yet the authors of this paper misleadingly claimed contrary to their own result (APC = 1.9, p = .00003): "The incidence rates for benign tumors (primarily meningiomas) were also stable, but did increase slightly over this period [44]. This is unlikely to be due to mobile phone use." The authors' assumption that this substantial increase in non-malignant brain tumors was caused simply by changes in screening practices and tumor classification criteria should be challenged.

Analysis of the SEER data also indicates that the trend for glioblastoma (GBM) varied by age group. Although it was quite stable for younger age groups, the GBM age-standardized incidence rate for seniors 65 years of age and older was 11.97 per 100,000 in 2001 and 13.66 in 2021, a 14% increase.

Since 1,000 hours of mobile phone use may be necessary to initiate a brain tumor and 20-30 years for a tumor to be diagnosed, continued surveillance is essential.

For more information see Brain Tumor Rates Are Rising in the US: The Role of Cellphone & Cordless Phone Use.

Zhang L, Muscat JE. Trends in Malignant and Benign Brain Tumor Incidence and Mobile Phone Use in the U.S. (2000–2021): A SEER-Based Study. *International Journal of Environmental Research and Public Health*. 2025; 22(6):933. doi: 10.3390/ijerph22060933.

Abstract

(1) Background: There has been an ongoing concern for several decades that radiofrequencies emitted from mobile phones are related to brain cancer risk. We calculated temporal trends in brain cancer incidence rates in adults and children and compared them to mobile phone subscription data over the same time period.

(2) Methods: We analyzed the Surveillance, Epidemiology and End Results (SEER 22) cancer database between 2000 and 2021. Age-standardized incidence rates (ASR) per 100,000 people were calculated and the annual percentage change (APC) for malignant and benign brain cancer and vestibular schwannomas (acoustic neuromas of the 8th cranial nerve) was established. The total number of mobile phone subscriptions in the United States was plotted for the period 1985–2024.

(3) Results: The APC for adolescents and adults was −0.6 (p = 0.0004) for malignant tumors, −0.06 (p = 0.551) for temporal lobe tumors, and **1.9 (p = 0.00003) for benign tumors**. The APC for benign acoustic neuroma was 0.09 (p = 0.8237), suggesting that mobile phone use is unlikely to be associated with this tumor type. There was a 1200-fold increase in the number of cell phone subscriptions during this period.

(4) Conclusions: These findings suggest that mobile phone use does not appear to be associated with an increased risk of brain cancer, either malignant or benign.

Open access paper: https://www.mdpi.com/1660-4601/22/6/933

--

5G Radio-Frequency-Electromagnetic-Field Effects on the Human Sleep Electroencephalogram: A Randomized Controlled Study in CACNA1C Genotyped Volunteers

Sousouri G, Eicher C, D' MariaAngelo R, Billecocq M, Fussinger T, Studler M, Capstick M, Kuster N, Achermann P, Huber R, Landolt H-P. 5G Radio-Frequency-Electromagnetic-Field Effects on the Human Sleep Electroencephalogram: A Randomized Controlled Study in CACNA1C Genotyped Volunteers. NeuroImage, 2025. doi: 10.1016/j.neuroimage.2025.121340.

Abstract

Background The introduction of 5G technology as the latest standard in mobile telecommunications has raised concerns about its potential health effects. Prior studies of earlier generations of radiofrequency electromagnetic fields (RF-EMF) demonstrated narrowband spectral increases in the

electroencephalographic (EEG) spindle frequency range (11-16 Hz) in non-rapid-eye-movement (NREM) sleep. However, the impact of 5G RF-EMF on sleep remains unexplored. Additionally, RF-EMF can activate L-type voltage-gated calcium channels (LTCC), which have been linked to sleep quality and EEG oscillatory activity.

Objective This study investigates whether the allelic variant rs7304986 in the CACNA1C gene, encoding the α1C subunit of LTCC, modulates 5G RF-EMF effects on EEG spindle activity during NREM sleep.

Methods Thirty-four participants, genotyped for rs7304986 (15 T/C and 19 matched T/T carriers), underwent a double-blind, sham-controlled study with standardized left-hemisphere exposure to two 5G RF-EMF signals (3.6 GHz and 700 MHz) for 30 min before sleep. Sleep spindle activity was analyzed using high-density EEG and the Fitting Oscillations & One Over f (FOOOF) algorithm.

Results T/C carriers reported longer sleep latency compared to T/T carriers. A significant interaction between RF-EMF exposure and rs7304986 genotype was observed, with 3.6 GHz exposure in T/C carriers inducing a faster spindle center frequency in the central, parietal, and occipital cortex compared to sham.

Conclusion These findings suggest 3.6 GHz 5G RF-EMF modulates spindle center frequency during NREM sleep in a CACNA1C genotype-dependent manner, implicating LTCC in the physiological response to RF-EMF and underscoring the need for further research into 5G effects on brain health.

Open access paper: https://www.sciencedirect.com/science/article/pii/S105381192500343X

--

Exposure to radiofrequency electromagnetic fields and IARC carcinogen assessment: Risk of Bias preliminary literature assessment for 10 key characteristics of human carcinogens

Simkó M, Repacholi MH, Foster KR, Mattsson MO, Croft RJ, Scarfi MR, Vijayalaxmi. Exposure to radiofrequency electromagnetic fields and IARC carcinogen assessment: Risk of Bias preliminary literature assessment for 10 key characteristics of human carcinogens. Mutat Res Rev Mutat Res. 2025 May 27;796:108545. doi: 10.1016/j.mrrev.2025.108545

Highlights

• We analyzed if the ten key characteristics (KCs) of human carcinogens according to IARC are influenced by RF-EMF exposure.
• We reviewed 159 articles by extracting relevant exposure and experimental data.
• A risk of bias (RoB) analysis was conducted using 6 criteria.
•There is a strong negative association between study quality and the likelihood of reporting effects of RF-EMF exposures.
• The heterogeneity and overall poor study quality suggest the need for high-quality studies.

Abstract

This is the first assessment of evidence needed to determine whether exposure to radiofrequency electromagnetic fields (RF-EMF) exposures, below the levels recommended in the ICNIRP (2020) guidelines, can influence any of the ten key characteristics (KCs) of human carcinogens developed by the

International Agency for Research on Cancer (IARC). We define the 10 KCs and their relevance to carcinogenesis; review in vivo and in vitro studies relevant to the KCs; and conduct a risk of bias (RoB) analysis using 6 criteria. We did not include KC studies on genotoxicity or oxidative stress since Romeo et al. (2024) and Meyer et al. (2024) recently published relevant systematic reviews, but note their respective conclusions. From the other 8 KCs we identified 119 in vitro and 40 in vitro measurements of in vivo studies through 30 June 2023, with 38 % reporting statistically significant effects of exposure. We identified a strong association between the quality of study and outcome, with those meeting more RoB criteria less likely to report statistically significant effects. Effects were reported over the entire frequency range, exposure levels, and biological endpoints with no apparent pattern of exposure parameters resulting in effects. Only KC10 (alters cell proliferation, cell death or nutrient supply) has sufficient studies to analyse, but the other KCs had few studies and diverse endpoints. A few relatively high-quality positive studies require follow-up through additional targeted studies. The heterogeneity and overall poor study quality suggest the need for high-quality studies on these endpoints, preferably adhering to standards such as the Organization for Economic Co-operation and Development [28].

Limitation of the review

A significant limitation of the present review is that it was not a PRISMA-compliant systematic review or scoping review. However, the present review did not fully adhere to the standard guidelines for a scoping review, although our approach bore some similarities to the approach adopted in Peters et al. [31]. It is not feasible to adjust the present review (eight reviews in total) to conform to such guidelines. This limitation is acknowledged, and the necessity for comprehensive scoping reviews and, where applicable, SRs, is emphasized.

Conclusions

This review aimed to assess the extent and adequacy of experimental data bearing on whether exposure to RF-EMF could influence any of the 10 IARC key characteristics of human carcinogenesis. Since two PRISMA-compliant SRs on the KCs genotoxicity and oxidative stress were recently published [35], [23], this present review was limited to the 8 remaining KCs for which no systematic reviews have been conducted so far. It became apparent that the present database of in vitro and in vivo studies was so diverse and scattered in their quality that helpful outcomes of SRs on most of the KCs would be unfeasible, and any conclusions from such reviews would have a very low confidence. The two PRISMA-compliant SRs as well as previous comprehensive reviews on in vitro and in vivo genotoxicity studies (e.g. [46], and our review of studies on the 8 KCs, have all reached the same conclusion: there is clearly a need for much higher quality RF-EMF bioeffects studies on these KCs.

There are, however, a few statistically significant results in the highest quality studies (Tier 1 studies in Romeo et al. [35] and studies meeting 5 or 6 RoB criteria (present study). These deserve close examination, and replication if warranted by stronger studies, preferably done under GLP [28], or are otherwise compliant with OHAT recommendations.

As a final comment, it is not useful for systematic reviews to examine hundreds of papers on a given topic, only to find that the overall quality of the large majority of the studies is too low to permit conclusions with a high level of confidence.

Open access paper: https://www.sciencedirect.com/science/article/pii/S138357422500016X

--

Does Electromagnetic Pollution in the ART Laboratory Affect Sperm Quality? A Cross-Sectional Observational Study.

Baldini GM, Lot D, Ferri D, Montano L, Tartagni MV, Malvasi A, Laganà AS, Palumbo M, Baldini D, Trojano G. Does Electromagnetic Pollution in the ART Laboratory Affect Sperm Quality? A Cross-Sectional Observational Study. Toxics. 2025 Jun 18;13(6):510. doi: 10.3390/toxics13060510.

Abstract

In recent decades, exposure to electromagnetic fields (EMFs) generated by standard devices has raised concerns about possible effects on reproductive health. This cross-sectional observational study examined the impact of EMFs on sperm motility in a sample of 102 healthy males aged 20-35 years in the IVF laboratory. Semen samples were exposed to different sources of EMF for one hour, and motility was assessed immediately thereafter. The results showed a significant reduction in progressive sperm motility after exposure to EMFs generated by mobile phones and Wi-Fi repeaters in the laboratory. In contrast, other equipment showed no significant effects. The study demonstrated a statistically significant reduction in progressive sperm motility following in vitro exposure to electromagnetic fields (EMFs) emitted by mobile communication devices and wireless local area network access points. Conversely, other electromagnetic emitting devices evaluated did not elicit significant alterations in this parameter. These findings suggest a potential negative impact of specific EMF sources on semen quality, underscoring the necessity for further comprehensive research to elucidate the clinical implications and to develop potential mitigation strategies aimed at reducing risks to male reproductive health. This study discourages the introduction of mobile phones in IVF laboratories and recommends positioning Wi-Fi repeaters on the ceiling.

Excerpt

- Group 3—iPhone Cell Phone. The samples were exposed to radiation emitted by an Apple iPhone 12 mobile phone (output power: 0.1 W) (Apple Computer, Cupertino, CA, USA). The device was kept at a distance of 10 cm from the semen sample for 1 h.
- Group 4—Ubiquiti Wi-Fi Repeater. The samples were exposed to EMFs emitted by a Ubiquiti UniFi 6 long-range Wi-Fi repeater (Ubiquiti, 685 Third Avenue, New York, NY, USA), utilising 2.4 GHz and 5 GHz Wi-Fi technology with an emission power of approximately 20 dBm (decibel milliwatts), which corresponds to approximately 100 mW (milliwatts). As in all other cases, the sample was placed at a distance of 10 cm for 1 h.

Open access paper: https://www.mdpi.com/2305-6304/13/6/510

--

A novel approach for assessments of radiofrequency electromagnetic fields exposure in buildings near telecommunication infrastructure

de F H Silva RQ, Rodrigues MEC, Pinheiro FSR, da Silva GS, da C Muniz M, Pinto LS, Mendonça HB, de Sousa VA Jr. A novel approach for assessments of radiofrequency electromagnetic fields exposure in

buildings near telecommunication infrastructure. Sci Total Environ. 2025 Jun 25;992:179853. doi: 10.1016/j.scitotenv.2025.179853. Epub ahead of print. PMID: 40570395.

Highlights

- Determines the worst-case exposure in buildings directly exposed to RF-EMF.
- Enables large-scale assessments by limiting specific measurement locations.
- Evaluates all buildings in a city using the proposed approach.
- Novel RF-EMF exposure results in vertical dwellings.
- Unprecedented discussions on recommendations for exposure to RF-EMF in buildings.

Abstract

This paper proposes to complement the current regulatory agencies' methodology for evaluating exposure to Radiofrequency Electromagnetic Fields (RF-EMF) in buildings under the direct incidence of emissions from Base Station (BS) antennas. The key contribution is the refinement of measurement point selection within buildings, ensuring that assessments more accurately capture exposure levels. The approach employs technical criteria for selecting target buildings, considering the location of the BSs and the configuration of the surrounding antenna structures. The proposed approach was applied to measurements in four buildings in the city of Natal, the capital of Rio Grande do Norte, located in the Northeast region of Brazil. The results show electric field intensity peaks up to 17.40 times higher and averages up to 14.13 times higher than values obtained from measurements conducted at ground level, such as those carried out by the National Telecommunications Agency (ANATEL). The highest exposure rates reached 82.27% and 59.43% of the limits established by the International Commission on Non-Ionizing Radiation Protection (ICNIRP) for the frequency bands of FM radio and mobile telephony, respectively. Our proposal can improve and complement the normative guidelines for RF-EMF exposure assessment, providing more representative evaluations of exposure in indoor building environments near modern telecommunication infrastructure.

This paper proposes to complement the current regulatory agencies' methodology for evaluating exposure to Radiofrequency Electromagnetic Fields (RF-EMF) in buildings under the direct incidence of emissions from Base Station (BS) antennas. The key contribution is the refinement of measurement point selection within buildings, ensuring that assessments more accurately capture exposure levels. The approach employs technical criteria for selecting target buildings, considering the location of the BSs and the configuration of the surrounding antenna structures. The proposed approach was applied to measurements in four buildings in the city of Natal, the capital of Rio Grande do Norte, located in the Northeast region of Brazil. The results show electric field intensity peaks up to 17.40 times higher and averages up to 14.13 times higher than values obtained from measurements conducted at ground level, such as those carried out by the National Telecommunications Agency (ANATEL). The highest exposure rates reached 82.27% and 59.43% of the limits established by the International Commission on Non-Ionizing Radiation Protection (ICNIRP) for the frequency bands of FM radio and mobile telephony, respectively. Our proposal can improve and complement the normative guidelines for RF-EMF exposure assessment, providing more representative evaluations of exposure in indoor building environments near modern telecommunication infrastructure.

8. Conclusions

This study presented a novel approach for assessing RF-EMF exposure in buildings directly exposed to emissions from BS antennas, in addition to criteria for selecting target buildings and determining the precise location of the potential worst-case exposure scenario within these structures. The measurement methodology conformed to internationally recognized standards (ICNIRP, 2020) and Brazilian guidelines for evaluating exposure to electromagnetic fields from radiofrequency services (ANATEL, 2023).

Applying the proposed approach for selecting target buildings resulted in obtaining an electrical field intensity peak value higher than those of studies found in the literature that carried out indoor measurements, in addition to presenting values close to those recorded by works that carried out outdoor measurements. These data were acquired from measures over 30 min, as recommended by ICNIRP (2020). This provides a time optimization in exposure assessment compared to approaches that select measurement locations randomly and/or perform measurements over 24 h. Furthermore, this approach provides a more accurate evaluation than methods that use measurements with 6 min or less.

The measurements conducted at the target floor yield average and peak electric field intensity values lower than ICNIRP (2020) exposure limits but up to 17.40 times higher than those obtained at the ground level. These findings indicate the necessity for specific investigations for the discussed exposure scenario and the importance of improving regulatory agencies' methods to consider the complexities and particularities of modern RF-EMF exposure contexts.

In addition, the proposed approach can be improved using geographic and structural datasets on buildings of the analyzed region, enabling the automation of the selection process of BSs and target buildings, as described in Sections 4.1 Target BSs selection, 4.2 Target buildings selection, respectively. Using methods for measuring in the external part of the façade of target buildings, such as drones, can reduce the number of authorizations required to carry out measurements, making more measurements feasible.

Finally, future work aims to conduct measurements in all 22 buildings in Natal and establish partnerships with other institutions to apply the methodology in different cities and countries, promoting a more comprehensive analysis of the exposure scenario in apartments and analyzing reproducibility.

https://pubmed.ncbi.nlm.nih.gov/40570395/

--

Effect of elevation on cumulative radiofrequency exposure from multiple communication towers

Osei S, Quarshie E, Azah CK, Fuseini A-R, Dogbey R, Deatanyah P, Hagan GB, Hodasi JAM, Sam F, Amoako JK. Effect of elevation on cumulative radiofrequency exposure from multiple communication towers, *Radiation Protection Dosimetry*, 2025. doi: 10.1093/rpd/ncaf068.

Abstract

A densely populated place like a public university needs good internet and communication connectivity for effective academic work. As such, University campuses in Ghana are inundated with communication antennas. This study investigated how radiofrequency (RF) power density levels are affected by the elevations of different floors of high-rise buildings of a public university. A spectrum analyser coupled to a log-periodic antenna was used. The RF power density decreased from the ground floor to the third

floor and only increased to maximum levels on the fourth floor. The variation across different floors indicates the influence of elevation on the measured EMF levels. The 900 MHz band produced the highest power density of 1.16E-03 W/m^2 on the last (fourth) floor, suggesting that communication applications in the 900 MHz band are the most used by the university community.

https://academic.oup.com/rpd/advance-article/doi/10.1093/rpd/ncaf068/8177309

--

Evaluation of Personal Radiation Exposure from Wireless Signals in Indoor and Outdoor Environments

Ruijie P, Sali A, Li L, Mohyedin MZ. Qahtan S. Evaluation of Personal Radiation Exposure from Wireless Signals in Indoor and Outdoor Environments. IEEE Access, doi: 10.1109/ACCESS.2025.3579085.

Abstract

With the development of wireless technology, the public is exposed to electromagnetic fields (EMF), which has led to concerns about the potential health effects of EMF exposures. This paper aims to evaluate personal EMF exposures from wireless signals in indoor and outdoor micro-environments in Malaysia. According to the influencing factors, four different types of micro-environments are selected. A radiation exposure meter called ExpoM-RF 4 is used to measure the electric field strength across these micro-environments. From the measurement campaigns, three machine learning (ML) techniques are simulated to model the Electric Field Strength in each micro-environment. The ML techniques are Fully connected neural network (FCNN), eXtreme Gradient Boosting (XG Boost), and Linear Regression (LR) to predict the RMS and Maximum radiation exposure. From the ML models, Total Emission Ratio (TER), Root Mean Square Error (RMSE) and Coefficient of Determination (R2) are evaluated to measure the performance of ML. By comparison, it is found that LR performs well with single and simple data set, while XG Boost and FCNN demonstrate superior capabilities in handling multiple types of data sets. The FCNN model provides the most accurate predictions, particularly in urban and suburban areas where extreme values are observed. Finally, the measured data and the predicted radiation exposure levels are compared against public exposure limit by International Commission on Non-Ionizing Radiation Protection (ICNIRP), Malaysian Communications and Multimedia Commission (MCMC) and Federal Communications Commission (FCC). The results demonstrate that typically personal radiation exposure is lower than the exposure limit (61.4 V/m), which is similar to the most research results. However, in areas with dense population and numerous base stations, the maximum exposure can approach 56.7365 V/m (measured data), which is close to the exposure limit.

CONCLUSION

LR has better prediction results under single and simple data sets, while XG Boost and FCNN have stronger analysis capabilities for multiple types of data sets. In addition, FCNN predicts best in the presence of extreme values and analyzing large and complex data.

By comparing and analyzing the measured data and predicted values across different micro-environments, it is observed that the highest levels of personal radiation exposure typically occur in outdoor urban areas, which is characterized by high population density, high concentration of base stations, and close proximity to these stations. In contrast, park areas with dense vegetation exhibit

significantly lower personal radiation exposure. The dense trees act as natural attenuators, absorbing and scattering the electromagnetic waves, which reduces their intensity. Indoor environments generally exhibit lower electromagnetic field strengths compared to outdoor environments. This can be attributed to structural shielding provided by building materials and fewer high-power sources.

Typically, Electric Field Strength is much lower than the international exposure limits, which is similar to the most research results [2, 6, 13, 14, 16]. However, in areas with dense population and base stations, the maximum value of Electric Field Strength would increase at some point, even close to the exposure limit.

Open access paper:
https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=11031433&isnumber=6514899

--

The use of different exposure metrics in the research about the health impacts of electromagnetic fields

European Research Cluster on EMF and Health (CLUE-H). The use of different exposure metrics in the research about the health impacts of electromagnetic fields. 2024, Policy brief, 1: 1-8.

No abstract.

Excerpts

Figure 5. Near-field sources contribute most to the RF-EMF energy absorbed by the human body. However, the contribution of individual near-field sources depends on their positioning with respect to the investigated organ or part of the body. The mean overall cumulative dose for whole-body was calculated at 0.29 J/kg/day and for brain it was 0.81 J/kg/day [1].

For RF-EMF, well-established effects include tissue heating, microwave hearing for highly pulsed radiation (e.g., from a radar), and tissue stimulation (e.g., from contact currents). Accordingly, different metrics are specified in the regulations to prevent these effects from happening. For instance, thermal effects are only a problem for health if a certain threshold is exceeded. Below this threshold, temperature rise is not expected to cause health effects.

In RF-EMF research, thermal effects are of limited interest because they are well understood. However, research has provided indications for biological effects below the thermal threshold such as effects on brain physiology or oxidative balance. Such biological effects, which per se are not health effects, could be the consequences of unknown biological mechanisms or could occur due to subtle warming of the tissue below the thermal damage threshold. Historically, epidemiology has been investigating health effects without prior knowledge of the underlying disease mechanism by comparing people who are exposed to a variable extent to the agent of interest. In this case a common approach for complex exposure situations is the time weighted average (TWA), i.e., exposure levels in different situations (e.g., at home, at work, during commuting) are averaged taking into account the time spent in these situations. This approach has also been applied in RF-EMF research dealing with far field exposures. However, it is not suitable for combining near and far field exposures since different metrics are used for

these two types of exposure. Thus, a cumulative dose metric was introduced in RF-EMF research a few years ago. In an approach to the TWA concept, SAR for various exposure situations (e.g., mobile phone call, WiFi access point exposure, etc.) is multiplied with the corresponding exposure duration to obtain a cumulative RF-EMF dose, often expressed per day (J/kg/day).

Calculation of cumulative dose in RF-EMF epidemiology allows combining different exposure situations into one metric. It considers magnitude and duration of each exposure situation and is based on the same philosophy as a time weighted average. In principle, cumulative dose refers to a linear-no-threshold model but is actually correlated to most other plausible effect models such as time spent above a certain threshold. It is a conservative approach, since it considers the possibility that long-term exposure to low levels might affect health, which is a common concern of parts of the population in relation to environmental RF-EMF exposure. It may also be helpful for risk communication as it enables to compare the contribution of various RF-EMF exposure situations to a combined metric of the absorbed RF-EMF.

The use of a cumulative dose metric in research should not be mistaken as an indication or proof that cumulative exposure to very low levels can be harmful to health. It just serves as the currently best metric to analyse if there could be effects on health, outside of a known biological mechanism.

In conclusion, the existence of various exposure measures in the field of RF-EMF research on health reflects the involvement of different biophysical concepts and exposure situations. Further, different metrics are used for different purpose:

• For biological research, physical quantities that best represent an underlying biophysical mechanism are usually the preference (e.g., SAR value, internal electrical field).
• For observational research aiming to explore a yet unknown effect on health, preference may be given to metrics that combine similar sources (e.g., time-weighted average, cumulatively absorbed dose) and are seen as superior in capturing exposures experienced over a long time period.).
• For regulation purposes, the suitable metric depends on the exposure situation (e.g., external electric field strength or power density for far field sources or spatially averaged SAR for localized near field exposures).
• For risk communication with the public, metrics which are intuitively understood, are considered most useful (e.g., fraction of regulatory limit).

There is a Council Recommendation [2] that is currently being amended due to the fact that new measurement methods have emerged. The work of CLUE-H contributes to improving our knowledge base on measurement methods and limit values, which will be useful for future updates of the Recommendation.

Open access: https://www.emf-health-cluster.eu/wp-content/uploads/2024/06/CLUE-H_First_Policy_Brief_DEF_approved.pdf

--

Auto-Induced Downlink Radiofrequency Electromagnetic Field Exposure at 3.5 GHz With Focusing Near the Head

Herssens H, Thielens A. Auto-Induced Downlink Radiofrequency Electromagnetic Field Exposure at 3.5 GHz With Focusing Near the Head. *IEEE Access*, vol. 13, pp. 56659-56670, 2025, doi: 10.1109/ACCESS.2025.3555388.

Abstract

The auto-induced downlink exposure at 3.5 GHz is evaluated in networks where the power density is focused at the user equipment (UE), leading to a highly localized exposure. To achieve this focusing, various precoding techniques can be applied. Two of these are compared using Finite-Difference Time-Domain simulations. For each technique, three different exposure scenarios are considered: UE located next to the ear (phone call), UE located in front of the eyes, and the nose (video call). The exposure is evaluated in terms of the localized specific absorption rate using a female and a male phantom. We find that the exposure highly depends on the location of the UE and precoding technique. According to the International Commission on Non-Ionizing Radiation protection (ICNIRP)'s guidelines, this exposure should be normalized to the maximum incident power density. In this work, we suggest normalization strategies of this highly focused exposure and compare these. We show that the choice of normalization can create a situation where the ICNIRP basic restrictions can be exceeded while complying with the reference levels. This is important because compliance with the reference levels should ideally ensure compliance with the basic restrictions.

Open access paper:

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=10943211&isnumber=10820123

--

In-Situ Measurements of Radiofrequency Electromagnetic Fields Measurements Around 5G Macro Base Stations in the UK

Calderon C, Addison D, Peyman A. In-Situ Measurements of Radiofrequency Electromagnetic Fields Measurements Around 5G Macro Base Stations in the UK. Bioelectromagnetics. 2025 Jul;46(5):e70012. doi: 10.1002/bem.70012.

Abstract

Radiofrequency (RF) electromagnetic field spot measurements were performed in line-of-sight to 56 active 5G macro base stations across 30 publicly accessible locations in the United Kingdom (UK). Four different exposure scenarios were assessed: background (no traffic instigation), streaming videos, downlink speed test, and extrapolation of SS-RSRP decoder measurements. Power density measurements across the 420 MHz-6 GHz frequency range were also performed at each site to assess the total exposure from various RF sources in the environment. Both total RF and 5G specific power density levels were found to be well within the 1998 ICNIRP public reference levels, even when extrapolating to worst-case scenario (≤ 5%). 4G downlink was the dominant contributor to total RF exposure, with 5G contributing on average less than 10%. No statistically significant difference was observed between beamforming and non-beamforming sites. Streaming did not seem to contribute materially to exposure levels, suggesting that background measurements are a good representation of typical downlink exposure at current urban and suburban 5G sites.

Open access: https://onlinelibrary.wiley.com/doi/10.1002/bem.70012

--

Exposure to 26.5 GHz, 5G modulated and unmodulated signal, does not affect key cellular endpoints of human neuroblastoma cells

Sannino A, Allocca M, Scarfì MR, Romeo S, Peluso V, Panariello G, Schettino F, Chirico G, Zeni O. Exposure to 26.5 GHz, 5G modulated and unmodulated signal, does not affect key cellular endpoints of human neuroblastoma cells. Sci Rep. 2025 Jul 1;15(1):20614. doi: 10.1038/s41598-025-04834-3.

Abstract

The fifth generation (5G) network is currently being worldwide spread out, raising questions about its potential health impact. The current study aimed to investigate the effects of a 26.5 GHz 5G electromagnetic field on key cellular endpoints of human neuroblastoma cells. A reverberation chamber-based exposure system was designed and realized which allowed the exposure/sham exposure of cell cultures under highly controlled exposure conditions of both electromagnetic and biological parameters. The suitability of the reverberation chambers to host cell cultures was verified by evaluating cell proliferation and cell cycle progression. The effect of 3 h exposure at specific absorption rate of 1.25 W/kg under both continuous wave and 5G modulated signal was evaluated in terms of cell cycle and DNA damage. In the latter case, the exposure was also given in combination with menadione to account for possible cooperative effects. Results showed absence of effects of exposure given alone and in combination with menadione, when both continuous wave and modulated signals were applied at the mentioned exposure level. Further investigations are needed by varying the exposure and biological parameters to strengthen the absence of effects due to 5G signals in the range of millimeter waves.

Open access: https://www.nature.com/articles/s41598-025-04834-3

--

Histomorphometric study of thyroid tissue in juvenile rats exposed to 5G electromagnetic fields

Ramelet M, Ronger L, Al-Salameh A, Pelletier A, Desailloud R, Seewooruttun C. Histomorphometric study of thyroid tissue in juvenile rats exposed to 5G electromagnetic fields. Annales d'Endocrinologie.86(3), 2025. doi: 10.1016/j.ando.2025.101755.

Abstract

Introduction 5G network deployment raises the issue of the health impact of these electromagnetic fields. Their effect on thermal regulation is of a “cold reesponse” type. In rats exposed to 900MHz, vasoconstriction was induced in the tail, with behavioral preference for warmer compartments. The hypothalamo-pituitary-thyroid axis is involved in thermogenesis: in response to cold, thyroid activity increases, stimulating heat production.

Objective To assess thyroid activity after 5G exposure.

Method Ten 3-week-old male Wistar rats were randomized between 2 weeks’ 5G exposure at 3.5GHz and 1.5V/m (5G group; n=5) and no exposure (controls; n=5). After sacrifice, the thyroid gland was harvested for histomorphometry. Thyroid follicle and colloid areas were measured and the thyroid activation index (TAI: ratio of follicle to colloid area) was assessed.

Results Follicle and colloid areas were significantly greater in 5G (2729µm2 vs 2444µm2, P=0.0063; and 1317µm2 vs 1015µm2, P=0.0027 respectively) and TAI was significantly lower (5.62 vs 7.07, P <0.001). These results indicate thyroid hypoactivity.

Discussion There was a 5G effect, with histomorphometry demonstrating thyroid hypoactivity, suggesting mechanisms different from those underlying cold response. Impact of electromagnetic fields on thyroid function was previously reported, but results were heterogeneous and discordant depending on the type of exposure.

Conclusion TSH and thyroid hormone assays are underway to reinforce assessment of thyroid function. Further studies are needed to explore the underlying mechanism.

https://www.sciencedirect.com/science/article/pii/S0003426625000745

--

Protective effects of quercetin against 3.5 GHz RF radiation-induced thyroid dysfunction and oxidative stress in rats

Bektas H, Bese Akgun BB, Cakir S, Dogu S, Ahnas B. Protective effects of quercetin against 3.5 GHz RF radiation-induced thyroid dysfunction and oxidative stress in rats. Electromagn Biol Med. 2025 Jul 8:1-12. doi: 10.1080/15368378.2025.2528732.

Abstract

The global expansion of 5 G communication networks has heightened concerns about the biological effects of high-frequency radiofrequency (RF) radiation, particularly on endocrine organs such as the thyroid gland. This study investigated the effects of 3.5 GHz RF radiation on thyroid hormone levels and oxidative stress markers in male Wistar rats and assessed the potential protective role of quercetin, a natural antioxidant. Twenty-eight rats were randomly assigned to four groups: Sham, RF, Quercetin, and RF + Quercetin. RF exposure was administered at 3.5 GHz (2 W) for 2 hours/day, 5 days/week, for 30 days. Quercetin (20 mg/kg) was administered intraperitoneally. Serum levels of T3, T4, and TSH, as well as thyroid tissue levels of TAS, TOS, GSH, and MDA, were analyzed using ELISA. RF exposure significantly decreased T3 and T4, increased TSH, elevated MDA and TOS, and reduced TAS and GSH levels. Quercetin treatment showed trends toward reversing some of these effects, although not all changes reached statistical significance. SAR simulations confirmed higher energy absorption in the thyroid region (average SAR: 1.128 W/kg). These findings suggest that 3.5 GHz RF radiation may impair thyroid function and redox homeostasis, and that quercetin may exert limited biochemical protection, though further studies are needed to confirm its efficacy. Further long-term molecular studies are warranted to elucidate the mechanisms involved.

Plain Language Summary

With the increasing rollout of 5 G networks, concerns have emerged regarding the potential health impacts of high-frequency radio signals, especially on sensitive organs like the thyroid gland. In this study, researchers explored how repeated exposure to 3.5 GHz RF radiation affects thyroid health in rats, and whether quercetin – a plant-based antioxidant – can offer protection. Rats were exposed to RF radiation for one month, and key hormone and oxidative stress indicators were measured. The results showed that RF exposure disrupted thyroid hormone levels and increased cellular stress. Quercetin

treatment showed some potential in alleviating certain changes, though its effects were not consistent across all measured parameters. Simulations also showed high absorption of RF energy in the thyroid area. These findings raise awareness about the possible biological effects of long-term 5 G exposure and suggest that natural antioxidants like quercetin may offer partial protection. Further research is needed to understand how these findings may apply to human health.

https://pubmed.ncbi.nlm.nih.gov/40625224/

--

The effects of short-term and long-term 2100 MHz radiofrequency radiation on adult rat auditory brainstem response

Er H, Basaranlar G., Derin N., Kantar D, Ozen S. (2025). The effects of short-term and long-term 2100 MHz radiofrequency radiation on adult rat auditory brainstem response. *Open Chemistry*, *23*(1), 20250173. doi: 10.1515/chem-2025-0173.

Abstract

Although mobile phones that work with RFR provide very important benefits for our lives, they may have negative effects. Namely, side effects, such as headaches, sleep disorders, dizziness, lower sperm quality, changes in brain potentials, an increase in oxidative stress levels, and a decrease in antioxidant parameters, have been reported due to mobile phone use. Accordingly, the aim of this research is to investigate the effects of acute and chronic 2100 MHz radiofrequency radiation (RFR) exposure on the auditory brainstem response (ABR) in adult rats. Study groups (*n* = 10 rats): Sh-1: sham for 1 week; Sh-10: sham for 10 weeks; 2100-1: 2100 MHz for 1 week; and 2100-10: 2100 MHz for 10 weeks. RFR groups were applied for 2 h/day (5 day/week) 2100 MHz RFR, whereas sham groups were kept under identical circumstances without RFR. ABR were recorded, and biochemical and ultrastructural examinations in the rat brain were carried out. In the acute RFR group, the latencies of all ABR waves were prolonged compared to the sham group. In the acute RFR group, brain 4-hydroxynonenal, thiobarbituric acid reactive substances, and protein carbonyl content levels increased and catalase and superoxide dismutase activities decreased compared to the acute sham group. Edema in acute RFR group neurons, astrocytes, astrocytic end-feet, and mitochondrial damage in astrocytes were observed. Our data imply that acute exposure to 2100 MHz RFR may have adverse impacts on the auditory system, while chronic exposure with certain rest days has no harmful effects.

Excerpt

In this study, the experimental apparatus and RFR application were identical to those previously reported [5]. However, the RFR frequency, the distance of the rats from the antenna, and therefore, the specific absorption rate (SAR) values were different. In this system, a 2100 MHz radiofrequency generator (Universal Mobile Telecommunication System [UMTS] Simulator 2100 MHz; Everest Company, Adapazari, Turkiye) was used to simulate exposure to UMTS radiation. The electric-field strength measured above the rat's head in the "signal-on" condition was found to be 35.2 V/m. The average SARs for the whole body and brain were 128 mW/kg and 0.27 W/kg, respectively. The computational analysis was conducted utilizing the finite difference time domain method [12]. Published sources were utilized to obtain data on electrical properties, dielectric constant, and conductivity [13,14].

Open access paper: https://www.degruyterbrill.com/document/doi/10.1515/chem-2025-0173/

--

Electromagnetic fields from mobile phones: A risk for maintaining energy homeostasis?

Seewooruttun C, Mai TC, Corona A, Delanaud S, de Seze R, Bach V, Desailloud R, Pelletier A. Electromagnetic fields from mobile phones: A risk for maintaining energy homeostasis? Annales d'Endocrinologie. 86(3). 2025. doi: 10.1016/j.ando.2025.101782.

Abstract

In the world, there is a near ubiquitous presence of a low-intensity radiofrequency electromagnetic field (RF-EMF) radiation, due to telecommunications as mobile phones. However, their rapid expansion raises concerns about possible interaction with biological mechanisms. The RF-EMF safety guidelines recommended limits to protect against the thermal heating, the most recognized effect at high intensity levels with a known biophysical mechanism. Among all the effects studied, the impact of RF-EMF exposure on thermoregulation is one of the most important aspects of this research. This review aims to present the complex relationship between RF-EMF exposure and thermoregulation, at intensity levels below the threshold to produce thermal effects. In fact, most studies showed that RF-EMF exposure at 900MHz seems to elicit physiological and biological effects similar to responses inducing by cold environment in two different rodent models. In this brief review, we will describe the effects and underlying mechanisms induced by RF-EMF exposure at low levels and discuss the potential implications for environmental health and safety.

Conclusion

In modern society, exposure to RF-EMF is very common and nearly impossible to avoid. Despite its major impact in driving technological advancements, it is important to address the growing public concern in regards to RF-EMF exposure. The thermal effects of high-intensity RF-EMF exposure (SAR > 4 W/kg) are well documented in the literature. However, there is now emerging evidence that low-intensity RF-EMF exposure (SAR < 4 W/kg) can also elicit thermoregulatory responses associated with a cold sensation. The present review highlights the behavioural thermoregulation associated with a cold sensation in rodents exposed to 900 MHz. We also provide molecular insights on RF exposure effects on different thermogenic mechanisms. The first results suggest that RF exposure primarily influences WAT browning rather than BAT thermogenesis, as reported by the up-regulation of UCP1 staining and reduced adipocyte size in WAT depots. In contrast, the cold-induced transcriptional changes on BAT thermogenesis were not observed after exposure to RF, although we detected increased plasma levels of noradrenaline, fatty acids in exposed rats. At short term, these adaptive responses do not seem to compromise homeostasis and therefore the health of organisms, but what about the long-term consequences?

With the introduction of 3.5 GHz for 5G and early use of wireless technology by infants, further research is needed to determine these thermoregulatory effects. Investigating the peripheral tail temperature and thermal preference of 5G-exposed rats at different ages could provide valuable insights into their thermoregulatory responses. Additionally, the implication of the thyroid on these responses need to be

more investigated. In fact, the hormones secreted by the thyroid regulate the basic metabolism and the processes implicated in lipolysis in adipose tissues to increase the production of energy in cells and heat by the thermogenesis.

https://www.sciencedirect.com/science/article/pii/S0003426625001015

--

Characterization of the Core Temperature Response of Free-Moving Rats to 1.95 GHz Electromagnetic Fields

Bala N, Croft RJ, McIntosh RL, Iskra S, Frankland JV, McKenzie RJ, Deng C. Characterization of the Core Temperature Response of Free-Moving Rats to 1.95 GHz Electromagnetic Fields. Bioelectromagnetics. 2025 Jul;46(5):e70013. doi: 10.1002/bem.70013.

Abstract

The present study investigated the core body temperature (CBT) response of free-moving adult male and female Sprague Dawley rats, during and following a 3-h exposure to 1.95 GHz radiofrequency electromagnetic fields (RF-EMFs) within custom-built reverberation chambers, using temperature capsules implanted within the intraperitoneal cavity and data transmitted via radiotelemetry. Comparing RF-EMF exposures (at Whole-Body Average-Specific Absorption Rate [WBA-SAR] levels of 0.1, 0.4, and 4 W/kg) to the sham exposed condition, we identified a statistically significant peak increase in CBT after 26 min of RF-EMF exposure at 4 W/kg (+0.49°C), but not in the 0.1 or 0.4 W/kg conditions at the same timepoint. In the last 30 min of the RF-EMF exposure, temperature was significantly increased in both the 4 W/kg (0.62°C) and 0.4 W/kg (0.14°C) conditions, but not 0.1 W/kg, when compared to sham. After 20 min following cessation of exposure, post temperature was still significantly higher in the 4 W/kg condition when compared to the sham (0.37°C), but not in either 0.1 or 0.4 W/kg. Based on our findings, it is apparent that rats can effectively compensate for increased thermal loads of up to 4 W/kg as the maximum temperature rise was substantially lower than 1°C. In addition, the elevated CBT during exposure in the 4 W/kg condition was significantly reduced immediately after exposure cessation, indicating that measures of CBT following RF-EMF exposure cessation may not reflect maximum RF-EMF-mediated changes in the CBT of rats.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.70013

--

The Influence of an Electromagnetic Field at a Radiofrequency of 900 MHz on the Behavior of a Honey Bee

Migdał P, Plotnik M, Bieńkowski P, Berbeć E, Latarowski K, Białecka N, Murawska A. The Influence of an Electromagnetic Field at a Radiofrequency of 900 MHz on the Behavior of a Honey Bee. *Agriculture*. 2025; 15(12):1266. doi: 10.3390/agriculture15121266.

Abstract

The development of wireless technology and the desire to improve communication electromagnetic fields (EMFs) of various frequencies have become common across the honey bee's foraging landscape. There has been discussion for many years about the possible impact of electromagnetic fields on living

organisms. Artificial radio fields emit frequencies ranging from 100 kHz to 300 GHz. The presented research aimed to demonstrate the influence of the radiofrequency electromagnetic field (RF-EMF) with a frequency of 900 MHz on the behavior of honey bees in laboratory conditions. For this experiment, we used wooden cages to house honey bee workers immediately after they emerged. Bee workers were divided into control and experimental groups. Bees in the control group were not exposed to RF fields, while the experimental groups were exposed to 900 MHz electromagnetic fields of different intensities and durations of exposure. Bees' behavior was analyzed with an appropriate computer program. Behavioral analysis of bees was performed immediately after exposure and seven days after exposure. Our research has shown that the radio field (900 MHz) affects the behavior of bees compared to the control group, although not all results are statistically significant. Significant effects were observed seven days after exposure in walking, flight, and individual contact. However, it is worth extending the study to include the impact of an RF-EMF on the expression of genes responsible for bee behavior.

Open access paper: https://www.mdpi.com/2077-0472/15/12/1266

--

Mitigation of 3.5 GHz Electromagnetic Field-Induced BV2 Microglial Cytotoxicity by Polydeoxyribonucleotide

Pachhapure S, Mufida A, Wei Q, Choi J-S, Jang B-C. Mitigation of 3.5 GHz Electromagnetic Field-Induced BV2 Microglial Cytotoxicity by Polydeoxyribonucleotide. *Current Issues in Molecular Biology*. 2025; 47(6):386. doi: 10.3390/cimb47060386

Abstract

Emerging evidence highlights the biological risks associated with electromagnetic fields (EMFs) generated by electronic devices. The toxic effects and mechanisms induced by exposure to EMFs on microglial cells and natural substances that inhibit them are limited to date. Here, we investigated whether exposure to 3.5 GHz EMF radiation, potentially generated by smartphones working in 5G communication or cooking using microwave ovens, affects the growth of BV2 mouse microglial cells and polydeoxyribonucleotide (PDRN), a DNA preparation derived from salmon sperm, inhibits it. Of note, exposure to 3.5 GHz EMF radiation for 2 h markedly inhibited the growth and triggered apoptosis in BV2 cells, characterized by the reduced number of surviving cells, increased genomic DNA fragmentation, increased reactive oxygen species (ROS) levels, and altered phosphorylation and expression levels of JNK-1/2, p38 MAPK, ERK-1/2, eIF-2α, and procaspase-9. Pharmacological inhibition studies revealed that JNK-1/2 and p38 MAPK activation and ROS generation were crucial for 3.5 GHz EMF-induced BV2 cytotoxicity. Of interest, PDRN effectively countered these effects by inhibiting the activation of JNK-1/2, p38 MAPK, and caspase-9, and the production of ROS, although it did not affect eIF-2 phosphorylation. In conclusion, this study is the first to report that PDRN protects against 3.5 GHz EMF-induced toxicities in BV2 microglial cells, and PDRN's protective effects on 3.5 GHz EMF-induced BV2 cytotoxicity are mediated primarily by modulating ROS, JNK-1/2, p38 MAPK, and caspase-9.

Open access paper: https://www.mdpi.com/1467-3045/47/6/386

--

The influence of Wi-Fi on the mesonephros in the 9-day-old chicken embryo

Almášiová V, Andrašková S, Karaffová V, Hudáková P, Molnár J, Tóth Š, Holovská K. The influence of Wi-Fi on the mesonephros in the 9-day-old chicken embryo. Vet Res Commun. 2025 Jun 10;49(4):216. doi: 10.1007/s11259-025-10777-x.

Abstract

The use of wireless devices has increased rapidly in recent times, especially in developed countries. As a result, all living systems are to some extent permanently exposed to this artificial electromagnetic non-ionizing radiation (NIR). These modern devices provide countless benefits to the users, but the disadvantage of their excessive use is the production of electrosmog. This physical pollutant of the environment can be particularly dangerous especially during the developmental period of the individual. The aim of the current study was to elucidate the effect of Wi-Fi radiation on the mesonephros development in the chicken embryo on day 9 of incubation. Continual 9-day application of radiation with a frequency of 2.4 GHz and a power density of 200-500 $\mu W/m^2$ had no adverse effect on the general development of the mesonephros, however moderate diffuse degenerative changes were found in the developing mesonephric corpuscles and tubules. Also congested blood vessels were present in the surrounding interstitium, but no signs of inflammatory infiltrate were detected. In the Wi-Fi group, we also noted a significantly increased number of apoptotic and proliferating cells as well as a significant up-regulation of caspase-1 gene expression. The results indicated that non-ionizing radiation at the frequency and power density used in the study can interfere with the key regulatory mechanisms involved in the normal development of tissues and organs.

https://link.springer.com/article/10.1007/s11259-025-10777-x

--

The proliferation rates of HT-1080 human fibrosarcoma cells can be accelerated or inhibited by weak static and extremely low frequency magnetic fields

Marek B, Nhat D, de Mingo Isabel L, Jason K, Hakki G, Ladislav J, Barnes F. The proliferation rates of HT-1080 human fibrosarcoma cells can be accelerated or inhibited by weak static and extremely low frequency magnetic fields. Frontiers in Public Health. Volume 13 - 2025. doi: 10.3389/fpubh.2025.1535155.

Abstract

Introduction Weak static and low-frequency magnetic fields (MFs) have been hypothesized to influence biological systems through mechanisms involving nuclear spin coupling. This study investigates how such fields modulate the proliferation of HT-1080 fibrosarcoma cells.

Methods HT-1080 cells were exposed in vitro for 4 days to weak MFs with a 10 μT amplitude and frequencies between 12 Hz and 33 Hz, superimposed on a 45 μT static background field. Changes in cell growth, mitochondrial superoxide (O2−), calcium ion (Ca2+) concentrations, and membrane potential were measured.

Results Results revealed that MFs could either increase or decrease fibrosarcoma cell growth in a frequency- and amplitude-dependent manner. Inversions in growth rates were observed near 16.5 Hz, where a 0.5 Hz shift or amplitude changes as small as 250 nT reversed effects relative to controls.

Reversing the static field direction also inverted growth outcomes. Changes in membrane potential, Ca2+, and mitochondrial superoxide levels supported a role for bioenergetic modulation.

Discussion These findings suggest that weak MFs affect cell proliferation through spin-dependent chemical reaction rate changes. The pronounced sensitivity of fibrosarcoma cells compared to normal fibroblasts points to potential therapeutic applications via selective MF-based modulation.

Open access paper: https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1535155

--

Impact of magnetic fields from tablets, laptops, smartphones, and household/leisure magnets on cardiac implantable electronic devices

Kamitani N, Miyazaki A, Tomida S, Shimizu K, Ohira N, Kondo K, Miura H, Koyama D, Tominaga S, Henmi R, Sugiura R, Masui H. Impact of magnetic fields from tablets, laptops, smartphones, and household/leisure magnets on cardiac implantable electronic devices. J Arrhythm. 2025 Jun 30;41(4):e70106. doi: 10.1002/joa3.70106.

Abstract

Background Cardiac implantable electronic devices (CIEDs) activate the magnet response at a magnetic flux density of ≥10 gauss (G), which may cause unintended pacing, leading to discomfort or even severe arrhythmias. Information processing devices have recently incorporated magnets, which may activate the magnet mode in patients with abdominally implanted devices, subcutaneous implantable cardioverter-defibrillators (ICDs), or extravascular ICDs.

Methods We investigated the effects of the magnetic fields generated by information processing devices (tablets, laptops, and smartphones) and household/leisure magnets on 13 models of CIEDs, analyzing their association with magnet mode activation in different manufacturers' CIEDs.

Results The tested magnet materials exhibited a maximum magnetic flux density of 290–1360 G. The magnetic flux density distribution in the information processing devices was as follows: accessory connectors, speakers, cameras, and microphones ($p = 0.0001$). The median activation distances for the magnet mode were 6.5 (range, 4–15), 5 (4–11.3), and 0.01 (activated only when attached; 0–7) mm for tablets and laptops, smartphones, and household/leisure magnets, respectively ($p < 0.0001$). The maximum distance at which the magnetic flux density decreased below 10 G was the longest for tablets and laptop computers at 18 mm.

Conclusion Information processing devices and household/leisure magnets can affect CIEDs when placed in close proximity. Among the devices tested, magnet mode activation did not occur at distances of ≥20 mm. Considering the increasing prevalence of information processing devices and the growing adoption of nonthoracic CIED placements, raising awareness among patients about potential interactions is crucial.

Open access: https://pmc.ncbi.nlm.nih.gov/articles/PMC12209319/

--

SAR Estimations in a Classroom with Wireless Computers

Soares NE, Bulla G, Fernández-Rodríguez CE, de Salles AAA. SAR Estimations in a Classroom with Wireless Computers. J. Microw. Optoelectron. Electromagn. Appl. 24 (02) • 202.
https://doi.org/10.1590/2179-10742025v24i3288526

Abstract

This research provides detailed examination of the peak spatial Specific Absorption Rate (psSAR) in different age students in a classroom with wireless computers. This is motivated by the escalating inclusion of electronic devices in educational settings and the necessity to estimate the implications of this on the overall radio-frequency exposure. Two classrooms with posable realistic human models are simulated. One filled with several 7 years old children and another with several 43 years old adult, each using a laptop. The 1 g and 10 g psSAR are calculated for the head, back and hands. The distances between students are varied and the results compared to one student alone. A small free distance between rows (10 cm) produces significant reduction in psSAR (around 13 dB). Results are less sensitives to changes in lateral distance. While the maximum simulated psSAR values are below the recommended limits, it is observed that, in some classroom arrangements, the psSAR can be substantially increased (e.g., up to 26 dB in the back) comparing to just one student with his laptop. One objective of this study is to provide guidelines for the design of safer classrooms in the context of widespread laptop usage.

Excerpts

"The psSAR simulation in a classroom should consider the interaction between multiple EMF sources and the matter in the environment. While other sources such as smartphones and routers can be present, for simplicity, only the laptop of each student connected in a Wi-Fi network is considered.

The 2.45 GHz band was chosen for computational reasons. Using the 5 GHz band would have increased the mesh by 10-fold.

According to the IEEE 802.11-2020 Standard, the maximum delivered power is 100 mW. This value is used in the simulations. The simulated signals are continuous waves (CW) at the center of the band or CW modulated by a gaussian pulse to cover the whole band."

"The psSAR values obtained in this study, even for the worst case, are below the safety limits recommended by the ICNIRP [2] and IEEE [14]. However, since children can stay in the classrooms for a long time, it is important to observe that long-term exposure to low levels of electromagnetic radiation should also be considered, since they can produce health effects too [1]. Among others, long time exposure can be of serious concern, due to possible biochemical effects not considered in the simulations. Further to that, other EMF sources, such as smartphones, can be used simultaneously in the classrooms and therefore should also be considered in the simulations."

"CONCLUSIONS

The results show that, generally the psSAR decreases as the distance between students' desks increases, as expected. Increased EMF exposure in the classroom were observed when comparing with one student alone, such as up to 4-fold in the hands, 45-fold in the head and 40-fold in the back/dorsal.

Fluctuations in the psSAR plots can be due to the EMF multipath random combinations in amplitude and phase. Also, it is observed that as the distance is increased, the positions of the psSAR are displaced along the body. For instance, in the head, the psSAR is displaced from one ear to the nose and to the other ear. Simulations show that hot spots position varies, since EMF may be increased or diminished when distance is varied, depending on the EMF phase.

Overall, simulations show that if the distance between the chair and the back desk is increased, then the EMF exposure is reduced. E.g., for 5 cm distance increase, the EMF exposure in the back can be reduced around 63 percent and for 50 cm distance increase, the EMF exposure can be reduced around 90 percent.

In order to reduce the EMF health risks, the authors suggest that wireless communications should be avoided in places where people could stay for long time, such as in the classrooms, libraries, offices, homes, etc., and the devices should then be connected using cables (e. g., ethernet or fiber optic), increasing therefore the bandwidth and reducing energy consumption."

Open access paper: https://www.scielo.br/j/jmoea/a/hS9VJvKddrTsTtyqLrVdNZx/?lang=en#

--

Bus-exposure matrix, a tool to assess bus drivers' exposure to physicochemical hazards

Remy VFM, Innocent G, Vernez D, Guseva Canu I. Bus-exposure matrix, a tool to assess bus drivers' exposure to physicochemical hazards. Ann Work Expo Health. 2025 Jun 20:wxaf036. doi: 10.1093/annweh/wxaf036.

Abstract

Swiss bus drivers suffer from musculoskeletal disorders, fatigue, and stress and have an excessive mortality from lung cancer and suicide compared to other workers. However, their occupational exposure is poorly documented. We created a bus-exposure matrix (BEM) to determine occupational exposures to 10 types of physical-chemical hazards for 705 bus models used in Switzerland since 1980. For this, we made a comprehensive bus inventory and review of 50 technical characteristics of each bus model, identified 10 bus models representative of the Swiss bus fleet evolution, and conducted static and dynamic exposure measurement campaigns in the representative buses. The measured values were then extended to the entire fleet using Integrated Nested Laplace Approximation (INLA) models. The choice of predictors and technical bus characteristics included in the models were based on directed acyclic graphs. To demonstrate the usefulness of the BEM as an exposure assessment tool, we used data from the 2022 survey of Swiss bus drivers who listed the bus models they had driven during their careers. The BEM linkage with these bus drivers' histories enabled us to estimate annual exposure to PM10 ratio (-), ultrafine particle ratio (-), whole-body vibration (m/s2), floor vibration (m/s2), equivalent noise (dB(A)), peak noise (dB(C)), high-frequencies electric fields (V/m), low-frequencies magnetic field (μT), low-frequencies electric fields (V/m), and air exchange rate (1/h) of 809 Swiss bus drivers. Historical data assessment from 1985 through 2022 showed that peak noise, high- and low-frequencies electric field levels have increased, while PM10 ratio, ultrafine particle ratio, equivalent noise, whole-body vibration levels, and air exchange rate have decreased. This, first in the world, BEM is an original tool for retrospective exposure assessment that will enable further research in the occupational health of bus drivers.

https://pubmed.ncbi.nlm.nih.gov/40578599/

--

Human achromatic flickers and phosphenes thresholds under extremely low frequency electric stimulations

Bouisset N, Carvallo A, Laporte M, Legros A. Human achromatic flickers and phosphenes thresholds under extremely low frequency electric stimulations. Sci Rep. 2025 Jul 3;15(1):23779. doi: 10.1038/s41598-025-06271-8.

Abstract

Human exposure to extremely low-frequency (< 300 Hz) electric/magnetic fields elicits a stroboscopic visual perception called electro/magneto phosphenes. The induction of phosphenes is the most exhaustively documented effect of in-situ electric fields. Thus, they are used by international guidelines as the basis for limiting human exposure to extremely low-frequency electric and magnetic fields. This study aimed to estimate the phosphene perception locus and threshold during an electric current stimulation for four different frequencies (20, 50, 60, and 100 Hz) and to estimate the associated in-situ electric field. Phosphene perception probabilities were calculated in 20 volunteers using binary logistic regressions applied to perceptual responses resulting from non-invasive transcranial alternating current stimulation between 0 and 2 mA delivered at 20, 50, 60, and 100 Hz. A dosimetry analysis was done to study the in-situ electric field induced in the retina during the electric stimulation. The data indicate that the stimulation current plays a significant role in the model's predictions across all frequencies. Phosphene perception thresholds were lowest at 20 Hz, while no perceptible phosphene were observed at 100 Hz. The findings of this study are crucial for understanding the mechanisms of phosphene induction and further support the retinal origin of phosphenes. The observed thresholds and trends will inform updates to international guidelines and standards.

Open access: https://www.nature.com/articles/s41598-025-06271-8

--

Influence of geomagnetic disturbances on myocardial infarctions in women and men from Brazil

Rezende LFC, De Paula ER, Muella MTAH, Dutra SLG, Rosa RR, Saldiva PHN, Ometto JPHB. Influence of geomagnetic disturbances on myocardial infarctions in women and men from Brazil. Commun Med (Lond). 2025 Jul 1;5(1):247. doi: 10.1038/s43856-025-00887-7.

Abstract

Background Understanding the role of space weather, specifically Geomagnetic Disturbances (GMDs) caused by solar activity, on health outcomes is unclear. One emerging link includes the impact of space weather on myocardial infarctions (MI). In this study we examined the correlation between MI and GMDs in Brazil.

Methods We used a database from the public health in Brazil, focusing on the city of São José dos Campos (23° 10′ 44″ S, 45° 53′ 13″ W), located in the state of São Paulo, during the period of 1998–2005. We focused on admissions for MIs, which included a total of 871 men and 469 women. We categorized

the MI data into three age groups: age 30 and younger, age 31–60, and age over 60. Additionally, we incorporated Planetary Index (Kp) data as an indicator of variations in the Earth's geomagnetic field resulting from solar disturbances, categorized as quiet, moderate, or disturbed days. In our analysis, we employed two methods: statistical counting and the unsupervised clustering known as K-Means, considering the attributes of age, sex, and geomagnetic condition.

Results Here we show that geomagnetic conditions have an impact on MI cases, particularly for women. The rate of relative frequency of MI cases during disturbed geomagnetic conditions is almost three times greater compared to quiet geomagnetic conditions. Using the unsupervised K-Means algorithm, the results indicate that the group associated with disturbed geomagnetic conditions has a higher incidence of MIs in women.

Conclusions Overall, our results provide evidence that women may exhibit a higher susceptibility to the effects of geomagnetic disturbances caused by solar activity on MI.

Open access: https://pmc.ncbi.nlm.nih.gov/articles/PMC12216950/

==

A comprehensive mechanism of biological and health effects of anthropogenic extremely low frequency and wireless communication electromagnetic fields

My note: This well-referenced literature review (373 refs.) explains how electromagnetic fields (EMF) produced in wireless communications--which are polarized, coherent microwaves with highly variable intensity and modulated and pulsed at extremely low frequencies--cause "cancer, infertility, electro-hypersensitivity, and various other pathologies."

Panagopoulos DJ, Yakymenko I, De Iuliis GN, Chrousos GP. A comprehensive mechanism of biological and health effects of anthropogenic extremely low frequency and wireless communication electromagnetic fields. Front. Public Health, Volume 13 – 2025. https://doi.org/10.3389/fpubh.2025.1585441.

Abstract

Exposure to anthropogenic electromagnetic fields (EMFs), especially those of wireless communications (WC) has increased tremendously. This is an unprecedented phenomenon throughout biological evolution because, all anthropogenic EMFs, being fully polarized, coherent, and, especially WC EMFs, highly variable, differ substantially from the natural EMFs. WC EMFs consist of Microwave (MW) carrier waves, modulated, by Extremely Low Frequency (ELF) signals, and included in on/off pulses repeated at various ELF rates. Moreover, they exhibit intense random variability, mainly in the Ultra Low Frequency (ULF) band. Thus, WC EMFs are a combination of MW and ELF/ULF EMFs. The combination of polarization/coherence and intense low-frequency (ELF/ULF) variability seems to be the key to EMF-bioactivity. Epidemiological and laboratory studies highlight a connection between ELF or WC EMF exposure and cancer, infertility, electro-hypersensitivity, and various other pathologies. Studies also find DNA damage and Oxidative Stress (OS) which explain these pathologies. While man-made EMFs cannot directly ionize molecules, they are capable of doing this indirectly in biological tissue, by triggering the biosynthesis of Reactive Oxygen Species (ROS) which can damage biomolecules, including DNA. The

(over)production of ROS and the consequent OS are triggered by irregular gating of Voltage-Gated Ion Channels (VGICs) in the cell membranes as described by the Ion Forced Oscillation (IFO)-VGIC mechanism: Mobile ions within VGICs forced to oscillate by the applied ELF/ULF EMFs exert forces on the voltage sensors of the VGICs, similar to or greater than the forces that physiologically gate those channels, resulting in their irregular gating (dysfunction). Dysfunction of ion channels disrupts intracellular ionic concentrations. This triggers ROS overproduction and OS by the ROS-generating systems/enzymes in the cells, such as the electron transport chain (ETC) in the mitochondria, or the NADPH/NADH oxidases (NOXs), the Nitric Oxide synthases (NOS), etc. The IFO-VGIC mechanism and the consequent OS constitute a comprehensive mechanism that explains all known adverse biological and health effects reported to be induced by anthropogenic EMFs.

Excerpts

"The recorded human and animal carcinogenicity, the DNA/genetic damage, the OS findings, and the reproductive declines due to DNA damage in ovarian or sperm cells or embryonic death, all point toward the same direction: Man-made EMFs induce OS and DNA damage, infertility, cancer, and other related pathologies. The reason why the same effects are observed in a wide variety of animals such as mammals, birds, insects, etc., and humans, is that all biological and health effects initiate in cells and all cells are essentially identical in all animals, humans, and even plants. They have identical membranes, ions, ion channels and pumps, biomolecules such as DNA, RNA, proteins, etc., water, ROS, identical cellular organelles such as nuclei, mitochondria, ribosomes, endoplasmic reticulum, etc., and very similar metabolic processes and regulatory mechanisms. These similarities at the cellular level between all animals and humans are much more fundamental than differences in volume, mass, shape, macroscopic functions, intelligence, etc. As a result, any effect induced by EMFs in animal cells such as OS, DNA damage, etc., is expected to be induced also in the human cells, and vice-versa (7, 66).

The exposure levels in the vast majority of all the aforementioned studies were significantly below the officially accepted exposure limits for ELF and RF EMFs, which are recommended by a private organization called the International Commission on Non-Ionizing Radiation Protection (ICNIRP) to prevent discharges on humans in the case of ELF and acute heating of living tissues in the case of RF/WC EMFs. It is remarkable that this organization arbitrarily ignores the overwhelming evidence of non-thermal effects which constitute the vast majority of effects of anthropogenic EMFs, and yet, governments adopt its recommendations instead of following the Precautionary Principle which dictates the obvious, that no new technology should be applied unless those who promote it have proven its safety beyond any doubt (50, 217–226).

The International Agency for Research on Cancer (IARC) branch of the World Health Organization (WHO), has, since long time, classified both ELF and “RF” (in fact WC) EMFs as possibly carcinogenic to humans (Group 2B) (204, 205, 227). Based on additional scientific evidence after the 2011 IARC classification for “RF” EMFs, several studies have argued that “RF”/WC EMFs should be re-evaluated and classified as probably carcinogenic (Group 2A) or carcinogenic (Group 1) to humans (50, 63, 64, 66, 118–120, 171, 179, 182, 183, 218, 219). Moreover, studies have asked for the urgent application of the Precautionary Principle, stricter exposure limits, especially for WC EMFs, and a moratorium on 5G rollout (50, 215, 217, 218, 225, 228)."

"In conclusion, the IFO-VGIC mechanism that explains VGIC dysfunction, and the subsequent OS, provide a comprehensive biophysical/biochemical mechanism explaining the plethora of experimental and epidemiological findings connecting anthropogenic EMF exposures with OS, DNA/cellular damage and related pathologies such as poor health, EHS, infertility, organic/neurodegenerative diseases, cancer, etc. Even though the mechanistic details of how exactly the ionic perturbations stimulate ROS production by their sources need to be further explored, the basic scheme of the complete EMF-bioeffects mechanism is revealed already. The long existing experimental and epidemiological findings connecting exposure to man-made EMFs and DNA damage, infertility, and cancer, are now explained by the presented comprehensive mechanism. We hope this provides a better understanding of the involved science, a basis for future research, and the establishment of biologically relevant EMF exposure guidelines for effective protection of public health and the environment."

Open access paper:

https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1585441/full

--

Rapid Deployment of 5G Wireless Communication and Risk Assessment on Human Health: Quid Novi?

Selmaoui B, Jamal L, Michelant L. Rapid Deployment of 5G Wireless Communication and Risk Assessment on Human Health: Quid Novi? Bioelectromagnetics. 2025 May;46(4):e70005. doi: 10.1002/bem.70005. PMID: 40276940.

No abstract.

Excerpt

"... the swift and widespread rollout of 5G has not been without controversy. Alongside the excitement over its potential, significant concerns have emerged regarding its potential impact on human health. These concerns stem from the increased exposure to electromagnetic fields (EMFs) associated with 5G technology, particularly as it operates on higher frequency bands, including millimeter waves. Consequently, given the lack of publications concerning the effects of frequencies implemented for 5G (3.5–26 GHz) for the general public, more in-depth studies need to be established due to the increased debates and inconclusive reports about the subject.

Given that 5G is a relatively new technology, short- and long-term studies are still in progress to assess its health implications comprehensively. For this purpose, the European Union Commission via their institutions has launched a call for proposals in the environmental health topic (HORIZON-HL-TH-2021-ENVHLTH-02). This program was implemented to answer to the public concern about the health effect of 5G exposure. The total amount of funding was 30 million euros from Horizon Europe 2021–2027. The results should fill the current knowledge gaps on the effects of wireless technologies on health and the environment. Four projects funded by Horizon Europe have been brought together under the CLUE-H network, involving more than 60 European research organizations across four research consortia: ETAIN, GOLIAT, NextGEM, and SEAWave. Additionally, scientists from outside Europe, including the USA, Korea, and Japan, are also collaborating on these projects...."

Conclusion

"The rapid deployment of 5G brings unprecedented opportunities for technological innovation but also necessitates thorough and ongoing risk assessment regarding its potential health impacts. While current scientific consensus generally supports the safety of 5G under existing guidelines, the evolving nature of the technology, coupled with the long-term uncertainty, underscores the importance of continued research, transparent communication, and adaptive regulatory frameworks. As 5G becomes more ubiquitous, balancing its benefits with precautionary health measures will be crucial to ensuring public trust and safety."

Open access paper: https://onlinelibrary.wiley.com/doi/full/10.1002/bem.70005

--

5G-exposed human skin cells do not respond with altered gene expression and methylation profiles

(My note: Contrary to the title of this paper and media reports, this was not a 5G study because it tested continuous waves in the millimeter band.)

Jyoti J, Gronau I, Cakir E, Hütt M-T, Lerchl A, Meyer V. 5G-exposed human skin cells do not respond with altered gene expression and methylation profiles, *PNAS Nexus*, 4(5), May 2025, doi: 10.1093/pnasnexus/pgaf127.

Abstract

Due to the ever-increasing wirelessly transmitted data, the development of new transmission standards and higher frequencies in the 5G band is required. Despite basic biophysical considerations that argue against health effects, there is public concern about this technology. Because the skin penetration depth at these frequencies is only 1 mm or less, we exposed fibroblasts and keratinocytes to electromagnetic fields up to ten times the permissible limits, for 2 and 48 h in a fully blinded experimental design. Sham-exposed cells served as negative, and UV-exposed cells as positive controls. Differences in gene expression and methylation due to exposure were small and not higher than expected by chance. These data strongly support the assessment that there is no evidence for exposure-induced damage to human skin cells.

This study was funded by the Bundesamt für Strahlenschutz (FKZ 3619S82470).

Open access paper: https://academic.oup.com/pnasnexus/article/4/5/pgaf127/8124427

--

Impact of mobile phone-emitted non-ionizing electromagnetic radiation on parotid gland function: A comprehensive study

Jacob RA, Jose M, Pai VR, Kalal BS. Impact of mobile phone-emitted non-ionizing electromagnetic radiation on parotid gland function: A comprehensive study. *International Journal of Risk & Safety in Medicine*. 2025;0(0). doi:10.1177/09246479251342488.

Abstract

Background and Objective The rapid proliferation of mobile technology has intensified debates on the potential adverse effects of non-ionizing electromagnetic radiation (NIER) from mobile phones on various human organs and cells. This study aimed to evaluate the impact of mobile phone-emitted NIER on parotid gland function.

Methodology This cross-sectional study included 104 student volunteers from the University campus, categorized based on their mobile phone usage and exposure duration. Saliva samples were collected from the dominant (test) and non-dominant (control) sides. Key metrics measured were salivary flow rate, pH, albumin, Ischemia-Modified Albumin (IMA), and the IMA/albumin ratio (IMAR). Data analysis was performed using one-way ANOVA and Fisher's LSD multiple comparisons, with $p \leq 0.05$ considered statistically significant.

Results The salivary flow rate and pH were higher on both sides in all groups, correlating with increased mobile phone usage duration. Although albumin levels were lower on the dominant side, they increased with longer mobile phone use. Salivary IMA and IMAR were higher on the dominant side in subjects using mobile phones for ≤3 years compared to those using them for >3 years.

Interpretation and Conclusion Consistent exposure to NIER and the heat generated by mobile phones adversely affects parotid gland function, as indicated by increased salivary flow rate, pH, and altered levels of albumin, IMA, and IMAR. Public health recommendations should encourage reducing long-duration conversations and using earphones to minimize NIER exposure. Further studies are needed to evaluate the long-term effects of NIER on parotid gland function.

https://journals.sagepub.com/doi/10.1177/09246479251342488

--

Epidemiological criteria for causation applied to human health harms from RF-EMF exposure: Bradford Hill revisited

Frank JW. Epidemiological criteria for causation applied to human health harms from RF-EMF exposure: Bradford Hill revisited. Front. Public Health, 26 May 2025. Volume 13. doi: 10.3389/fpubh.2025.1559868.

Abstract

Purpose: This paper reviews the applicability of standard epidemiological criteria for causation, to the multidisciplinary studies of RF-EMF exposure and various adverse biological and health effects, with the aim of demonstrating that these criteria, although 60 years old, are still helpful in this context—albeit in some cases not entirely straightforward to apply.

Methods: This is a commentary, based on Bradford Hill's criteria for assessing evidence of causation, applied to recent primary studies and systematic reviews of the RF-EMF/health-effects literature. Every

effort has been made to use non-epidemiological language to reach a wide readership of biologists, physicists, and engineers now active in this field.

Results: A rapidly growing number of human observational epidemiological studies have assessed the association of diverse adverse health effects with RF-EMF exposures. However, existing systematic reviews and meta-analyses of these primary studies have substantially diverged in their conclusions. The application of Bradford Hill's epidemiological criteria for assessing evidence of causation, originally designed for use in occupational and environmental health, casts light on some of reasons for this divergence, mostly reflecting the key weaknesses in the primary literature, which are discussed in detail. As a result of these threats to their validity—particularly the facts that (1) exposure measurement is typically subject to substantial error, and (2) insufficient time has elapsed, since modern cell phone use began in earnest, to allow tumors of longer latency to develop—most primary studies to date, and therefore many published systematic reviews of them, probably underestimate the true potential for causation, if in fact this association is causal.

Conclusion and recommendations: In view of these findings, international experts representing professional and scientific organizations in this field should convene an independent Guidelines development process to inform future epidemiological studies of associations between RF-EMF exposures and human health outcomes. Wide dissemination of such Guidelines could help researchers, journals and their reviewers in this field to execute, review and publish higher-quality studies to better inform evidence-based policy.

Open access paper:

https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1559868

--

Skin Fibroblasts from Individuals Self-Diagnosed as Electrosensitive Reveal Two Distinct Subsets with Delayed Nucleoshuttling of the ATM Protein in Common

Sonzogni L, Al-Choboq J, Combemale P, Massardier-Pilonchéry A, Bouchet A, May P, Doré J-F, Debouzy J-C, Bourguignon M, Dréan YL, et al. Skin Fibroblasts from Individuals Self-Diagnosed as Electrosensitive Reveal Two Distinct Subsets with Delayed Nucleoshuttling of the ATM Protein in Common. *International Journal of Molecular Sciences*. 2025; 26(10):4792. doi: 10.3390/ijms26104792.

Abstract

Electromagnetic hyper-sensitivity (EHS) and its causal link with radio-frequencies raise a major question of public health. In the frame of the clinical study DEMETER, 26 adult volunteers self-diagnosed as EHS-positive agreed to reply to a self-assessment questionnaire and to provide a skin biopsy sampling to establish a primary fibroblast cell line. The questionnaire and the biological data revealed, independently, 2 subsets of donors associated each with a low background, highly responsive (LBHR) and a high background, lowly responsive (HBLR) phenotype. A couple of subsets based on questionnaire data and based on the yield of spontaneous DNA double-strand breaks were found to be composed of the same donors at 64% identity. After exposure to X-rays, and application of anti-γH2AX, pATM, and MRE11 immunofluorescence, all the DEMETER fibroblasts (26/26) elicited a delayed radiation-induced

ATM nucleoshuttling (RIANS). The use of RIANS biomarkers showed that the 2 phenotypes described above corresponded to DEMETER donors with a high risk of cancer (LBHR) or high risk of accelerated aging (HBLR). By exposing DEMETER cells to H_2O_2 followed by an antioxidative agent, we confirmed that EHS may be related to the management of DNA strand breaks. A preliminary molecular model of EHS inspired by the RIANS model was proposed.

Open access paper: https://www.mdpi.com/1422-0067/26/10/4792

--

Exposure Perception and Symptom Reporting in Idiopathic Environmental Intolerance Attributed to Electromagnetic Fields Using a Co-Designed Provocation Test

Ledent M, Bordarie J, Vatovez B, Dieudonné M, Prignot N, Vanderstraeten J, Bouland C, De Clercq EM. Exposure Perception and Symptom Reporting in Idiopathic Environmental Intolerance Attributed to Electromagnetic Fields Using a Co-Designed Provocation Test. Bioelectromagnetics. 2025 Apr;46(3):e70006. doi: 10.1002/bem.70006.

Abstract

Idiopathic Environmental Intolerance Attributed to Electromagnetic Fields (IEI-EMF) is a syndrome that defines people who report symptoms that they attribute to their exposure to EMF sources, without any identified underlying medical condition to explain these symptoms. To date, provocation protocols have failed to demonstrate a consistent relationship between EMF exposure and reported symptoms, raising questions among some researchers and individuals with IEI-EMF about the relevance of these protocols for studying the syndrome. To address these criticisms, a provocation protocol was co-designed in collaboration with individuals with IEI-EMF. This study presents the results of the tests, with a focus on exposure perception and symptom reporting among IEI-EMF volunteers. A total of 47 IEI-EMF volunteers were enrolled and participated in an open-field habituation session. Of these, 27 completed the first double-blind controlled exposure session, while 26 and 16 volunteers, respectively, participated in three sessions for collective analyses and 12 sessions for individual-level analyses. At the individual level, no consistent association was found between exposure perception certainty level and exposure status, except for one volunteer whose perception was mostly consistent with exposure status. Similarly, symptom reporting did not align with exposure status, except for the same volunteer, whose symptom reporting showed a borderline significant result with exposure status. However, for half of the volunteers, symptom reporting was significantly correlated with exposure perception certainty level, supporting a nocebo hypothesis. At the collective level, no consistency was observed between exposure perception certainty level, symptom reporting, and exposure status. This study discusses the conditions necessary for future provocation protocols to enhance their relevance, acceptability, and potential utility in a possible care-oriented approach. It also considers criticisms of using exposure perception and symptom reporting as outcomes in provocation protocols, despite their central role in how individuals identify themselves as individuals with IEI-EMF.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.70006

--

Electrohypersensitivity: what is belief and what is known?

de Vocht F, Martin Röösli M. Electrohypersensitivity: what is belief and what is known? Frontiers in Public Health. Vol. 13. 2025. doi:10.3389/fpubh.2025.1603692.

No abstract

https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1603692/full

--

What is the effect of alarmist media and radiofrequency electromagnetic field (RF-EMF) exposure on salivary cortisol and non-specific symptoms?

Verrender A, Wallace NK, Loughran SP, Wallace C, Beange J, Croft RJ (2025). What is the effect of alarmist media and radiofrequency electromagnetic field (RF-EMF) exposure on salivary cortisol and non-specific symptoms? *Applied Psychology: Health and Well-Being*, 17(3), e70044. https://doi.org/10.1111/aphw.70044.

Abstract

While there is consistent evidence that the symptoms reported by people who experience Idiopathic Environmental Intolerance attributed to Electromagnetic Fields (IEI-EMF) are closely associated with a nocebo effect, and that alarmist media reports may contribute to this nocebo effect, some methodological criticisms remain to be resolved. This study aimed to replicate previous findings and determine whether viewing an alarmist media report and being openly exposed to radiofrequency electromagnetic fields (RF-EMF) could induce a salivary cortisol response. A total of 144 participants were randomly assigned to watch either an alarmist or control video before completing an open-label provocation trial where they were either exposed or not exposed to RF-EMF. Personality factors, RF-EMF risk perception (pre- and post-video), symptoms and salivary cortisol were assessed. Consistent with previous studies, participants who were aware that they were being exposed had increased symptoms compared to participants who were aware they were not being exposed. However, the current study failed to replicate an effect of viewing an alarmist media report and being openly exposed to RF-EMF on symptoms and failed to identify an effect on salivary cortisol. This suggests that awareness and belief of exposure play a more important role in symptom perception than underlying physiological processes.

CONFLICT OF INTEREST STATEMENT

Sarah P. Loughran receives funding from the National Health and Medical Research Council of Australia (NHMRC). She is the Director of Radiation Research and Advice at the Australian Radiation Protection and Nuclear Safety Agency (ARPANSA), a member of the Scientific Expert Group at the International Commission on Non-Ionizing Radiation Protection (ICNIRP), and a member of the World Health Organisation Task Group on Radiofrequency Fields and Health Risks. Rodney J. Croft receives funding from the National Health and Medical Research Council of Australia (NHMRC) and is a member of the Scientific Expert Group at the International Commission on Non-Ionizing Radiation Protection (ICNIRP). No potential competing interests were reported by the remaining authors.

Open access paper: https://iaap-journals.onlinelibrary.wiley.com/doi/full/10.1111/aphw.70044

--

Exploring the impact of environmental factors on male reproductive health through epigenetics

Zhang Y, Song JY, Sun ZG. Exploring the impact of environmental factors on male reproductive health through epigenetics. Reprod Toxicol. 2025 Mar;132:108832. doi: 10.1016/j.reprotox.2025.108832. Epub 2025 Jan 6. PMID: 39778664.

Abstract

Male infertility has become an increasingly severe global health issue, with its incidence significantly rising over the past few decades. This paper delves into the crucial role of epigenetics in male reproductive health, focusing particularly on the effects of DNA methylation, histone modifications, chromatin remodeling and non-coding RNAs regulation on spermatogenesis. Exposure to various environmental factors can cause sperm DNA damage, leading to epigenetic abnormalities. Among these factors, we have discussed heavy metals (including Zinc, Cadmium, Arsenic, Copper), phthalates, electromagnetic radiation, and temperature in detail. Notably, aberrations in DNA methylation are closely associated with various symptoms of male infertility, and histone modifications and chromatin remodeling are essential for sperm maturation and function. By synthesizing existing literature and experimental data, this narrative review investigates how environmental factors influence male reproductive health through epigenetic mechanisms, thus providing new theoretical foundations and practical guidelines for the early diagnosis and treatment of male infertility.

Excerpt

3.3 Electromagnetic radiation and sperm epigenetics

The operating frequencies for analog mobile phones are 450–900 MHz, digital phones (Global System for Mobile Communications) range from 850 to 1900 MHz, and third-generation phones operate around 2000 MHz[144]. Electromagnetic waves (EMW) can affect sperm epigenetics through two mechanisms: 1. EMW-specific effects 2. Thermal molecular effects [145].

Experiments have found that when male rats were exposed to Wi-Fi from laptops for seven hours daily, there was a reduction in sperm count and motility within a week [146,147]. Additionally, the thermal effects of electromagnetic radiation increase free radicals around the testes, which oxidize the phospholipids in the sperm outer membrane, thus reducing membrane fluidity and impairing motility [148].

Furthermore, oxidative stress mediated by electromagnetic radiation leads to the phosphorylation of heat shock proteins, causing chromatin damage [148], thereby altering sperm epigenetics. Mice lacking Hsp90 or with highly phosphorylated Hsp proteins show cessation of spermatogonia during meiosis, leading to loss of germ cells and reduced testicular size [149]. Research indicates that Hsp90 possesses ATPase activity, stimulated by nuclear autoantigenic sperm protein (NASP) in mouse sperm nuclei [150]. A deficiency in Hsp90 or high phosphorylation of Hsp proteins can halt spermatogonia during meiosis, leading to germ cell loss and reduced testicular size [149].

In vitro animal studies by Liu et al. [151] concluded that radiofrequency radiation could have detrimental effects on human semen quality. Research has found that carrying a mobile phone or Bluetooth headset in a trouser pocket or on the waist can decrease the proportion of sperm with rapid progressive motility and increase the rate of abnormal sperm morphology [152]. Therefore, men should avoid placing mobile phones near the trousers pocket or close to the testicles to prevent potential harmful effects of radiofrequency radiation on the male reproductive system [153].

https://pubmed.ncbi.nlm.nih.gov/39778664/

--

Laptop and tablet use and their influence on total motile sperm count parameters: are laptops linked to infertility in Jamaican men?

Sterling L, Carroll K, Harris LR. Laptop and tablet use and their influence on total motile sperm count parameters: are laptops linked to infertility in Jamaican men? Rev Int Androl. 2024 Dec;22(4):25-32. doi: 10.22514/j.androl.2024.027.

Abstract

Background: This study aims to determine the impact of laptop and tablet use on total motile sperm count (TMSC) in men being investigated for assisted reproduction.

Methods: A cross-sectional study was conducted on 156 men attending a fertility clinic in Jamaica. Routine semen analyses were performed and parameters specific to TMSC assessed. All data analyses were performed using SPSS Version 26. Logistic regression analyses were performed to independently predict the impact of quantifiable measures of laptop and tablet use. The main outcome measures were the parameters associated with TMSC.

Results: Overall, 64% of the participants reported using laptops and 36% reported using tablets. There was a significant relationship seen with time spent on laptops and time trying to conceive ($p = 0.015$). Regression analyses showed that persons who used their laptops for 2 to 5 h daily were approximately 16 times (adjusted Odds Ratio (aOR) = 15.9; 95% Confidence Interval (CI), 2.5-103.3, $p = 0.004$) more likely to be diagnosed with low semen volume (hypospermia). Although no significant association was found between total motile sperm count (TMSC) and laptop use, a trend towards significance was observed with high laptop use ($p = 0.052$), suggesting potential implications for TMSC as a predictor of pregnancy outcomes.

Conclusions: Our findings highlight the need for clinicians to take into consideration wireless device usage in men undergoing fertility investigations.

Open access paper: https://files.intandro.com/files/article/20241230-28/pdf/RIA20240615001.pdf

--

Electromagnetic fields from mobile phones: a risk for maintaining energy homeostasis?

Seewooruttun C, Mai TC, Corona A, Delanaud S, Seze R, Bach V, Desailloud R, Pelletier A. Electromagnetic fields from mobile phones: a risk for maintaining energy homeostasis? Ann Endocrinol (Paris). 2025 May 6:101782. doi: 10.1016/j.ando.2025.101782.

Abstract

In the world, there is a near ubiquitous presence of a low-intensity radiofrequency electromagnetic field (RF-EMF) radiation, due to telecommunications as mobile phones. However, their rapid expansion raises concerns about possible interaction with biological mechanisms. The RF-EMF safety guidelines recommended limits to protect against the thermal heating, the most recognized effect at high intensity levels with a known biophysical mechanism. Among all the effects studied, the impact of RF-EMF exposure on thermoregulation is one of the most important aspects of this research. This review aims to present the complex relationship between RF-EMF exposure and thermoregulation, at intensity levels below the threshold to produce thermal effects. In fact, most studies showed that RF-EMF exposure at 900 MHz seems to be elicited physiological and biological effects similar to responses inducing by cold environment in two different rodent models. In this brief review, we will describe these responses observed during or after RF-EMF exposure to give physiological and molecular mechanisms of the interaction between the organism and RF-EMF exposure at low levels and the potential implications for environmental health and safety.

Conclusion

In modern society, exposure to RF-EMF is very common and nearly impossible to avoid. Despite its major impact in driving technological advancements, it is important to address the growing public concern in regards to RF-EMF exposure. The thermal effects of high-intensity RF-EMF exposure (SAR>4 W/kg) are well documented in the literature. However, there is now emerging evidence that low-intensity RF-EMF exposure (SAR<4 W/kg) can also elicit thermoregulatory responses associated with a cold sensation. The present review highlights the behavioural thermoregulation associated with a cold sensation in rodents exposed to 900 MHz. We also provide molecular insights on RF exposure effects on different thermogenic mechanisms. The first results suggest that RF exposure primarily influences WAT browning rather than BAT thermogenesis, as reported by the increased UCP1 staining and reduced adipocyte size in WAT depots. In contrast, the cold-induced transcriptional changes on BAT thermogenesis were not observed after exposure to RF, although we detected an increased plasma levels of noradrenaline, NEFA in exposed rats. In the short term, these adaptive responses do not seem to compromise homeostasis and therefore the health of organisms, but what about the long-term consequences?

With the introduction of 3.5 GHz for 5G and early use of wireless technology by infants, further research is needed to determine these thermoregulatory effects. Investigating the peripheral tail temperature and thermal preference of 5G-exposed rats at different age could provide valuable insights into their thermoregulatory responses. Additionally, the implication of the thyroid on these responses need to be more investigate since this organ could allow a long-term adaptative response.

https://pubmed.ncbi.nlm.nih.gov/40339686/

--

Potential Impacts of Radiofrequency Electromagnetic Fields on the Central Nervous System, Brain Neurotransmitter Dynamics and Reproductive System

Eskandarpour M, Nooramin A. Potential Impacts of Radiofrequency Electromagnetic Fields on the Central Nervous System, Brain Neurotransmitter Dynamics and Reproductive System. *AUT Journal of Electrical Engineering*, 2025. doi: 10.22060/eej.2025.23753.5631.

Abstract

Human life has been increasingly affected by the rapid advancement of electronic technology and the widespread use of devices emitting electromagnetic radiation (EMR), such as Wi-Fi and mobile phones. While much remains unclear, studies suggest that electromagnetic fields (EMFs) can influence human health, particularly reproduction and the nervous system. EMF exposure, including from non-ionizing radiation produced by Wi-Fi and mobile phones, has been linked to potential effects on the male and female reproductive systems, embryonic development, and neuronal health. Key mechanisms include oxidative stress, thermal effects, changes in neurotransmitter metabolism, receptor function, nerve cell apoptosis, and ion channel dynamics. However, the long-term health risks, especially in children and adolescents due to prolonged exposure, remain a topic of debate. Despite current studies not confirming that RF-EMW from Wi-Fi exceeds safety guidelines, further research is essential to fully understand the implications of RF-EMW exposure on human health, particularly regarding reproduction and neurological effects. This review highlights the need for updated safety standards, more refined regulatory frameworks, and long-term investigations to clarify the potential biological and neurobiological consequences of EMF exposure.

Conclusion

RF-EMR is a form of nonionizing radiation that does not have the energy to destroy biological structures directly. However, it impressively and importantly acts on the human nervous system. Although RF-EMR itself does not act by ionizing molecules, the exposure to it can further bring about thermal effects at higher frequencies whereby raised temperature of the tissue may have an impact on cellular processes. Other than thermal effects, RF-EMR exposure can cause non-thermal biological responses through mechanisms such as oxidative stress, altered neurotransmitter activity, and changes in the flux of calcium ions within neurons. These cellular disruptions can lead to neuronal apoptosis, which hampers appropriate brain function and can potentially affect cognition and behavior. While the effects on human health are still debated, it is evident that the nervous system, given its sensitivity to electromagnetic disturbances, may be vulnerable to prolonged exposure. While radiofrequency electromagnetic radiation from Wi-Fi, for example, does involve much smaller power compared with mobile phones, and exposure is much farther from the body, there remains considerable ignorance regarding its long-term outcomes, particularly to sensitive populations of children and teenagers.

While various studies have tended to address a number of questions about RF-EMR risk, especially for brain function and reproductive health, current evidence remains conflicting and thus has necessitated further detailed investigation. It is not possible to decide on a definite conclusion because of the complexity of the biological systems implicated and the difficulty in extrapolating from animal models to human health. However, while neutral or even positive effects can be envisaged at specific levels of RF-

EMR exposure, particularly in medical uses, there is still concern about possible risks that may occur above threshold values, particularly in the nervous and reproductive systems. These are potentially very disruptive to neurotransmitter functions, neuronal health, and even sperm and egg cell integrity. Larger, more comprehensive, long-term studies are urgently needed to ascertain the level of exposure that is not harmful and for a full understanding of all biological effects. In summary, exposure to RF-EMR continues to increase with each new generation of technology, including 5G. It is, therefore, of crucial importance that accurate safety guidelines and regulatory frameworks are updated. Continued research into the possible health risks, in particular the cumulative effects of long-term exposure, is essential to safeguarding public health, especially for the younger generation who are exposed more frequently and for longer periods.

Open access paper: https://eej.aut.ac.ir/article_5734.html

--

Assessing RF EMF exposure in multiple microenvironments across ten European countries with a focus on 5G

Veludo AF, Stroobandt B, Van Bladel H, Sandoval-Diez N, Deprez K, Aerts S, ..., de Vocht F, Joseph W, Röösli M. Assessing radiofrequency electromagnetic field exposure in multiple microenvironments across ten European countries with a focus on 5G. Environment International, 2025. https://doi.org/10.1016/j.envint.2025.109540.

Abstract

To evaluate the implementation of 5G in Europe, we have systematically measured environmental, auto-induced downlink (DL) and uplink (UL) radiofrequency electromagnetic field (RF-EMF) exposure in more than 800 microenvironments in ten European countries. Outdoor, indoor, and public transport microenvironments were measured in two cities and three villages in each country. Exposure was measured during three mobile-phone user scenarios: flight mode (non-user), inducing maximum DL traffic (max DL) or maximum UL traffic (max UL). The mobile phone was carried in a backpack, placed 30 cm from an ExpoM-RF 4 that continuously measured 35 frequency bands (87.5 MHz–6 GHz). For each user scenario, mean exposure levels were calculated. In the non-user scenario, mean exposure levels ranged from 0.33 to 1.72 mW/m2 per country and were lower in Switzerland, Belgium and Italy. RF-EMF levels were, on average, 80 % lower in villages compared to cities, with DL bands contributing the most in this scenario. During max DL, exposure increased mainly due to the 5G band at 3.5 GHz (mean exposure per country 2.61–11.12 mW/m2). However, the time-division nature of this band prevents distinguishing between DL and UL signals with the ExpoM-RF4. Exposure levels were the highest during max UL, particularly in the Netherlands, Italy and Belgium, with 50 % of the mean levels per country above 16 mW/m2. Exposure was, on average, 35 % higher in villages compared to cities. Environmental exposure levels were below international guideline values. Countries with precautionary limits had lower environmental exposure levels but higher auto-induced uplink exposure during data upload.

Conclusions

In this comprehensive European RF-EMF measurement study, environmental exposure levels during a non-user scenario are in line with previous research in Europe and well below international guidelines, suggesting that environmental exposure has not substantially changed since the introduction of 5G.

Moreover, we could demonstrate that environmental RF-EMF exposure is mainly attributed to the Mobile DL frequency bands emitted from base stations and increases with increased urbanisation, a result of the dense network in urban settings. Additionally, we observed that environmental exposure levels are generally lower in countries with precautionary limits, namely Switzerland, Italy, and Belgium. Inducing DL traffic resulted in higher personal exposure levels, attributed to the Mobile TDD 3.5 GHz in countries where 5G is widely available. However, more studies are needed to understand the real contribution of beamforming when inducing traffic. RF-EMF exposure from an own phone during data upload was attributed mainly to 4G Mobile UL bands, and to a lower extent to the 5G band at 3.5 GHz. Additionally, in this scenario exposure levels decreased with increasing urbanisation.

Future research should monitor the evolution of environmental and auto-induced RF-EMF exposure in the different countries alongside the ongoing deployment of 5G networks in Europe.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412025002910

--

5G EMF Exposure at 3.6 GHz in Greece Using Data From Frequency-Selective Monitoring Sensors

Iakovidis S, Manassas A, Apostolidis C, Samaras T. 5G EMF Exposure at 3.6 GHz in Greece Using Data From Frequency-Selective Monitoring Sensors. Bioelectromagnetics. 2025 May;46(4):e70008. doi: 10.1002/bem.70008.

Abstract

The introduction of 5G networks has raised public concerns about potential changes in environmental electromagnetic field (EMF) exposure. This study analyzes continuous monitoring data collected over 2 years (August 2022-October 2024) from 13 frequency-selective monitoring sensors located in Greece's five largest cities. Focusing on the 3.6 GHz band, we evaluated trends and weekly variations in EMF levels. Results indicated a gradual increase in EMF exposure at 3.6 GHz, driven by the growing penetration of 5G infrastructure and devices. Notably, this band exhibited higher maximum-to-median power density ratios compared to other frequency bands, attributable to active antenna systems' characteristics and traffic variations. Applying the ICNIRP 2020 guidelines, we found that 30-min averaged values significantly reduced these variations. All measured EMF levels, including maximum values, remained well below Greek and international safety limits. These findings, especially the increasing trend identified for the EMF levels, underscore the importance of continuous monitoring networks for assessing EMF exposure to existing and emerging telecommunications networks and ensuring compliance with safety standards.

Summary

- Over the last 2 years, 5G EMF levels at the 3.6 GHz frequency band increased due to network expansion but stayed well below ICNIRP and Greek reference levels.
- The 3.6 GHz band showed much higher maximum-to-median E-field variability than other frequency bands, driven by active antenna systems and traffic variations.

- Using 30-min averaging, as recommended by ICNIRP (2020), significantly reduced exposure variability at 3.6 GHz, aligning it with the one observed in frequency bands used by cellular networks of previous generations.

Excerpts

"A more representative view of all 13 monitoring sensors used in this study is provided in Figure 4. The figure shows clearly that in the 3.6 GHz frequency band the ratio of the maximum 6-min-averaged power density in a week can become 12 times higher than the median 6-min-averaged power density of that week."

"The maximum-to-median ratio for the 30-min averaged values at the 3.6 GHz frequency band, became of the same order of magnitude (≈3) as those at frequency bands of older generation cellular networks (≈2)."

"The highest 6-min-averaged power density registered over 2 years was 0.053 W/m^2, which is much lower than both the ICNIRP and the Greek reference levels for general public exposure, that is, 10 and 6 W/m^2, respectively. It's worth mentioning that any E-field value used in this study corresponds to the 6-min RMS, that is, it is immediately comparable to established safety limits in Greece. However, the significantly higher E-field variability at the 3.6 GHz frequency band, compared to other cellular network bands, highlights the need for application of “extrapolation to the maximum” techniques when 6-min spot (in situ) measurements are performed for compliance assessment. This point should also be considered when exposure assessment studies are performed."

"Conclusions

In this study, continuous measurements of E-field from 13 monitoring sensors located in the five largest Greek cities and collected over the last 2 years were analyzed with a focus on the 3.6 GHz frequency band, where 5G networks operate. An increasing trend in environmental EMF levels at this band over these years was identified but the maximum registered incident power density remained well below the reference levels of ICNIRP and the Greek legislation. A considerably higher variation of exposure levels within a week at the 3.6 GHz band compared to other frequency bands was observed in terms of weekly maximum-to-median incident power ratio. This result highlights the necessity of extrapolation to the actual maximum when short-time measurements (e.g., of 6 min) are used as a proxy for compliance assessment. However, this high maximum-to-median ratio considerably decreases by considering 30-min averaging of the measurements, as suggested by the 2020 ICNIRP guidelines. Notably, the median E-field values at the 3.6 GHz band are lower than those of legacy networks.

No significant trends at older frequency bands were identified possibly due to limited spatial sampling, with only 13 measurement locations. Therefore, further research with broader geographical coverage and additional monitoring sensors would be valuable to evaluate these trends of exposure in the context of 5G rollout."

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.70008

--

Deduction of Extrapolation Factors in Realistic Scenarios for In-Situ Assessment of 5G Base Stations

Villaescusa-Tebar A, Peris JR, Garcia-Pardo C. Deduction of Extrapolation Factors in Realistic Scenarios for In-Situ Assessment of 5G Base Stations. 2025 19th European Conference on Antennas and Propagation (EuCAP), Stockholm, Sweden, 2025, pp. 1-5, doi: 10.23919/EuCAP63536.2025.10999421.

Abstract

Accurate assessment of electromagnetic field (EMF) exposure in 5G networks presents challenges due to beamforming and advanced technologies. Traditional frequency-selective methods provide actual exposure values but cannot distinguish beams or cells. Code-selective techniques differentiate cells and beams, but require complex extrapolation processes based on unavailable network parameters like beamforming patterns. This study investigates extrapolation factors like Fextbeam in realistic indoor and outdoor 5G environments, using frequency-selective and code-selective measurements. Measurements under different traffic conditions evaluated the effect of beam patterns and power. Results showed higher variability in extrapolated values due to differences in antenna radiation patterns. Outdoor environments exhibited more stable values, indicating the critical role of antenna patterns, especially indoors.

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=10999421&isnumber=10999162

--

An approach for annual analysis of EMF exposure in highly sensitive areas of kindergartens and schools

Kljajic D, Djuric N, Pasquino N, Solmonte N, Djuric S. An approach for annual analysis of EMF exposure in highly sensitive areas of kindergartens and schools. Radiat Prot Dosimetry. 2025 May 19:ncaf047. doi: 10.1093/rpd/ncaf047.

Abstract

Continuous increase of electromagnetic field (EMF) emission in the environment causes worldwide research on various short- and long-term EMF measurements and exposure assessments. Measurements are usually performed in sensitive areas, with conventional measuring approaches and worst-case exposure assessments, which are unable to provide any deeper EMF behavior analyses. Therefore, this paper brings an innovative EMF exposure analysis approach, based on time-averaging of time series data, disclosing some concealed information on long-term exposure. The averaging was done on the set of EMF time series acquired by three sensors in the EMF Regulatory Agency for Electronic Communications and Postal Services (RATEL) monitoring network. Sensors are installed at extremely sensitive locations-on buildings of two kindergartens and an elementary school. Analyses revealed patterns of daily/weekly exposure changes and their differences between working days and weekends, as well as annual decrease/increase, at each location. Comparative analysis among locations showed differences in exposure behavior regarding its peaks, slopes of the decrease/increase, and time intervals of the day with higher exposure.

https://pubmed.ncbi.nlm.nih.gov/40384509/

--

Assessment of RF EMF Exposure to Car Driver from Monopole Array Antennas in V2V Communications Considering Thermal Characteristics

My note: This paper only addresses estimated RF-EMF exposure relative to inadequate ICNIRP guidelines which only protect against heating risk.

Wang S, Lu M. Assessment of RF Electromagnetic Exposure to Car Driver from Monopole Array Antennas in V2V Communications Considering Thermal Characteristics. *Sensors*. 2025; 25(10):3247. https://doi.org/10.3390/s25103247.

Abstract

Vehicles are rapidly evolving into objects of intelligent interconnection. Vehicle-to-Vehicle (V2V) communications enable the interconnection between vehicles, while also leading to new electromagnetic exposure scenarios. This paper integrates a monopole array antenna into a shark-fin antenna on the car roof for V2V communications and evaluates the specific absorption rate (SAR) and temperature rise of a human body in a smart mobility communication scenario operating at 5.9 GHz. The V2V antenna is modeled and placed on a 3D vehicle model using COMSOL Multiphysics (v.6.2) to numerically estimate the SAR in the head and body regions of the human body model (adult male) inside the vehicle. Both the localized and whole-body 30 min average SAR are lower than the International Commission on Non-Ionizing Radiation Protection (ICNIRP) occupational restrictions for electromagnetic field exposure from 100 kHz to 6 GHz, being equal in the worst-case scenario to 0.981 W/kg (for the head), which is 9.81% of the ICNIRP limit (10 W/kg), and 0.008728 W/kg (for the whole-body average), which is 2.18% of the ICNIRP limit (0.4 W/kg). The 30 min average human core temperature rise is 0.055 °C, which is 5.5% of the ICNIRP limit. This indicates that, in typical automotive scenarios, the electromagnetic exposure from a monopole array antenna for V2V communications does not pose threat to the human body. This study provides knowledge related to emerging exposure scenarios in intelligent mobility communication, which is beneficial for evaluating possible health impacts and designing public health management policies.

Open access paper: https://www.mdpi.com/1424-8220/25/10/3247

--

Exposure to radiofrequency electromagnetic fields and IARC carcinogen assessment: Risk of Bias preliminary literature assessment for 10 key characteristics of human carcinogens

My note: Apparently, ICNIRP supporters plan to dismiss the IARC Group 2B classification of RF-EMF carcinogenicity and the recent WHO systematic review on animal laboratory carcinogenicity studies (Mevissen et al., 2025) by claiming that all of the *in vitro* and *in vivo* studies that found evidence for carcinogenicity are poor quality.

Simkó M, Repacholi MH, Foster KR, Mattsson M-O, Croft RJ, Scarfi MR, Vijayalaxmi. Exposure to radiofrequency electromagnetic fields and IARC carcinogen assessment: Risk of Bias preliminary literature assessment for 10 key characteristics of human carcinogens. Mutation Research - Reviews in Mutation Research, Volume 796, 2025, https://doi.org/10.1016/j.mrrev.2025.108545

Highlights

We analyzed if the ten key characteristics (KCs) of human carcinogens according to IARC are influenced by RF-EMF exposure.

We reviewed 159 articles by extracting relevant exposure and experimental data.

A risk of bias (RoB) analysis was conducted using 6 criteria.

There is a strong negative association between study quality and the likelihood of reporting effects of RF-EMF exposures.

The heterogeneity and overall poor study quality suggest the need for high-quality studies.

Abstract

This is the first assessment of evidence needed to determine whether exposure to radiofrequency electromagnetic fields (RF-EMF) exposures, below the levels recommended in the ICNIRP (2020) guidelines, can influence any of the ten key characteristics (KCs) of human carcinogens developed by the International Agency for Research on Cancer (IARC). We define the 10 KCs and their relevance to carcinogenesis; review in vivo and in vitro studies relevant to the KCs; and conduct a risk of bias (RoB) analysis using 6 criteria. We did not include KC studies on genotoxicity or oxidative stress since Romeo et al. (2024) and Meyer et al. (2024) recently published relevant systematic reviews, but note their respective conclusions. From the other 8 KCs we identified 119 in vitro and 40 in vitro measurements of in vivo studies through 30 June 2023, with 38 % reporting statistically significant effects of exposure. We identified a strong association between the quality of study and outcome, with those meeting more RoB criteria less likely to report statistically significant effects. Effects were reported over the entire frequency range, exposure levels, and biological endpoints with no apparent pattern of exposure parameters resulting in effects. Only KC10 (alters cell proliferation, cell death or nutrient supply) has sufficient studies to analyse, but the other KCs had few studies and diverse endpoints. A few relatively high-quality positive studies require follow-up through additional targeted studies. The heterogeneity and overall poor study quality suggest the need for high-quality studies on these endpoints, preferably adhering to standards such as the Organization for Economic Co-operation and Development [28].

6. Conclusions

This review aimed to assess the extent and adequacy of experimental data bearing on whether exposure to RF-EMF could influence any of the 10 IARC key characteristics of human carcinogenesis. Since two PRISMA-compliant SRs on the KCs genotoxicity and oxidative stress were recently published [35], [23], this present review was limited to the 8 remaining KCs for which no systematic reviews have been conducted so far. It became apparent that the present database of in vitro and in vivo studies was so diverse and scattered in their quality that helpful outcomes of SRs on most of the KCs would be unfeasible, and any conclusions from such reviews would have a very low confidence. The two PRISMA-compliant SRs as well as previous comprehensive reviews on in vitro and in vivo genotoxicity studies (e.g. [46], and our review of studies on the 8 KCs, have all reached the same conclusion: there is clearly a need for much higher quality RF-EMF bioeffects studies on these KCs.

There are, however, a few statistically significant results in the highest quality studies (Tier 1 studies in

Romeo et al. [35] and studies meeting 5 or 6 RoB criteria (present study). These deserve close examination, and replication if warranted by stronger studies, preferably done under GLP [28], or are otherwise compliant with OHAT recommendations.

As a final comment, it is not useful for systematic reviews to examine hundreds of papers on a given topic, only to find that the overall quality of the large majority of the studies is too low to permit conclusions with a high level of confidence.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

Authors MS, MOM, and MRS acknowledge funding from the NextGEM project from the Horizon Europe Research and Innovation programme under Grant Agreement No 101057527 for part of this study. Views and opinions expressed are those of the author only and do not necessarily reflect those of the European Union or the European Health and Digital Executive Agency. Neither the European Union nor the granting authority can be held responsible for them. The other authors received no external funding for this research.

Open access paper: https://www.sciencedirect.com/science/article/pii/S138357422500016X

--

Electromagnetic fields regulate iron metabolism: From mechanisms to applications

Wang S, Yang J, Zhen C, Wang H, Shang P. Electromagnetic fields regulate iron metabolism: From mechanisms to applications. Journal of Advanced Research. 2025, https://doi.org/10.1016/j.jare.2025.04.044.

Highlights

• Electromagnetic fields (EMFs) can influence the progression of various diseases by regulating iron metabolism.

• The regulation of iron metabolism by EMFs is highly parameter-specific and varies across different tissues and cells.
• The electromagnetic properties of iron-containing molecules/proteins and iron-rich tissues influence iron metabolism modulation by EMFs.
• EMFs can modulate iron metabolism through their effects on membrane structure and ion channels.
• EMFs can modulate iron metabolism by regulating the generation and activity of Reactive Oxygen Species (ROS).

Abstract

Background Electromagnetic fields (EMFs), as a form of physical therapy, have been widely applied in biomedicine. Iron, the most abundant trace metal in living organisms, plays a critical role in various

physiological processes, and imbalances in its metabolism are closely associated with the development and progression of numerous diseases. Numerous studies have demonstrated that EMF exposure induces significant changes in both systemic and cellular iron metabolism.

Aim of Review This review aims to summarize the evidence and potential biophysical mechanisms underlying the role of EMFs in regulating iron metabolism, thereby enhancing the understanding of their biological mechanisms and expanding their potential applications in biomedical fields.

Key Scientific Concepts of Review In this review, we have synthesized research findings and proposed the hypothesis that the biophysical mechanisms of EMFs regulate iron metabolism involve the special electromagnetic properties of iron-containing proteins and iron-enriched tissues, as well as the modulation of membrane structure and function, ion channels, and the generation and activity of Reactive Oxygen Species (ROS). Then, the review summarizes the latest advances in the effects of EMFs on iron metabolism and their safety, as well as their impact on immunoregulation, cardiovascular diseases, neurological diseases, orthopedic diseases, diabetes, liver injury, and cancer.

Conclusion and perspective

EMFs have been widely applied as a non-invasive physical alternative or adjunctive therapy in translational medicine. The regulation of iron metabolism by EMFs is a crucial mechanism underlying their biological effects and plays a significant role in their translational applications. On one hand, iron metabolism-related indicators can be used to evaluate the biological safety of EMFs. On the other hand, the effects of EMFs on iron metabolism have potential applications in the research and treatment of various diseases, including immunoregulation, cardiovascular diseases, neurological diseases, orthopedic diseases, diabetes, liver injury, and cancer. Moreover, the synergistic regulation of iron metabolism through the combination of EMFs with drugs—particularly iron-based NPs—has notably enhanced the potential of EMFs in the biomedical field. However, conflicting results in many studies regarding the effects of EMFs on iron metabolism make it challenging to systematically summarize the regulatory patterns of EMFs on iron metabolism. Despite these inconsistencies, several general characteristics of EMF-mediated regulation of iron metabolism can still be summarized. First, the effects of EMFs on iron metabolism vary depending on the properties of the EMFs (such as intensity, gradient, frequency, etc.) and exposure time [17]. Additionally, the effects of EMFs on iron metabolism appear to exhibit a biological “window”. It has been reported that the biological effects induced by EMFs of specific intensities, frequencies, and exposure durations within this “window” may differ significantly from those outside it [123]. Furthermore, the regulation of iron metabolism by EMFs also shows significant tissue- and cell-type specificity. Compared to changes in biochemical indicators, biophysical mechanisms are typically used to provide a deeper explanation of the biological effects and patterns of EMFs. In this review, we have summarized several potential biophysical mechanisms through which EMFs regulate iron metabolism, aiming to better understand the underlying reasons for the complex regulatory characteristics of EMFs on iron metabolism, and to provide research insights for utilizing EMFs in the modulation of iron metabolism in the future. Furthermore, the descriptions of EMFs parameters and exposure forms in many studies are often vague, which limits the comparability of results across different studies. Therefore, more precise descriptions of EMF parameters and exposure methods are needed in future research on the effects of EMFs on iron metabolism and related diseases. We expect that with extensive and in-depth studies on the effects and mechanisms of EMFs in iron metabolism, the scope of EMF applications in disease prevention and treatment is likely to expand.

Open access paper: https://www.sciencedirect.com/science/article/pii/S2090123225002887

--

Active matter as the underpinning agency for extraordinary sensitivity of biological membranes to electric fields

Mathew A, Kulkarni Y. Active matter as the underpinning agency for extraordinary sensitivity of biological membranes to electric fields. Proc Natl Acad Sci U S A. 2025 Mar 25;122(12):e2427255122. doi: 10.1073/pnas.2427255122.

Abstract

Interaction of electric fields with biological cells is indispensable for many physiological processes. Thermal electrical noise in the cellular environment has long been considered as the minimum threshold for detection of electrical signals by cells. However, there is compelling experimental evidence that the minimum electric field sensed by certain cells and organisms is many orders of magnitude weaker than the thermal electrical noise limit estimated purely under equilibrium considerations. We resolve this discrepancy by proposing a nonequilibrium statistical mechanics model for active electromechanical membranes and hypothesize the role of activity in modulating the minimum electrical field that can be detected by a biological membrane. Active membranes contain proteins that use external energy sources to carry out specific functions and drive the membrane away from equilibrium. The central idea behind our model is that active mechanisms, attributed to different sources, endow the membrane with the ability to sense and respond to electric fields that are deemed undetectable based on equilibrium statistical mechanics. Our model for active membranes is capable of reproducing different experimental data available in the literature by varying the activity. Elucidating how active matter can modulate the sensitivity of cells to electric signals can open avenues for a deeper understanding of physiological and pathological processes.

Significance

What is the minimum electric field that biological cells can detect? This question has inspired significant research as certain cells and organisms are known to possess extraordinary sensitivity to weak electrical signals. However, existing models that consider this threshold to be the electrical noise arising from equilibrium thermal fluctuations yield puzzling results that are many orders of magnitude larger than experimental measurements. We resolve this paradox by proposing a nonequilibrium statistical mechanics model in which cell membranes can exploit energy-driven active mechanisms to modulate their electrical sensitivity to sense electric fields that are far weaker than the thermal electrical noise. Our study could open avenues for harnessing the sensitivity of biological cells to electric fields for diagnostics and therapeutics.

Conclusion

In summary, we investigated the role of activity in the sensitivity of biological membranes (and cells) to electric fields by way of a nonequilibrium statistical mechanics–based model. Our study reveals that activity, attributed to different sources, can endow the membranes with the ability to sense and respond to electric fields that are far weaker than the thermal noise limit estimated from purely equilibrium statistical mechanics. Our model for active membranes is capable of reproducing different

experimental data available in the literature by varying the activity and thus provides a possible resolution for a long-standing discrepancy between theoretical studies and experimental measurements for the minimum electric field that can be detected by cells. As part of future study, we intend to enrich the model to incorporate electromechanical coupling through flexoelectricity and investigate its role in the interaction of active membranes with electric fields. Understanding electromechanical biological phenomena that may be impacted by increased fluctuations due to active noise in polarization is another interesting direction of study. Elucidating how cells can modulate their interaction with electric signals through activity can proffer deeper insights into physiological processes and open avenues for applications in biotechnology and medicine.

https://pubmed.ncbi.nlm.nih.gov/40117314/

--

Single exposure to near-threshold 5G millimeter wave modifies restraint stress responses in rats

Matsumoto A, Endo I, Ijima E, Hirata A, Kodera S, Ichiba M, Tokiya M, Hikage T, Masuda H. Single exposure to near-threshold 5G millimeter wave modifies restraint stress responses in rats. Environ Health Prev Med. 2025;30:33. doi: 10.1265/ehpm.24-00321.

Abstract

Background: In response to growing concerns about the health effects of quasi-millimeter waves (qMMW) used in 5th-generation wireless systems, conservative whole-body exposure thresholds based on indirect evidence have been proposed. The guidelines define a whole-body average specific absorption rate (WBA-SAR) of 4 W/kg which causes a 1 °C increase in core temperature, as the operational threshold for adverse health effects. To address the lack of direct evidence, we recently reported that a 30-minute exposure to qMMW at 4.6 W/kg resulted in a 1 °C increase in rat core temperature. Here, we further analyzed the near-threshold stress response for the first time, using biological samples from the aforementioned and additional experiments.

Methods: A total of 59 young Sprague-Dawley rats (240-322 g) were exposed to 28 GHz for 40 minutes at WBA-SARs of 0, 3.7, and 7.2 W/kg, under normal (22.5 °C, 45-55% humidity), and heat (32 °C, 70% humidity) conditions. Rats were restrained in acrylic holders for dose control. We repeatedly measured serum and urinary biomarkers of stress response, aggregated the data, and analyzed them using a single statistical mixed model to subtract the effects of sham exposure and between-subject variation.

Results: Sham exposure induced stress responses, suggesting an effect of restraint. After the subtraction of the sham exposure effect, 28 GHz appeared to induce stress responses as evidenced by elevated serum-free corticosterone 1 or 3 days after the exposure, which was more evident in animals with a change in rectal temperature exceeding 1 °C. Urinary-free catecholamines demonstrated an inhibitory property of 28 GHz frequency exposure on the stress response as evidenced by noradrenaline on the day of exposure. Heat exposure enhanced this effect, suggesting a possible role of noradrenaline in heat dissipation by promoting cutaneous blood flow, a notion supported by the correlation between noradrenaline levels and tail surface temperature, a critical organ for heat dissipation.

Conclusions: This study is the first to demonstrate that qMMW whole-body exposure can alter the stress

response as indicated by corticosterone and noradrenaline at near-threshold levels. Our findings may provide insight into the biological basis of the whole-body exposure thresholds in the international guidelines.

Open access paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC12062831/

--

Effects of non-ionizing radiation on the thyroid gland in rats

Sarhad ZS, Ebrahimbabaei A, Tavassoli A, Shojaeifard M. Effects of non-ionizing radiation on the thyroid gland in rats. BMC Res Notes. 2025 May 24;18(1):231. doi: 10.1186/s13104-025-07297-x.

Abstract

Objectives: This study evaluated the effects of non-ionizing electromagnetic radiation on rat thyroid function and histopathology. Forty female and thirty male Sprague-Dawley rats (200-220 g, 2 months old) were exposed to 2.45 GHz Wi-Fi, mobile jammer radiation, or a sham condition. In Group A, male rats were exposed to Wi-Fi or mobile jammers for 2 h daily for two weeks. The devices were located within a one-meter radius of the animal cage, either on or off. In the Sham group, the experimental setup was like the other groups, but the irradiating devices were turned off. Group B included non-pregnant females, pregnant (exposed and control subgroups), and their offspring. Thyroid hormones in the serum were measured, and the histology was microscopically analyzed, focusing on areas of colloid and epithelium in the thyroid follicles.

Results: Compared to the control group, T4 hormone levels were significantly different in male rats exposed to mobile jammer radiation (p-value = 0.037). In group B, significant differences were found solely in the male offspring regarding T3 levels due to jammer exposure (average = 109.00 for male offspring in the experiment vs. average = 65.50 for those in the control, p-value < 0.001). Additionally, histopathological findings indicated significant differences as well. These results highlight a potential link between exposure to electromagnetic radiation and changes in thyroid endocrine and histological parameters. Our findings suggest that ongoing assessment of existing safety guidelines on non-ionizing radiation exposure is necessary, especially concerning its effects on thyroid hormone levels and follicular histology.

Open access paper: https://bmcresnotes.biomedcentral.com/articles/10.1186/s13104-025-07297-x

--

An 1800 MHz Electromagnetic Field Affects Hormone Levels, Sperm Quality, and Behavior in Laboratory Rats

Pawlak K, Bojarski B, Jagusiak W, Wojnar T, Nieckarz Z, Arent Z, Ludwiczak M, Lasko M. An 1800 MHz Electromagnetic Field Affects Hormone Levels, Sperm Quality, and Behavior in Laboratory Rats (*Rattus norvegicus*). *Applied Sciences*. 2025; 15(9):5160. https://doi.org/10.3390/app15095160.

Abstract

In addition to natural electromagnetic fields (EMFs), so-called artificial electromagnetic fields exist in the biosphere, with mobile communications being one of their main sources. This study aimed to determine the impact of EMF at a frequency of 1800 MHz on the concentrations of selected hormones, sperm motility, viability, morphology and behaviors in laboratory rats. We used 28 rats divided into two equinumerous groups: control (n = 14) and experimental (n = 14). The rats in the experimental group were exposed to EMF for 12 weeks (for 10 min, four times daily); at the same time, the control specimens were kept in standard conditions. After 12 weeks, half of each group was killed, while the other half was maintained for another 4 weeks with no EMF emission. Elevated corticosterone levels and decreased thyroid-stimulating hormone levels were observed in the experimental specimens, which persisted for 2 weeks after the cessation of EMF emission. Exposure to EMF also resulted in decreased sperm motility and viability, as well as increased rat anxiety. This study shows that exposure to EMF (1800 MHz) may affect the endocrine status of the body and the behavior and reproductive functions of animals. However, hormonal disorders appear to be reversible.

Excerpts

"The experimental group was exposed to electromagnetic waves for 10 min, four times daily: 8:00–8:10, 14:00–14:10, 18:00–18:10, and 22:00–22:10. The source of electromagnetic radiation was a generator emitting electromagnetic waves in the frequency range of the GSM cellular network (1100–2100 MHz). The average output power of the generator was 330 mW, with an instantaneous maximum power of 2 W. The load for the power stage was a Yagi GSM antenna with omnidirectional characteristics. The distance between the field generator antenna and the animals (depending on their position in the cage) was 30–40 cm."

"The electromagnetic field emitted by the generator within the rat cages was characterized by the following parameters: the average electric field intensity ranged from 2.8 to 5.7 V/m (0.1 V/m), the average magnetic field intensity ranged from 7.2 to 14.0 mA/m (±0.2 mA/m), and the average power density ranged from 17.7 to 63.6 mW/m^2 (±0.5 mW/m^2) at a frequency of 1800 MHz. Throughout the experiment, the EMF background was monitored, with an average value of 0.22 ± 0.07 V/m."

Open access paper: https://www.mdpi.com/2076-3417/15/9/5160

--

Impairment of Oogenesis and Folliculogenesis in Neonatal Rats after Maternal Exposure to Mobile Phones

Yousefi B, Jadidi M, Nabizadeh Z, Amjad MHT, Ardekanian M. Impairment of Oogenesis and Folliculogenesis in Neonatal Rats after Maternal Exposure to Mobile Phones. Reprod Sci. 2025 May 19. doi: 10.1007/s43032-025-01880-0.

Abstract

Today's lifestyle has led to an increase in infertility. The study investigated the effects of cell phone radiation on oogenesis, folliculogenesis, and gestation stage. Also, its impact on neonates' ovarian hormones and their ovarian development during gestation was investigated. Fifty-four virgin female Wistar rats were randomly divided into three groups: the study group (conversation and stand-by mode, n = 24), the control group (stand-by mode, n = 24), and the sham (turn off, n = 6). The study and control

groups were separated into 4 subgroups, including early stage (1st week), mid-stage (2nd week), late stage (3rd week), and all stages (3 weeks). The results showed that the concentration of plasma estrogen and progesterone, ovarian primordial follicle/primary oocyte, the number of primordial follicles, and nuclei diameters in the study and control subgroups decreased significantly in comparison with the sham group. The most significant reduction was observed in subgroups in which mothers were exposed to radiation for a long period during their pregnancy. Compared to the control and sham groups, the number of primordial follicle apoptosis markedly increased in the study subgroups. The pregnant rats showed significant effects in the entire stages of pregnancy, especially during the first stage (the first week) of development, which has not been reported previously. Exposure to cell phones during the different gestation stages probably decreases ovarian hormone secretion and may harm oogenesis. It also inversely increases the apoptosis of primordial follicles. Therefore, the gestational stages and duration of exposure to cell phone radiation affect the risk of ovarian harm.

Conclusion

This study examined the effects of cell phone exposure on estrogen, progesterone, and ovarian follicles in neonatal rats immediately after birth (natal day). Compared to the sham group, the results showed that cell phone radiation in both the study and control groups caused a decrease in the number and diameter of the primordial follicle, primary oocyte, its nucleus, and the amount of estrogen and progesterone hormones in neonatal rats during different and entire weeks of gestation. Also, these radiations cause an increase in apoptosis (atretic) of primordial follicles and a reduction in the diameter of the neonates' ovaries, which depends on the duration of radiation. Meanwhile, for the first time, the impact of mobile phones during different weeks of pregnancy was examined in this study. In sum, the most effects were observed in the third week of pregnancy. The results showed that the first week of pregnancy was also affected by cell phone effects. Although this study has investigated the effects of EMF on infertility in an animal model (rat), these findings can highlight the potential risks of exposure to mobile phone radiation during pregnancy on the future fertility of the fetus in the womb. However, further research is needed to establish a direct link between EMR exposure and infertility. Non-human primate models (monkeys) are recommended to achieve more reliable results.

Open access paper: https://link.springer.com/article/10.1007/s43032-025-01880-0

--

Analyzing the Impact of Occupational Exposures on Male Fertility Indicators: A Machine Learning Approach

Mohammadi H, Khoddam S, Golbabaei F, Dehghan SF. Analyzing the Impact of Occupational Exposures on Male Fertility Indicators: A Machine Learning Approach. Reprod Toxicol. 2025 May 29:108959. doi: 10.1016/j.reprotox.2025.108959.

Highlights

• Combines occupational epidemiology with cutting-edge machine learning for risk stratification
• Provides actionable thresholds for workplace exposure limits (e.g., WBGT ≥ 28°C linked to decreased testosterone levels)

• Bridges methodological gaps in reproductive health research through model interpretability (SHAP, feature importance analysis)

Abstract

Occupational exposures are critical factors affecting workers' reproductive health. This study investigates the impact of magnetic fields, electric fields, whole-body vibration, noise levels, and heat stress on male reproductive indicators using advanced machine learning models. The aim is to identify key risk factors and provide predictive insights into workers' reproductive health over the next decade. Data were collected from 80 male workers in an automobile part manufacturing plant, capturing demographic characteristics, occupational exposures, biochemical markers, hormone levels, and sperm parameters. Five machine learning models logistic regression, bagging classifier, extreme gradient boosting, random forest, and support vector machine were trained and evaluated using 5-fold cross-validation to determine effective predictors of reproductive health outcomes. Exposure to whole-body vibration, magnetic fields, electric fields, and heat stress closely affected free testosterone levels, with SHAP importance indicating: Magnetic Field Exposure (0.339) and Wet Bulb Globe Temperature (0.138). Worker Age (0.244) was the most influential demographic factor negatively impacting Free Testosterone. The XGBoost and random forest achieved the highest AUC (0.99), outperforming other models in predictive accuracy. The Random Forest model Importance (% Increase in MSE) predicted that Electric Field Exposure (5%) and Magnetic Field Exposure (4.7%) would have the most substantial negative impact on Free Testosterone, followed by Worker Age (4.1%). This study underscores the need for targeted interventions, such as improved workplace safety protocols and regular health monitoring, to protect workers' reproductive health.

Conclusion

This study concludes that machine learning, particularly tree-based models like Random Forest and XGBoost, can effectively identify key occupational and demographic factors influencing male reproductive health. Electric and magnetic field exposures, age, work experience, and oxidative stress biomarkers emerged as the most critical predictors. Explainable AI methods revealed complex interactions among these factors. The 10-year forecast highlighted electric field exposure as the most significant long-term risk. These findings emphasize the need for targeted interventions to reduce electromagnetic and vibration exposures and to protect aging workers.

https://pubmed.ncbi.nlm.nih.gov/40449633/

--

A Mini-Review of the Potential Health Impacts of Indoor Radiation Exposure in Companion Animals

Maipas S, Vardiambasis IO, Nikolopoulos CD, Kavantzas N. A Mini-Review of the Potential Health Impacts of Indoor Radiation Exposure in Companion Animals. Environ Health Insights. 2025 May 28;19:11786302251342972. doi: 10.1177/11786302251342972.

Abstract

Poor indoor environmental quality is a major concern for both humans and companion animals-yet its impact on the latter is often overlooked. This mini review sheds light on indoor radiation sources that may affect the health of companion animals. Radon, primarily from the ground and possibly also from

cat litter, naturally occurring radionuclides in feed, radiofrequency radiation from mobile and cordless phones, Wi-Fi networks, pet tracking devices, solar radiation, and various sources of extremely low frequency radiation are all common examples. Indoor radiation, an odorless and tasteless pollutant, may have the potential to negatively impact the health and well-being of companion animals, which are involuntarily exposed to this environmental health risk factor. Therefore, preventive and precautionary measures are necessary.

Open access paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC12120279/

--

Melatonin ameliorates RF-EMR-induced reproductive damage by inhibiting ferroptosis through Nrf2 pathway activation

Wang J, Dong J, Xu Q, Yan S, Wang H, Lei H, Ma X, Yang T, Wang K, Li Z, Wang X. Melatonin ameliorates RF-EMR-induced reproductive damage by inhibiting ferroptosis through Nrf2 pathway activation. Pathol Res Pract. 2025 May 7;270:156003. doi: 10.1016/j.prp.2025.156003.

Abstract

In recent years, there has been increased attention to the deleterious impacts of radiofrequency electromagnetic radiation (RF-EMR) on male reproductive ability, necessitating the exploration of effective protective measures. Melatonin has antioxidant and anti-apoptotic effects, and there is growing evidence of its benefit to the reproductive process. However, the biochemical mechanisms by which melatonin protects against reproductive damage from RF-EMR exposure are unknown. Here, we found that prolonged (8 weeks) exposure to RF-EMR [2.45 GHz; power density, 2.5 W/m^2; whole-body specific absorption rate (SAR), 0.125-0.5 W/kg] induced ferroptosis and oxidative stress in testicular tissue, leading to a decrease of sperm quality in male mice. Notably, the administration of melatonin mitigated the oxidative harm to the testicles and ferroptosis caused by RF-EMR in mice. Mechanistically, melatonin could inhibit ROS production and ferroptosis by stimulating the nuclear factor-erythroid 2-related factor 2 (Nrf2) signaling pathway through its receptors (MT1/MT2). Taken together, these results indicate that melatonin could potentially improve RF-EMR-induced reproductive damage in male mice by blocking ferroptosis through activation of the Nrf2 pathway.

Conclusions

In conclusion, this study demonstrated that RF-EMR harms male reproduction by triggering ferroptosis, a cell death process, in mouse testes. Melatonin, by activating the Nrf2 pathway via MT1/MT2 receptors, suppresses ferroptosis and protects against RF-EMR damage (Fig. 6). These findings suggest melatonin's potential clinical value. Future controlled trials will explore its efficacy in treating male infertility caused by RF-EMR exposure, making melatonin a promising candidate therapy. ...

https://pubmed.ncbi.nlm.nih.gov/40344840/

--

Exposure at 3.6 GHz in Greece Using Data From Frequency-Selective Monitoring Sensors

Iakovidis S, Manassas A, Apostolidis C, Samaras T. 5G EMF Exposure at 3.6 GHz in Greece Using Data From Frequency-Selective Monitoring Sensors. Bioelectromagnetics. 2025 May;46(4):e70008. doi: 10.1002/bem.70008.

Abstract

The introduction of 5G networks has raised public concerns about potential changes in environmental electromagnetic field (EMF) exposure. This study analyzes continuous monitoring data collected over 2 years (August 2022-October 2024) from 13 frequency-selective monitoring sensors located in Greece's five largest cities. Focusing on the 3.6 GHz band, we evaluated trends and weekly variations in EMF levels. Results indicated a gradual increase in EMF exposure at 3.6 GHz, driven by the growing penetration of 5G infrastructure and devices. Notably, this band exhibited higher maximum-to-median power density ratios compared to other frequency bands, attributable to active antenna systems' characteristics and traffic variations. Applying the ICNIRP 2020 guidelines, we found that 30-min averaged values significantly reduced these variations. All measured EMF levels, including maximum values, remained well below Greek and international safety limits. These findings, especially the increasing trend identified for the EMF levels, underscore the importance of continuous monitoring networks for assessing EMF exposure to existing and emerging telecommunications networks and ensuring compliance with safety standards.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.70008

--

Terahertz Radiation Affects the Dynamics of Neurons by Decreasing Membrane Area Ratio

Ma S, Li S, Wang H, Li Y, Lu C, Li X. Terahertz Radiation Affects the Dynamics of Neurons by Decreasing Membrane Area Ratio. Brain Res Bull. 2025 May 6:111373. doi: 10.1016/j.brainresbull.2025.111373.

Abstract

Terahertz radiation at specific frequencies and energies can mediate cellular morphology or function changes by exciting nonlinear resonance effects in proteins or DNA. However, the effects of terahertz radiation on neuronal morphology and function are currently unknown, and the correlation between neuronal morphology and kinetic properties after terahertz radiation remains to be elucidated. In this paper, we first characterized the changes in neuronal morphology by the relative ratio of neuronal cytosol to protruding membrane area. Analyzed the pattern of the influence of terahertz radiation on neuronal morphology and the cumulative effect. On this basis, this paper constructs a kinetic model of neurons regulated by terahertz radiation, investigates the influence law of terahertz radiation on the kinetic properties of neurons, and analyzes the correlation between neuronal morphology and kinetic properties. The results showed that terahertz radiation caused a decrease in the membrane area ratio of neuronal cytosol to protrusion, and this effect started on the first day of terahertz radiation and lasted until the end of terahertz radiation; terahertz radiation changed the neuronal discharge pattern by decreasing the membrane area ratio of neuronal cytosol to protrusion and lowered the frequency of neuronal inter-cluster discharges and amplitude of action potentials, and increased the neuronal intra-cluster discharge. In addition, terahertz radiation can increase the peak value of neuronal postsynaptic currents by decreasing the membrane area ratio. In summary, terahertz radiation can modulate neurons' morphology and change their firing patterns and kinetic properties by affecting their

morphology. These predict that terahertz radiation at specific frequencies and energies can be developed as a novel, molecular-level neuromodulation technique for intervening or treating neuronal degenerative diseases.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0361923025001856?via%3Dihub

--

Exploring the influence of Schumann resonance and electromagnetic fields on bioelectricity and human health

Nelson I. Exploring the influence of Schumann resonance and electromagnetic fields on bioelectricity and human health. Electromagn Biol Med. 2025 May 20:1-11. doi: 10.1080/15368378.2025.2508466.

Abstract

This article explores the relationship between electromagnetic fields (EMF) and biological systems, focusing on the influence of extremely low-frequency electromagnetic frequencies (ELF), particularly Schumann's resonance (SR) at 7.83 hz. Cells and proteins may have evolved to take advantage of frequencies naturally present in the Earth's EMF, potentially enhancing cellular energy levels and affecting resting membrane potential (RMP). Thus, changes in or absence of SR may have adverse effects on the functioning of the whole organism. Bioelectricity, independent of genes, has been shown to modulate health, suggesting the potential for using controlled application of EMF frequencies in treating certain types of cancer or conditions affecting the RMP. Research indicates that human brainwave activity is highly dependent on the SR, implying a correlation between atmospheric electromagnetic frequencies and brain activity. ELF, including SR, appears to modulate cellular calcium influx/efflux, likely via indirect mechanisms involving field-sensitive molecules or radical pairs that affect ion channel behavior which plays a critical role in cell signaling and regulation of various processes. It can also trigger a cascade of molecular events that ultimately lead to the generation of action potentials, affecting consciousness and behavior. The influence of atmospheric electromagnetic frequencies on human brainwave activity, modulation of cellular calcium influx/efflux, and potential effects on cellular energy levels and RMP highlight the significance of ELF in biological systems. However, further research is required to fully understand these mechanisms and their implications for human health and well-being.

Plain Language Summary

This study looks at how very low-frequency electromagnetic fields – especially the Schumann Resonance at 7.83 hz – may affect our health. These natural frequencies come from the Earth’s atmosphere and may play an important role in how our bodies and cells work. The research explores how these signals could influence the electrical activity inside our cells, including how ions like calcium move in and out. This process is essential for brain function, heart rhythm, sleep patterns, and overall health. The study suggests that human cells may have evolved to respond to these natural frequencies. Changes in, or the absence of, these signals – such as when we’re exposed to artificial electromagnetic fields from modern technology – might disrupt important processes in the body. For example, the paper highlights evidence that some low-frequency signals may help stop cancer cells from growing or improve healing, while unnatural frequencies might do harm. A key point is that our brain activity seems to align with Earth’s

natural electromagnetic rhythms. This could explain why changes in these fields can affect our mood, sleep, or concentration. The study also points out that future space missions should consider simulating these natural frequencies to help astronauts stay healthy outside Earth's environment. In short, this research helps us understand how our bodies interact with the invisible electromagnetic environment around us – and how both natural and man-made signals could influence our health in powerful ways.

Conclusions

Solar and geomagnetic activities appear to affect biological organisms in diverse ways. SR appears to be a biophysical mechanism that coordinates such a relationship.

Human EEG is highly dependent on SR, which suggests that the natural electromagnetic frequencies in the Earth's atmosphere, known as SR, influence human brainwave activity. This suggests a correlation between atmospheric electromagnetic frequencies and brain activity.

While earlier models focused on calcium influx as the primary target of ELF fields, more recent findings indicate that weak magnetic fields likely act first on radical pairs within mitochondria. This radical pair mechanism (RPM) may influence reactive oxygen species (ROS) dynamics, which are tightly linked to intracellular calcium signaling via mitochondria-associated membranes (MAMs). These pathways can alter membrane potential, gene expression, and cellular communication.

Given this context, VGCC activation and calcium fluxes may be secondary consequences of mitochondrial redox modulation. Since calcium ions are involved in nearly all critical cellular processes – from apoptosis to signal transduction – these downstream effects can profoundly influence bioelectric regulation and health outcomes.Biological systems may have evolved to be influenced by electromagnetic frequencies in the ELF range that are naturally present in the environment. These frequencies, particularly the SR at 7.83 hz, could potentially enhance cellular energy levels and affect the RMP. Healthy cells typically have a higher RMP than those with dysfunction or disease.

During the silent phase, thalamocortical neurons might be entrained with SR, which suggests that during this phase, the electromagnetic frequencies of SR could potentially synchronize or entrain the activity of thalamocortical neurons, thereby justifying the coordination between SR and EEG.

There are ongoing studies on the potential health effects of exposure to electromagnetic fields. With the information collected from these studies, it seems plausible that unnatural EMF might affect organisms and cells in multiple ways, such as abnormal ion channels that function to coordinate the whole organism, and improper circadian rhythms.

https://www.tandfonline.com/doi/10.1080/15368378.2025.2508466

--

Effect of Occupational Exposure to Low-frequency Electromagnetic Fields on Cataract Development

Validad MH, Mahjoob M, Pishjo M, Diani M, Rakhshandadi T. Effect of Occupational Exposure to Low-frequency Electromagnetic Fields on Cataract Development. J Ophthalmic Vis Res. 2025 Apr 25;20:1-6. doi: 10.18502/jovr.v20.12281.

Abstract

Purpose: Cataracts are the second leading cause of visual impairment worldwide. This study aimed to examine the impact of occupational exposure to low-frequency electromagnetic fields on cataract development.

Methods: One hundred employees of Zahedan Electricity Company participated in this study. They were assigned to four groups based on their level of exposure: regular, operational, operator personnel, and non-exposure. Based on LOCS III grading, the risk of developing different types of cataracts (i.e., nuclear, posterior subcapsular, and cortical) was evaluated for all participants.

Results: The frequency of cataracts was 62.2% in the exposure group (which includes three subgroups: the regular, operational, and operator personnel) and 53.8% in the non-exposure group. There was a significant difference between the study groups in terms of nuclear opacity grading ($P = 0.003$). The correlation between nuclear and posterior subcapsular cataract grading and work experience duration in the exposure group was statistically significant ($P < 0.018$).

Conclusion: This study's findings indicate that exposure to low-frequency electromagnetic fields such as power lines, power plants, and power distribution posts may be a risk factor for cataract development, particularly nuclear cataracts.

https://pubmed.ncbi.nlm.nih.gov/40400659/

--

Carcinogenicity of extremely low-frequency magnetic fields: A systematic review of animal studies

Luukkonen J, Roivainen P, Nieminen V, Naarala J, Mustafa E, Juutilainen J. Carcinogenicity of extremely low-frequency magnetic fields: A systematic review of animal studies. Environ Res. 2025 May 13:121819. doi: 10.1016/j.envres.2025.121819.

Abstract

Possible carcinogenicity of extremely low frequency (ELF) magnetic fields (MFs), associated with the use and transmission of electricity, has been under scientific and public debate for decades. This review aims to provide an update on studies testing carcinogenicity of ELF MFs in experimental animals. Emphasis was placed on identifying possible connections between study characteristics and the results obtained. This review followed the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) statement, and the methodological quality of the studies was evaluated using the Risk of Bias Rating Tool for Human and Animal Studies developed by the National Toxicology Program's Office of Health Assessment and Translation (OHAT). Publication bias was assessed using the caliper test. Fifty-four eligible studies were identified. Despite poor ratings in certain aspects of the risk of bias evaluation, the quality of the studies was generally relatively good, with only four studies receiving the weakest rating. Overall, there was very little evidence that ELF MFs alone are carcinogenic. Evidence of co-carcinogenicity, from studies that have used ELF MFs in combination with other agents, remains inconclusive. A clear indication of publication bias was observed, though it is unlikely that this bias alone explains all reported MF-modified effects. Based on the current literature, future studies on co-carcinogenicity of ELF MFs should employ approaches other than classical initiation-promotion models.

Additionally, well-designed studies to better understand the reported effects of ELF MFs and the underlying mechanisms are warmly welcomed.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0013935125010709

--

Weak anthropogenic electric fields affect honeybee foraging

Mallinson VJ. Woodburn FA, O'Reilly KJ. Weak anthropogenic electric fields affect honeybee foraging. Cell Press. 2025. https://doi.org/10.1016/j.isci.2025.112550.

Highlights

• Weak anthropogenic-like electric fields reduce honeybee floral landing

• AC and positive DC fields deter honeybee landing, but negative DC does not

• Electric fields near power lines match experimental levels, spanning tens of meters

• These findings highlight electric pollution as a potential disruptor of pollination

Summary

Aerial electroreception, the detection of airborne electric fields (E-fields), is an emerging sensory system in arthropods, including bees, which can use floral E-fields as foraging cues. However, the influence of anthropogenic E-fields on these interactions remains underexplored. Through field experiments in urban meadows, we demonstrate that weak anthropogenic E-fields, including alternating current (AC) and direct current (DC) fields, significantly alter honeybee floral landing behaviors. AC and positive DC fields reduced landings by 71% and 53%, respectively, whereas negative DC fields had no statistically significant impact. Measurements of E-fields near high-voltage transmission lines revealed persistent field strengths comparable to those used experimentally, spanning tens of meters at heights relevant for bee foraging. These findings underscore the potential for anthropogenic E-fields to affect and potentially disrupt plant-pollinator interactions, threatening pollination efficiency, a cornerstone of agriculture and biodiversity. Our study highlights the need for advancing research on the ecological impacts of electric pollution.

Open access paper: https://www.cell.com/iscience/fulltext/S2589-0042(25)00811-9

--

Effect of short-term extremely low-frequency electromagnetic field on respiratory functions

Sirinyildiz F, Cesur G, Elmas O, Elmas S, Comlekci S, Yazıcı O, Keskin A, Keskin HE. Effect of short-term extremely low-frequency electromagnetic field on respiratory functions. Rev Assoc Med Bras (1992). 2025 Jun 2;71(4):e20241812. doi: 10.1590/1806-9282.20241812. PMID: 40465999.

Abstract

Objective: Exposure to extremely low-frequency electromagnetic fields can cause harmful or beneficial effects on living organisms. The aim of this study was to examine the potential effects of extremely low-frequency electromagnetic fields on respiratory physiology by investigating possible changes in respiratory function parameters during and after short-term extremely low-frequency electromagnetic field exposure.

Methods: Twenty Wistar albino rats were included in the study, and these rats were randomly divided into two groups: control and electromagnetic field. A noninvasive head-out plethysmography technique was used to accurately assess lung function in rats. Rats in the electromagnetic field group were exposed to electromagnetic field at a frequency of 50 Hz and a magnetic flux density of 0.3 mT for 2 min. Respiratory function parameters of both groups were recorded in three separate periods before, during and after electromagnetic field exposure. Respiratory rate, respiratory cycle duration, inspiration time, expiration time, tidal volume, minute volume, peak inspiratory flow, and peak expiratory flow were measured in these periods.

Results: There was no significant difference in the parameters measured before electromagnetic field exposure between the groups. During the electromagnetic field exposure period, the mean respiratory rate measured in the electromagnetic field group was lower compared to the control group data, while the mean respiratory cycle duration, inspiration time, and tidal volume measured in the electromagnetic field group were higher compared to the control group data. There was no significant difference in the parameters measured after electromagnetic field exposure between the groups.

Conclusion: Short-term extremely low-frequency electromagnetic field exposure decreases respiratory rate and increases respiratory cycle duration, inspiration time, and tidal volume.

Open access paper: https://www.scielo.br/j/ramb/a/9mQPKdGLX8DJ3mPwSqygwTG/?lang=en

--

Electromagnetic wireless remote control of mammalian transgene expression

Lin Z, Guha Ray P, Huang J. *et al.* Electromagnetic wireless remote control of mammalian transgene expression. *Nat. Nanotechnol.* (2025). https://doi.org/10.1038/s41565-025-01929-w.

Abstract

Communication between wireless field receivers and biological sensors remains a key constraint in the development of wireless electronic devices for minimally invasive medical monitoring and biomedical applications involving gene and cell therapies. Here we describe a nanoparticle–cell interface that enables electromagnetic programming of wireless expression regulation (EMPOWER) of transgenes via the generation of cellular reactive oxygen species (ROS) at a biosafe level. Multiferroic nanoparticles coated with chitosan to improve biocompatibility generate ROS in the cytoplasm of cells in response to a low-frequency (1-kHz) magnetic field. Overexpressed ROS-responsive KEAP1/NRF2 biosensors detect the generated ROS which is rewired to synthetic ROS-responsive promoters to drive transgene expression. In a proof-of-concept study, subcutaneously implanted alginate-microencapsulated cells stably expressing an EMPOWER-controlled insulin expression system normalized blood-glucose levels in a mouse model of type 1 diabetes in response to a weak magnetic field.

Open access paper: https://www.nature.com/articles/s41565-025-01929-w

--

Effects of radiofrequency electromagnetic field exposure on cancer in laboratory animal studies, a systematic review

Mevissen M, Ducray A, Ward JM, Kopp-Schneider A, McNamee JP, Wood AW, Rivero TM, Straif K. Effects of radiofrequency electromagnetic field exposure on cancer in laboratory animal studies, a systematic review. Environment International. 2025, doi: 10.1016/j.envint.2025.109482.

Abstract

Background More than ten years ago, the World Health Organization's (WHO) International Agency for Research on Cancer (IARC) published a monograph concluding there was limited evidence in experimental animals for carcinogenicity of Radio Frequency Electromagnetic Field (RF EMF).

Objective The objective of this review was to systematically evaluate the effects of RF EMF exposure on cancer in experimental animals.

Methods

Eligibility criteria: Based on pre-established Populations, Exposures, Comparators, Outcomes, and Study Type (PECOS) criteria, studies in experimental animals of the following study types were included: chronic cancer bioassays, initiation-(co–)promotion studies, and studies with tumor-prone animals.

Information sources: MEDLINE (PubMed), Science Citation Index Expanded and Emerging Sources Citation Index (Web of Science), and the EMF Portal.

Data abstraction and synthesis: Data are publicly available online as interactive visuals with downloadable metadata. We adapted the risk-of-bias (RoB) tool developed by Office of Health Assessment and Translation (OHAT) to include considerations pertinent to the evaluation of RF EMF exposure and cancer bioassays. Study sensitivity was assessed with a tool adopted from the Report on Carcinogens (RoC). We synthesized studies using a narrative approach. Effect size was calculated as the 1% Bayesian Average benchmark dose (BMD) of a respective study when dose–response or a trend was identified (see BMDAnalysisSupplementaryMaterial) (Supplement 1).

Evidence Assessment: Certainty of the evidence (CoE) was assessed using the Grading of Recommendations, Assessment, Developing and Evaluations (GRADE) approach, as refined by OHAT. Evidence from chronic cancer bioassays was considered the most directly applicable to evaluation of carcinogenicity.

Results We included 52 studies with 20 chronic bioassays. No studies were excluded based on risk of bias concerns. Studies were not considered suitable for meta-analysis due to heterogeneity in study design, species, strain, sex, exposure characteristics, and cancer outcome. No or minimal evidence of RF EMF exposure-related cancer outcomes was found in most systems or organs in any study (these included gastrointestinal/digestive, kidney, mammary gland, urinary, endocrine, musculoskeletal, reproductive, and auditory). For lymphoma (18 studies), with 6 chronic bioassays (1,120 mice, 1,780 rats) inconsistency between two chronic bioassays was not plausibly explainable, and the CoE for lymphoma was rated 'moderate'. For brain tumors (20 studies), including 5 chronic bioassays (1,902

mice, 6,011 rats), an increase in glial cell-derived neoplasms was reported in two chronic bioassays in male rats. The CoE for an increased risk in glioma was judged as high. The BMD analysis was statistically significant for only one study and the BMD was 4.25 (95% CI 2.70, 10.24). For neoplasms of the heart (4 chronic bioassays with 6 experiments), 3 studies were performed in rats (~2,165 animals), and 1 in mice (~720 animals). Based on 2 bioassays, statistically significant increases in malignant schwannomas was judged as high CoE for an increase in heart schwannomas in male rats. The BMDs from the two positive studies were 1.92 (95 %CI 0.71, 4.15) and 0.177 (95 %CI 0.125, 0.241), respectively. Twelve studies reported neoplasms in the adrenal gland (5 chronic bioassays). The CoE for an increased risk in pheochromocytoma was judged as moderate. None of these findings were dose-dependent when compared to the sham controls. Sixteen studies investigated tumors of the liver with 5 of these being chronic bioassays. The CoE was evaluated as moderate for hepatoblastomas. For neoplasms of the lung (3 chronic bioassays), 8 studies were conducted in rats (~1,296 animals) and 23 studies in mice (~2,800 animals). In one chronic bioassay, a statistically significant positive trend was reported for bronchoalveolar adenoma or carcinoma (combined), which was rated as moderate CoE for an increase in lung neoplasms with some evidence from 2 initiation-(co–)promotion studies.

Discussion Meta-analysis was considered inappropriate due to the heterogeneity in study methods. The GRADE/OHAT CoE framework has not been frequently applied to animal studies and experience to date suggests refinements are needed. We deferred to standard methods in environmental health where CoE is framed in the context of strength of the evidence providing positive support for carcinogenicity. High CoE can be interpreted as the true effect is highly likely to be reflected in the apparent relationship. Moderate CoE indicates the true effect may be reflected in the apparent relationship. Cancer bioassays conducted in experimental animals are commonly used to identify potential human carcinogens. We note that the two tumor types with high CoE in animals in this systematic review are the same as those identified with limited evidence in humans by the IARC Working Group. However, even in cases where the animal evidence demonstrates high CoE, the extrapolation of risk from cancer bioassays to humans is particularly complex for RF EMF. Without a better understanding of the mechanism of the carcinogenicity of RF-EMF, the choice of exposure metric for risk extrapolation (whole body versus localized), intensity or cumulative exposure, whether or not a monotonic dose–response holds for carcinogenic effects, and whether SAR is the appropriate dose metric for adverse effects induced by RF-EMF may be critical.

Other This review was partially funded by the WHO radioprotection programme. The protocol for this review was registered in Prospero reg. no. CRD42021265563 and published in Environment International 2022 (Mevissen et al. 2022).

Final conclusions

The findings of this systematic review indicate that there is evidence that RF EMF exposure increases the incidence of cancer in experimental animals with the CoE being strongest for malignant heart schwannomas and gliomas.

Despite the high level of certainty that evidence of carcinogenicity in experimental animals may predict a carcinogenic hazard to humans, extrapolation of risk from cancer bioassays to humans is particularly complex for RF EMF. Without an understanding of the mechanism of the carcinogenicity of RF-EMF the choice of exposure metric for risk extrapolation (whole body versus localized), intensity or cumulative

exposure whether or not a monotonic dose–response holds for carcinogenic effects, and whether SAR is the appropriate dose metric for adverse effects induced by RF-EMF may be critical.

Besides the integration of the sensitivity domain adopted from the RoC, more work is needed to tailor the GRADE approach to assessing the CoE from animal cancer bioassays designed to identifying risks of environmental agents.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412025002338

--

The association of widely used electromagnetic waves exposure and pregnancy and birth outcomes in Yazd women: a cohort study

Razavimoghadam M, Sefidkar R, Ehrampoush MH, Teimouri F, Hassanabadi MHZ, Nokhostin F. The association of widely used electromagnetic waves exposure and pregnancy and birth outcomes in Yazd women: a cohort study. BMC Pregnancy Childbirth. 2025 Apr 11;25(1):427. doi: 10.1186/s12884-025-07512-4.

Abstract

Background: The present study investigated the association between exposure to electromagnetic waves from widely used devices and pregnancy and birth outcomes among women in Yazd City.

Methods: This study was conducted on a total of 1,666 participants enrolled in the Yazd Mother and Child Cohort Center between 2015 and 2019. Cell phones, cordless phones, and Wi-Fi instruments were considered electromagnetic wave sources. Pregnancy and birth outcomes examined in this study include miscarriage, preterm labor, abnormal birth weight, and deviations in newborn height and head circumference. SPSS version 24 and R-studio version 4.3.1 software were used for statistical analysis.

Results: The current study revealed that 41 (2.5%) mothers had miscarriage and 174 (10.4%) mothers experienced preterm labor. Furthermore, 181 (10.9%) infants with abnormal birth weights, 117 (7%) infants with abnormal height, and 124 (7.4%) infants with abnormal head circumference were observed in this study. Individuals with longer cell phones call duration during pregnancy had higher risk of miscarriage ($p < 0.001$, RR (95% CI), 1.0061(1.003-1.0093)), abnormal birth weight ($p = 0.002$, RR (95% CI), 1.0012 (1.0004-1.002)) and height ($p = 0.003$, RR (95% CI), 1.0014 (1.0004-1.0023)) in comparison with those with lower cell phone call duration in this period.

Conclusion: The findings of the study revealed that an increase in exposure to cell phones during pregnancy can increase the risk of pregnancy and birth outcomes, namely miscarriage, abnormal weight and height of the infant. Furthermore, cordless phone conversations could increase the risk of abnormal weight in newborns.

Open access paper: https://bmcpregnancychildbirth.biomedcentral.com/articles/10.1186/s12884-025-07512-4

--

Effects of Mobile Electromagnetic Exposure on Brain Oscillations and Cortical Excitability: Scoping Review

Torkan A, Zoghi M, Foroughimehr N, Yavari A, Jaberzadeh S. Effects of Mobile Electromagnetic Exposure on Brain Oscillations and Cortical Excitability: Scoping Review. *Sensors*. 2025; 25(9):2749. doi: 10.3390/s25092749.

Abstract

With the widespread adoption of smartphones, concerns about increased exposure to non-ionizing radiofrequency have emerged. This scoping review examines the effects of mobile phone exposure on neural oscillations and cortical excitability, focusing on both motor and non-motor regions of the cerebral cortex. A scoping review identified seventy-eight studies that involved healthy individuals and employed electroencephalography and only two studies that investigated transcranial magnetic stimulation as primary technical tools. The findings suggest that mobile phone exposure may affect brain oscillations and cortical excitability. However, inconsistencies in experimental methods across studies make it difficult to draw definitive conclusions. Additionally, research on fifth-generation technology, particularly mmWave exposure from next-generation mobile networks, remains limited and needs further exploration. These gaps highlight the need for more in-depth studies on how mobile phone exposure impacts brain function.

Conclusions and Future Work

Our literature review indicates that mobile EM exposure can induce changes in brain oscillations, primarily increasing the amplitude of alpha, beta, theta, and delta bands, particularly in the eyes closed condition. This review suggests that mobile EM exposure may enhance CSE due to strengthened facilitator mechanisms and weakened inhibitory mechanisms. Despite extensive research over the years, several gaps remain in the literature, which require further investigation, particularly regarding the effects of 5G as an emerging technology. This is linked to the non-thermal effect even at exposure levels below current thermal thresholds. These findings underscore the need to revise SAR limits and to incorporate neurophysiological EEG and TMS metrics as non-invasive measures. Furthermore, the advent of 5G technology, with its use of higher frequency mmWave bands (>24 GHz), demands urgent investigation into potential unique tissue interactions and indirect neural impacts, which warrants precautionary measures until robust safety data are available. The conclusions of this review may be judged as consistent with the evidence. However, they should be treated with caution due to key limitations, such as limited TMS studies (only two), which resulted in a lack of data and some methodological and technical problems in EEG studies, as mentioned in the discussion and limitations section. Additionally, the lack of research verifying 5G frequencies above 6 GHz prevents definitive claims.

The following recommendations are suggested for future research in this field: (1) Conduct more comprehensive examinations of the effects of 5G mobile EM exposure on brain oscillations and brain excitability. This requires additional experimental research, including exposure systems operating at 5G frequencies and post-exposure assessments. (2) Study the impact of age factors on brain structure and physiology by assessing mobile EM exposure effects on both young and older adults to better understand age-related differences. (3) Investigate potential gender effects of mobile EM exposure on brain structure and function. (4) Enhance the precision of future research by refining methodological

aspects, from data acquisition to result analysis, in EEG and TMS studies. This includes careful attention to study design, optimal sample size determination, and comprehensive study setup, all contributing to improved accuracy in future investigations. (5) Standardize exposure parameters (frequency, SAR, distance from the head), sham/control conditions (double-blind protocols), and reporting EEG/TMS technical details (electrode placement, filtering) in EEG and TMS protocols, which are critical updates to international EM exposure guidelines.

Open access paper: https://www.mdpi.com/1424-8220/25/9/2749

--

Sleep and Arousal Hubs and Ferromagnetic Ultrafine Particulate Matter and Nanoparticle Motion Under Electromagnetic Fields: Neurodegeneration, Sleep Disorders, Orexinergic Neurons, and Air Pollution in Young Urbanites

Calderón-Garcidueñas L, Cejudo-Ruiz FR, Stommel EW, González-Maciel A, Reynoso-Robles R, Silva-Pereyra HG, Pérez-Guille BE, Soriano-Rosales RE, Torres-Jardón R. Sleep and Arousal Hubs and Ferromagnetic Ultrafine Particulate Matter and Nanoparticle Motion Under Electromagnetic Fields: Neurodegeneration, Sleep Disorders, Orexinergic Neurons, and Air Pollution in Young Urbanites. Toxics. 2025 Apr 8;13(4):284. doi: 10.3390/toxics13040284.

Abstract

Air pollution plays a key role in sleep disorders and neurodegeneration. Alzheimer's disease (AD), Parkinson's disease (PD), and/or transactive response DNA-binding protein TDP-43 neuropathology have been documented in children and young adult forensic autopsies in the metropolitan area of Mexico City (MMC), along with sleep disorders, cognitive deficits, and MRI brain atrophy in seemingly healthy young populations. Ultrafine particulate matter (UFPM) and industrial nanoparticles (NPs) reach urbanites' brains through nasal/olfactory, lung, gastrointestinal tract, and placental barriers. We documented Fe UFPM/NPs in neurovascular units, as well as lateral hypothalamic nucleus orexinergic neurons, thalamus, medullary, pontine, and mesencephalic reticular formation, and in pinealocytes. We quantified ferromagnetic materials in sleep and arousal brain hubs and examined their motion behavior to low magnetic fields in MMC brain autopsy samples from nine children and 25 adults with AD, PD, and TDP-43 neuropathology. Saturated isothermal remanent magnetization curves at 50-300 mT were associated with UFPM/NP accumulation in sleep/awake hubs and their motion associated with 30-50 μT (DC magnetic fields) exposure. Brain samples exposed to anthropogenic PM pollution were found to be sensitive to low magnetic fields, with motion behaviors that were potentially linked to the early development and progression of fatal neurodegenerative diseases and sleep disorders. Single-domain magnetic UFPM/NPs in the orexin system, as well as arousal, sleep, and autonomic regions, are key to neurodegeneration, behavioral and cognitive impairment, and sleep disorders. We need to identify children at higher risk and monitor environmental UFPM and NP emissions and exposures to magnetic fields. Ubiquitous ferrimagnetic particles and low magnetic field exposures are a threat to global brain health.

Open access paper: https://www.mdpi.com/2305-6304/13/4/284

--

Understanding Electromagnetic Hypersensitivity (EHS) From Mobile Phone Radiofrequency Radiation (RFR) Exposure: A Mixed-Method Study Protocol

Razak NSZM, Rahman AA, Minhat HS, Fauzi FA. Understanding Electromagnetic Hypersensitivity (EHS) From Mobile Phone Radiofrequency Radiation (RFR) Exposure: A Mixed-Method Study Protocol. Bioelectromagnetics. 2025 May;46(4):e70007. doi: 10.1002/bem.70007.

Abstract

More than 7 billion mobile phone users were recorded worldwide in 2022. Radiofrequency radiation (RFR) from mobile phones can produce radiobiological effects, which may lead to electromagnetic hypersensitivity (EHS) in humans. This study employs a mixed-method approach: the quantitative component identifies predictors of EHS from mobile phone RFR using the biopsychosocial model, whereas the qualitative component explores individual experiences that contribute to EHS. Quantitative analysis is conducted using the Statistical Package for the Social Sciences (SPSS) version 25.0 for descriptive and analytical statistics, whereas qualitative analysis is carried out with NVivo version 14 for thematic analysis. Findings from both analyses are compared and discussed to highlight similarities and differences, providing a richer, in-depth understanding of EHS resulting from mobile phone RFR exposure among undergraduate students. The study sample consists of 351 undergraduate students aged 18-25 from Universiti Putra Malaysia (UPM), conducted between April 2023 and December 2023. Approval was obtained from the Ethics Committee for Research Involving Human Subjects at UPM before the study. The quantitative component uses a proportionate simple random sampling method with a validated questionnaire, whereas the qualitative component utilizes purposive sampling through in-depth interviews. Enhancing understanding of EHS may contribute to new knowledge and raise awareness of its potential effects on the population.

https://pubmed.ncbi.nlm.nih.gov/40293416/

--

Temporal Change of Outdoor Rf-Emf Levels Four in European Countries: A Microenvironmental Measurement Study

Belackova L, Veludo AF. Aminzadeh R, Van Bladel H, Griffon V, Cardis E. et al. Temporal Change of Outdoor Rf-Emf Levels Four in European Countries: A Microenvironmental Measurement Study. Available at SSRN: https://ssrn.com/abstract=5146655 or http://dx.doi.org/10.2139/ssrn.5146655.

Abstract

Introduction Over the past two decades, the amount of transmitted mobile data have increased rapidly. It is unknown whether the implementation of the new technologies enabling this has resulted in changes of outdoor radio-frequency electromagnetic fields (RF-EMF) exposure. Therefore, microenvironmental measurements were used to investigate temporal trends in RF-EMF exposure between 2016 and 2023, in the Netherlands, Switzerland, Belgium and Spain, following a similar protocol across campaigns. This allowed us to compare exposure trends between countries and years.

Methods The data was collected as part of the ACCEDERA (2016-2018), ETAIN (2023), and GOLIAT (2023) projects, walking repeatedly the same routes with RF-EMF exposimeters. Identical

microenvironments were identified in each country and measurements of the exposure of mobile base station, contribution of mobile phones and the total levels were compared across years.

Results Comparing measurements between 6 to 14 unique microenvironments in each country, our data did not suggest significant changes in the exposure from the mobile base station origin (total downlink exposure) between 2016 and 2023 for the four countries. Across all countries the median values of the mobile base station exposure ranged from 0.12 mW/m2 in Switzerland and 0.24 mW/m2 in Belgium in 2016 to 0.28 mW/m2 in the Netherlands in 2023, 0.20 mW/m2 in Belgium, 0.11 mW/m2 in Switzerland and 0.19 mW/m2 in Spain in 2023. There was no consistent trend in the individual microenvironments across the countries.

Conclusions Our measurements of across included environment groups RF-EMF outdoor exposure levels do not indicate change in exposure levels between 2016 and 2023 despite an increase in mobile data traffic by a factor of 8 in Western Europe.

https://papers.ssrn.com/sol3/papers.cfm?abstract_id=5146655

--

Assessing EMF Exposure in Greek Urban and Suburban Areas During 5G Deployment: A Focus on 5G EMF Levels and Distance Correlation

Manassas A, Christopoulou M, Papanikolaou N, Delidimitriou S, Karabetsos E, Samaras T. Assessing EMF Exposure in Greek Urban and Suburban Areas During 5G Deployment: A Focus on 5G EMF Levels and Distance Correlation. *Electronics*. 2025; 14(8):1554. doi: 10.3390/electronics14081554

Abstract

This study presents an extensive analysis of electromagnetic field (EMF) exposure in Greek urban and suburban areas during the ongoing deployment of the 5G technology. A total of 400 in situ measurements of electric field strength were conducted at ground level across four distinct regions. The first part of the study provides a statistical analysis of the overall EMF exposure, highlighting the contributions of different cellular generations (2G–5G) to the total field. The results show that 4G networks are the dominant contributors, while the impact of 5G remains limited. The second part investigates the correlation between electric field levels in the 3.5 GHz band and the distance from 5G base stations (BS). Using geospatial analysis, we computed distances to the nearest BS and examined their relationship with measured EMF levels. The findings indicate a general inverse correlation between EMF levels and BS distance, although variations occur due to factors such as line-of-sight conditions and urban built environment characteristics. These results confirm that distance remains a key predictive feature for AI-based EMF exposure models, even in the case of 5G networks.

Conclusions

This study presents a descriptive statistical analysis of electric field measurements in four different regions in Greece, focusing on both the existing cellular networks and the emerging 5G technology. The findings reveal that electric field levels vary considerably across different environments, with urban areas exhibiting higher values due to greater population density and a higher concentration of BS. Cellular networks, especially 4G, contribute significantly to the overall electric field levels, whereas the 5G network shows only a minor contribution. Recording the electric field at the 3.5 GHz band without

generating data traffic and examining its correlation with distance from BS, we observed that, in general, as distance increased, the recorded electric field decreased. Our results confirm that distance from BS antennas should be included as a predictive feature in AI models for EMF levels, not only for 2G, 3G, and 4G networks, but also for 5G. This finding highlights the enduring significance of distance as a key factor for accurately predicting EMF levels in the 5G band, where its influence on field strength is clearly observed.

Open access paper: https://www.mdpi.com/2079-9292/14/8/1554

--

Impact of expectancy on fatigue by exposure to the fifth generation of mobile communication signals

Yang L, Ding X, Zhang S, Wu T. Impact of expectancy on fatigue by exposure to the fifth generation of mobile communication signals. Electromagn Biol Med. 2025 Apr 23:1-12. doi: 10.1080/15368378.2025.2496151.

Abstract

There is a long-standing debate about the relationship between Radio Frequency Electromagnetic Field (RF-EMF) exposure and fatigue. Past studies primarily rely on self-report scales to assess fatigue, but these methods are often susceptible to personal biases. Notably, the role of psychological factors in the fatigue response induce by RF-EMF exposure remains unclear. Therefore, our study focuses on exploring the impact of 5G signal exposure on human fatigue, particularly considering the influence of expectancy induced by psychological priming on the outcomes. In this study, we recruited 21 healthy subjects who were tested in three sessions. Each session included two 30-min exposures to either real or sham 5 G signals, with the order randomized. The experiment was conducted under varying informational conditions: subjects were provided with correct, false, or no information about the order of exposure. Additionally, subjects completed a fatigue scoring questionnaire and underwent Electroencephalogram (EEG) measurements during the experiment. The statistical comparison indicates that 5 G RF-EMF exposure at routine levels does not lead to changes in EEG power. The finding reveals that the report of fatigue can be altered by the conveyed information of being exposed by 5 G signals although there is no real exposure and no detectable electrophysiological indicator. Our findings suggest that it is necessary to prevent psychological priming in any kind or to take its possible consequence into consideration, to reveal this effect of RF-EMF exposure.

https://pubmed.ncbi.nlm.nih.gov/40269539/

--

Determining the relationship between mobile phone network signal strength and RF-EMF exposure: protocol and pilot study to derive conversion functions

Sandoval-Diez N, Belácková L, Fernandes Veludo A, Jalilian H, Guida F, Deltour I, Thielens A, Zahner M, Fröhlich J, Huss A, Röösli M. Determining the relationship between mobile phone network signal strength and radiofrequency electromagnetic field exposure: protocol and pilot study to derive conversion functions. Open Res Eur. 2025 Mar 31;4:206. doi: 10.12688/openreseurope.18285.2.

Abstract

Mobile phones continuously monitor and evaluate indicators of the received signal strengths from surrounding base stations to optimise wireless services. These signal strength indicators (SSIs) offer the potential for assessing radiofrequency electromagnetic field (RF-EMF) exposure on a population scale, as they can be related to exposure from both base stations and handset devices. Within the ETAIN (Exposure To electromAgnetic fields and plaNetary health) project, an open-access RF-EMF exposure app for smartphones, named "ETAIN 5G-Scientist", has been developed using citizen science. This paper delineates a measurement protocol for deriving formulas to convert the app SSIs into electric field values to estimate RF-EMF exposure. It presents pilot study results from measurements taken at four locations in France (FR), and 14 locations in the Netherlands (NL), using three different phone models and the most common network providers in each country. The measurements were conducted while executing different usage scenarios, such as calls or data transmission. The exposimeter ExpoM-RF4 and on-body electric field probes were used to measure exposure from far-field sources and the handset, respectively. Two-minute aggregates were considered the sample unit for analyses (n=891 in NL, n=395 in FR). Regression analyses showed a positive log-linear relationship between Long Term Evolution (LTE) Received Signal Strength Indicator (RSSI) and far-field RF-EMF exposure when aggregating data by location (coefficients for normalised RSSI: 0.91 [95% CI: 0.55 - 1.28] in FR, 1.09 [95% CI: 0.96 - 1.22] in NL). Negative log-linear trends were observed for handset-related RF-EMF exposure at the ear (-0.31 [95% CI: -0.46 - -0.16]) and chest (-0.20 [95% CI: -0.37 - -0.03]) during data transmission scenarios. These results demonstrate that the ETAIN 5G-Scientist app can be implemented for smartphone-based RF-EMF estimation. However, uncertainties in individual measurement points highlight the need for further data collection and analysis to improve the accuracy of exposure estimates.

Plain language summary

In this study, we looked at how mobile phones can help us measure exposure to radiofrequency electromagnetic fields (RF-EMF) from mobile wireless communication. As part of the ETAIN project (Exposure To electromAgnetic fields and plaNetary health), we developed an app called "5G Scientist" to keep track of the signal strength indicators (SSIs) that smartphones continuously use to establish and improve wireless connections. Our goal is to study how these indicators are related to RF-EMF exposure and to develop formulas for converting them into exposure values. To achieve this, we developed a measurement protocol and tested it in a pilot study. In this pilot study, we took spot measurements in France and the Netherlands using various phone models and network providers, while applying different usages of the phone, e.g. voice or video calls and data transmission. We used a personal exposimeter to measure exposure from far-field sources (such as base stations) and on-body electric field probes to measure exposure from near-field sources (such as own phone use). Our measurements indicate that as signal quality improves, exposure from mobile phone base stations increases, whereas exposure from one's own device decreases in both strength and duration. These results suggest that smartphone apps could serve as valuable tools for large-scale exposure assessment, as they show the potential to capture ambient exposure as well as exposure coming from the own mobile phone. However, more extensive measurement campaigns are needed to make this method more accurate.

Open access paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC12032521/

--

Chicken or egg? Attribution hypothesis and nocebo hypothesis to explain somatization associated to perceived RF-EMF exposure

Ariccio S, Traini E, Portengen L, Martens A, Slottje P, Vermeulen R, Huss A. Chicken or egg? Attribution hypothesis and nocebo hypothesis to explain somatization associated to perceived RF-EMF exposure. Front Public Health. 2025 Apr 9;13:1561373. doi: 10.3389/fpubh.2025.1561373.

Introduction: The aim of this study is to understand the temporal relationship between the somatization usually attributed to RF-EMFs, and to evaluate the attribution hypothesis and the nocebo hypothesis in this context.

Method: In this longitudinal study, data from the Dutch Occupational and Environmental Health Cohort Study (AMIGO) was analyzed, consisting of a baseline questionnaire collected in 2011 (14,829 participants) and a follow-up questionnaire collected in 2015 (7,904 participants). Participants completed a questionnaire providing information on their health status, perceived environmental exposures, and demographics. Two sets of multiple regressions were conducted to evaluate the two hypotheses.

Results: Results show that the attribution hypothesis overall explained symptom reporting in association to perceived RF-EMF base station exposure and perceived electricity exposure more frequently than the nocebo hypothesis.

Discussion: This finding stands out from most of the existing literature, which primarily points to the nocebo effect as the main explanation for somatization in response to RF-EMF exposure. While this does not exclude, in absolute terms, the existence of a nocebo effect, potentially at other time scales, this finding has relevant consequences at the policy making level. The emerging relevance of the attribution hypothesis moves the focus on the discomfort of people with unexplained symptoms and their need to find a plausible explanation for their discomfort.

Open access paper: https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1561373/full

--

Electromagnetic fields regulate iron metabolism: From mechanisms to applications (Review)

Wang S, Yang J, Zhen C, Wang H, Shang P. Electromagnetic fields regulate iron metabolism: From mechanisms to applications. J Adv Res. 2025 Apr 29:S2090-1232(25)00288-7. doi: 10.1016/j.jare.2025.04.044.

Abstract

Background: Electromagnetic fields (EMFs), as a form of physical therapy, have been widely applied in biomedicine. Iron, the most abundant trace metal in living organisms, plays a critical role in various physiological processes, and imbalances in its metabolism are closely associated with the development and progression of numerous diseases. Numerous studies have demonstrated that EMF exposure induces significant changes in both systemic and cellular iron metabolism.

Aim of review: This review aims to summarize the evidence and potential biophysical mechanisms underlying the role of EMFs in regulating iron metabolism, thereby enhancing the understanding of their

biological mechanisms and expanding their potential applications in biomedical fields.

Key scientific concepts of review: In this review, we have synthesized research findings and proposed the hypothesis that the biophysical mechanisms of EMFs regulate iron metabolism involve the special electromagnetic properties of iron-containing proteins and iron-enriched tissues, as well as the modulation of membrane structure and function, ion channels, and the generation and activity of Reactive Oxygen Species (ROS). Then, the review summarizes the latest advances in the effects of EMFs on iron metabolism and their safety, as well as their impact on immunoregulation, cardiovascular diseases, neurological diseases, orthopedic diseases, diabetes, liver injury, and cancer.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S2090123225002887?via%3Dihub

--

Efficient design of electromagnetic field exposure maps with multi-method evolutionary ensembles

Guillén-Pina J, Pérez-Aracil J, Chocano-del-Cerro R, Sánchez-Montero R, López-Espí P-L, Salcedo-Sanz S. Efficient design of electromagnetic field exposure maps with multi-method evolutionary ensembles. Environmental Research, 2025, doi: 10.1016/j.envres.2025.121636.

Abstract

Radio-frequency electromagnetic field (EMF) exposure is a growing concern among the population. This concern has led to a need for practical tools to contribute to an adequate risk perception. Representing spatial variations after measurements from fixed sites and interpolating using different techniques is the most suitable method for obtaining EMF high-quality exposure maps. This paper uses evolutionary computation to obtain the optimal set of points to construct high-quality electromagnetic field exposure maps (minimizing an error measure with respect to a reference exposure map). A multi-method ensemble evolutionary approach, able to combine different search operators in a single population (PCRO-SL), is introduced for this particular problem, and it has been tested over actual measurements at Meco town, Madrid, Spain, obtaining good quality electromagnetic field exposure map reconstructions in terms of the differences with a reference EMF exposure map. The results obtained show that reducing the number of measurement points necessary to obtain significant exposure maps, while maintaining their representativeness, is possible.

Conclusions

In this work, an evolutionary multi-method ensemble algorithm for optimization (PCRO-SL) has been used to determine the best points that allow a true representation of the EMF exposure values to be obtained in an ur- ban area. This choice makes it possible to optimize the effort involved in producing these type of maps, for which measurements are taken from fixed locations. The density of points required has been reduced from the initial 8-10 points/km^2 to just over 5 points/km^2.

The division of the territory to be sampled into 250-meter squares can be useful to establish a first analysis of the area, but it is necessary to determine those points of largest interest in order to optimize the sampling effort. The analysis of the results obtained with the proposed multi-method ensemble for

optimization, has made possible to establish the following criteria for the preparation of future exposure maps:

- At least one point in direct line-of-sight (LOS) conditions shall be cho- sen within a first radius of 250 meters.
- Within a second radius of 500 meters, sampling points should be chosen in different radial directions, regardless of their LOS/NLOS status, to determine the rate of variation in the different directions.
- If they are not included in the above criteria, points should be chosen on the perimeter of the surface to be analyzed to avoid artifacts due to interpolation at the extremes of the exposure map.

The above criteria allow obtaining a spatial representation of EMF expo- sure while also maintaining the basic statistical properties of the chosen set of points: mean, variance and approximation to the lognormal distribution.

The effort required to produce these type of maps has been reduced using an evolutionary-type algorithm. This has make possible to obtain exposure maps leading to an adequate perception of the risk of exposure by the general public. For future work, the specific analysis depending on the type of mobile technology or radio service, and the comparison of the results obtained when the measurements are taken by means of exposure meters or other meters are still pending.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0013935125008874

--

High Radiofrequency Radiation in the Surroundings of 10 Schools in Örebro, Sweden

Hardell L, Nilsson M. High Radiofrequency Radiation in the Surroundings of 10 Schools in Örebro, Sweden. Fortune Journal of Health Sciences. 8 (2025): 306-310.

Abstract

In Sweden as well as in many other countries the fifth generation, 5G, for wireless communication is rolled-out since 2019/2020. This has caused increasing environmental exposure to radiofrequency (RF) radiation. Children and adolescents are especially vulnerable to RF radiation. Exposure sources in schools such as Wi-Fi routers, wireless connected mobile phones, and computers may cause high RF-exposure inside the schools but also outside sources, such as mobile phone base stations, may contribute significantly. In this study outside environmental exposure to RF radiation from 4G and 5G base stations was measured in October 2024 in close proximity to 10 schools in Sweden. The maximum levels ranged from 10,716 to 68,452 μ/m^2. These levels far exceed the EUROPAEM EMF guidelines for daytime RF radiation exposure 10-1,000 μW/m^2, nighttime 1-100 μW/m2, and for sensitive persons 0.1-10 μW/m^2.

Excerpts

The measurements were made with the Narda broadband field meter NBM-550, with the probe EF-1891, measuring frequencies between 3 MHz-18 GHz and measurement range 0.6 V/m (955 μW/m2) to

65 V/m (11,206,897 µW/m2). This meter shows results in root mean square (RMS) for both minimum, maximum and average RF radiation level. The frequencies used for 5G in city environments in Sweden are in general around 3.5 GHz while frequencies used for 4G are primarily around 2.6 GHz (https://www.induo.com/b/lte-band-mobilt-bredband/)...

The Narda meter RMS results were considerably lower than Safe and Sound maximum peak exposure values. As one example, at two different spots, the Safe and Sound meter showed max peak levels of 749,000 µW/m 2 (sleeping room, desk, daughter) and 504,000 µW/m2 (sleeping room, pillow, daughter). The Narda RMS maximum levels at the same spots were 137,889 µW/m2 and 88,616 µW/m2, respectively (22). Both measurements were made at the same time.

Open access paper: https://cdn.fortunejournals.com/articles/high-radiofrequency-radiation-in-the-surroundings-of-10-schools-in-oumlrebro-sweden.pdf

--

The effect on rat peripheral nerve morphology and function of a 900-MHz electromagnetic field applied in the prenatal period

Hancı H, Yenilmez E, Demir S, Yıldırım M, Gedikli Ö, Kaya H. (2025). The effect on rat peripheral nerve morphology and function of a 900-MHz electromagnetic field applied in the prenatal period. *Electromagnetic Biology and Medicine*, 1–16. doi: 10.1080/15368378.2025.2479517

Abstract

The purpose of this study was to investigate the effects of a 900 megahertz (MHz) electromagnetic field (EMF) applied in the prenatal period on rat peripheral nerve morphology, nerve conduction velocity, and locomotor activity. Nine pregnant Sprague Dawley rats were assigned into three groups. No procedure was applied to the first group (control). The second (sham) group was placed inside an EMF cage for 1 h a day throughout the experiment (days 1–21 of pregnancy), but was not exposed to EMF. The third group (EMF) was placed inside the EMF cage for 1 h a day throughout the experiment (days 1–21 of pregnancy) and exposed to a 900 MHz EMF. No procedure was applied to the newborn pups until postnatal (PN) day 21, and new groups were constituted from among these. All the newly established groups were subjected to the open field and rotarod tests on PN days 21 and 60, after which electrophysiological measurements were performed on the groups in line with the study protocol. Sciatic nerves obtained from the animals sacrificed on PN day 60 were subjected to histopathological, histomorphometric, immunohistochemical, and biochemical analyses. In light of the study results, we concluded that prenatal application of a 900 MHz EMF adversely affects rat peripheral nerve development, and that these effects persist up to adulthood such as to be detectable in the sciatic nerve morphology, but that these morphological changes are not sufficiently severe to affect functional associated with the sciatic nerve.

Plain Language Summary

This study explores the impact of a 900 MHz electromagnetic field (EMF) applied during the prenatal period on the peripheral nerve morphology, nerve conduction velocity, and locomotor activity of rats. Nine pregnant Sprague Dawley rats were divided into three groups: control, sham, and EMF. The control group received no treatment, the sham group was placed in an EMF cage without exposure, and the

EMF group was exposed to the 900 MHz EMF for 1 hour daily throughout pregnancy. The offspring were tested on postnatal days 21 and 60 using open field and rotarod tests, followed by electrophysiological assessments. Sciatic nerves from the sacrificed animals on postnatal day 60 underwent histopathological, histomorphometric, immunohistochemical, and biochemical analyses. The results indicated that prenatal EMF exposure negatively affects peripheral nerve development in rats, with these effects persisting into adulthood, though not severe enough to impair functional outcomes associated with the sciatic nerve. Additionally, the study provides context on the evolution of communication technologies, emphasizing the increasing use of EMF-emitting devices like cell phones. Given that these devices emit EMF even in standby mode, understanding their potential effects on peripheral nerve integrity is crucial, especially considering the vital role of Schwann cells in nerve development and function. The findings underscore the need for further investigation into the long-term impacts of EMF exposure on peripheral nerve health, particularly during critical developmental periods.

https://doi.org/10.1080/15368378.2025.2479517

--

Experimental Study of Animal Behavior under the Influence of the Electromagnetic Field of the 5G Mobile Communication Standard

Nikitina VN, Kalinina NI, Dubrovskaya EN *et al.* Experimental Study of Animal Behavior under the Influence of the Electromagnetic Field of the 5G Mobile Communication Standard. *Biol Bull Russ Acad Sci* **51**, 3473–3480 (2024). doi: 10.1134/S1062359024701875.

Abstract

A pilot experimental study was carried out to assess the effect of unmodulated and modulated electromagnetic fields of the 5G mobile communication standard on the central nervous system of animals. Mature male Wistar rats were exposed to electromagnetic fields with a frequency of 4.9 GHz and an intensity of 250 μW/cm^2 for 15 days for two hours a day. The animals of the two irradiated groups and the control group were in identical conditions in a semi-anechoic shielded chamber. The effect of the electromagnetic field on the functional state of the central nervous system was assessed by a comprehensive analysis of behavior patterns in the “open field” test before and after irradiation. The study did not reveal clear differences in animal behavior when exposed to unmodulated and modulated 5G electromagnetic fields. Statistically significant changes in animal behavior were recorded in the dynamics of the experiment in two irradiated and control groups, which may be due to a change in the natural electromagnetic background in a semi-anechoic shielded chamber.

https://link.springer.com/article/10.1134/S1062359024701875

--

Effect of Repeated Exposure to Complexly Organized Electromagnetic Radiation on the Rat Behavior in the “Open Field” Test

Kolganova OI, Panfilova VV, Izmestyeva OS *et al.* The Effect of Repeated Exposure to Complexly Organized Electromagnetic Radiation on the Rat Behavior in the “Open Field” Test. *Biol Bull Russ Acad Sci* **51**, 3467–3472 (2024). doi: 10.1134/S1062359024701826.

Abstract

The effect of threefold (1 hour a day with an interval of three days) and multiple (8 weeks, 5 days a week, 2 hours a day) pulsed-modulated electromagnetic radiation on the orienting-exploratory response was studied on males and females of Wistar rats by the “open field” method. The animals were irradiated in an anechoic chamber in the zone of the formed wave of the microwave source (a block of 10 generators carrying frequencies from 1 to 4 GHz, the total pulse power density 300 μW/cm^2, pulse duration 25 ms, sweep cycles 1–6 Hz, cycle turn-around time 30 s). It was shown that the exposure to such parameters could cause a stress reaction and an impairment of long-term memory in some irradiated animals, female rats being more sensitive to the action of microwaves than males. This reaction was transient and the animal behavior was restored 1.5–2 months after cessation of irradiation. However, there is a danger that in the case of constant exposure to microwaves, which, for example, occurs in mobile users, such behavior deviations may be fixed and clinically significant.

https://link.springer.com/article/10.1134/S1062359024701826

--

Experimental Study of Stress Reactions Induced by Low-Intensity Microwave Radiation

Izmestieva OS, Kolganova OI, Chibisova OF *et al.* Experimental Study of Stress Reactions Induced by Low-Intensity Microwave Radiation. *Biol Bull Russ Acad Sci* **51**, 3457–3466 (2024). doi: 10.1134/S1062359024701802.

Abstract

A study was made of the severity and dynamics of changes in indicators reflecting the state of the blood system (the number of myelokaryocytes in the femur, the level of CFU-S in the exo and endotest, hemolytic resistance of erythrocytes, as well as the mass and cellularity of the thymus) in mice exposed to monochromatic pulsed electromagnetic fields of the frequency range 2.27–2.78 GHz with an average power flux density of 60 μW/cm^2 at doses from 0.086 to 0.86 J/g. It is shown that, under repeated exposure to electromagnetic fields, the bioeffect cumulates, leading to the development of a stress-like adaptive reaction in animals.

https://link.springer.com/article/10.1134/S1062359024701802

--

Effects of 700MHz radiofrequency radiation (5G lower band) on the reproductive parameters of female Wistar rats

Jha N, Sarsaiya P, Tomar AK, Pardhiya S, Nirala JP, Chaturvedi PK, Gupta S, Rajamani P. Effects of 700MHz radiofrequency radiation (5G lower band) on the reproductive parameters of female Wistar rats. Reprod Toxicol. 2025 Apr 10:108910. doi: 10.1016/j.reprotox.2025.108910.

Abstract

The advent of 5G technology has raised concerns about its potential biological effects, particularly reproductive health being one key area of focus. This study investigated the impact of 700MHz, a lower

5G frequency band, on the reproductive health of female Wistar rats. The experiment analyzed the effects of short-term and long-term exposure to 700MHz mobile radiation on female Wistar rats. Rats were divided into three groups (control, sham-exposed, and exposed), with sample sizes of n=6 for short-term and n=8 for long-term exposure. For short-term exposure, rats were subjected to 6 hrs of radiation daily for 10 days, while for long-term exposure, rats exposed 4 hrs daily for 60 days. Physiological parameters, including estrous cyclicity, were monitored, and histopathological and biochemical analyses were conducted on harvested ovaries. Comet assay was performed to assess DNA damage. The results indicated no changes in estrous cycles or comet assay parameters in either exposure group. Serological hormone levels, including estradiol and progesterone, remained within normal ranges, but a slight yet significant increase in testosterone levels was observed in exposed groups. Oxidative stress markers revealed elevated malondialdehyde (MDA) levels and significant decreases in superoxide dismutase (SOD), total sulfhydryl content, and ferric reducing antioxidant power (FRAP) in exposed ovaries. Histopathological analysis showed no significant changes in ovarian morphology in short-term exposure but revealed alterations, including cystic follicles and abnormal vasculature, in long-term exposure. These findings suggest that 700MHz radiation may induce oxidative stress and tissue changes in ovarian samples over prolonged exposure

Highlights

• The length of estrous cycle remained unaffected in both short-term and long-term exposure to 700 MHz.
• Short-term exposure showed no major histoarchitectural changes. However, in long term exposure the exposed group showed abnormal structures when compared to control and sham group.
• For both short-term and long-term exposure significant changes in the testosterone level of the exposed group observed. However, the change in level remained within the normal range.
• The comet assay results indicated no DNA damage in both short-term and long-term exposure.

https://pubmed.ncbi.nlm.nih.gov/40220970/

--

Possible effects of RF EMR on contextual fear conditioning, hippocampal perivascular space, apoptosis and adrenal gland microarchitecture in rats

Narayanan SN, Kumar RS, Kumar N, Prabhakar P, Nayak SB, Bhat PG. Possible effects of radiofrequency electromagnetic radiation on contextual fear conditioning, hippocampal perivascular space, apoptosis and adrenal gland microarchitecture in rats. Behav Brain Res. 2025 Mar 12;481:115424. doi: 10.1016/j.bbr.2025.115424.

Abstract

Whilst the world sees the tremendous growth of mobile phone technology, radiofrequency electromagnetic radiation (RF-EMR) induced possible health effects have emerged as a topic of recent day debate. The current study is designed to test the hypothesis that chronic 900 MHz radiation exposure would potentially dysregulate the stress response system (HPA axis) in vivo, via, its non-thermal mechanisms, leading to alterations in the microarchitecture of the adrenal gland, vulnerable brain regions such as the hippocampus which may results in altered behaviours in rats. Male albino

Wistar rats aged four weeks, weighing 50-60 g were subjected to 900 MHz radiation from a mobile phone for four weeks at a rate of one hour per day. On the 29th day, animals from the control, sham exposed and RF-EMR exposed groups were tested for contextual fear conditioning. They were later euthanized to study hippocampal and adrenal gland cytoarchitecture. Bright and dark compartment transitions in the avoidance box were considerably elevated in the RF-EMR exposed group and they exhibited a significant decrease in the latency to enter the dark compartment during the contextual fear conditioning test. Apoptosis was apparent in the CA3 region and perivascular space was significantly increased in the hippocampus of the radiation-exposed group. In addition to lymphocytic infiltrates, congested sinusoids, apoptotic-like changes were evident in the zona fasciculata of the adrenal gland. However, the cytoarchitecture of the adrenal medulla was comparable in all three groups. Chronic RF-EMR exposure caused changes in contextual fear conditioning, enlargement of hippocampal perivascular space, apparent CA3 apoptosis, and apoptotic-like changes in the zona fasciculata of the adrenal gland in rats.

Highlights

Chronic 900 MHz radiation-induced altered contextual fear memory.

Exposure to 900 MHz radiation resulted in increasing the hippocampal perivascular space.

Altered memory is attributed to hippocampal apoptosis induced by 900 MHz radiation.

900 MHz radiation triggered apoptosis and lymphocytic infiltration in the adrenal gland.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0166432825000105?via%3Dihub

--

Impact of a Terahertz electromagnetic field on the ion permeation of potassium and sodium channels

Song Z, Xue L, Ouyang Q, *et al.* Impact of a Terahertz electromagnetic field on the ion permeation of potassium and sodium channels. *Commun Chem* 8, 101 (2025). doi: 10.1038/s42004-025-01503-4.

Abstract

Ion channels are essential for various physiological processes, and their defects are associated with many diseases. Previous research has revealed that a Terahertz electromagnetic field can alter the channel conductance by affecting the motion of chemical groups of ion channels, and hence regulate the electric signals of neurons. In this study, we conducted molecular dynamics simulations to systematically investigate the effects of terahertz electromagnetic fields on the ion permeation of voltage-gated potassium and sodium channels, particularly focusing on the bound ions in the selectivity filters that have not been extensively studied previously. Our results identified multiple new characteristic frequencies and showed that 1.4, 2.2, or 2.9 THz field increases the ion permeability of $K_v1.2$, and 2.5 or 48.6 THz field increases the ion permeability of $Na_v1.5$. Such effects are specific to the frequencies and directions of the electric field, which are determined by the intrinsic oscillation motions of the permeating ions in the selectivity filter or certain chemical groups of the ion channels. The amplitude of the THz field positively correlates with the change in ion permeation. This study

demonstrates that THz fields can specifically regulate ion channel conductances by multiple mechanisms, which may carry great potential in biomedical applications.

Open access paper: https://www.nature.com/articles/s42004-025-01503-4

--

Residential exposure to magnetic field due to high-voltage power lines and childhood leukemia risk in mainland France - GEOCAP case-control study

Mancini M, Hémon D, Faure L, Clavel J, Goujon S. Residential exposure to magnetic field due to high-voltage power lines and childhood leukemia risk in mainland France - GEOCAP case-control study, 2002-2010. Environ Res. 2025 Apr 16:121638. doi: 10.1016/j.envres.2025.121638.

Abstract

Background: Environmental exposure to extremely low frequency magnetic fields (ELF-MF) is suspected of being a risk factor of childhood acute leukemia (AL) and classified as possible carcinogen. Results of recent epidemiological studies remain however heterogeneous.

Objective: The present study aimed to evaluate AL risk in children exposed to ELF-MF by living close to high-voltage overhead power lines (HVOL) in France.

Methods: We included 4,117 AL cases under the age of 15 diagnosed in 2002-2010 and 44,838 contemporary controls representative of the French pediatric population, drawn from the national registry-based GEOCAP study. The distance between the geocoded address of residence and the closest 63-400kV HVOL, and the closest 225-400kV HVOL, were evaluated. ELF-MF exposure was also calculated at the geocoded addresses considering the characteristics of the neighboring HVOLs. Logistic regression models adjusted for age were used to estimate odds-ratios (OR) and 95% confidence intervals. Sensitivity analyses were carried out to account for geocoding error and potential confounders.

Results: 0.7% of the controls lived within 50 meters of HVOL and 0.3% were exposed to more than 0.3 µT. Living within 50 m of HVOL was associated with an increased risk of AL for children under 5 (OR=1.6 (1.0-2.7)), an association more marked when restricting to the high-quality geocoded addresses (OR=3.2 (1.3-7.9)). ELF-MF was not associated with AL risk (≥ 0.3 µT, OR=0.6 (0.3-1.3)). The results remained stable in all the sensitivity analyses.

Conclusion: Our study brings new evidence that ELF-MF are probably not associated with AL risk, and cannot explain an association with distance to HVOL.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0013935125008898?via%3Dihub

--

Use of Electrical Household Appliances and Risk of All Types of Tumours: A Case-Control Study

Noori S, Aleem A, Sultan IN, Tareen AK, Ullah H, Khan MW. Use of Electrical Household Appliances and Risk of All Types of Tumours: A Case-Control Study. Med Sci (Basel). 2025 Apr 1;13(2):36. doi: 10.3390/medsci13020036.

Abstract

Introduction: The use of electrical appliances using extremely low frequency (ELF) electromagnetic fields (EMF) has increased in the past few years. These ELF MF are reported to be linked to several adverse health effects. However, only a couple of studies have been conducted on the association between risk of tumours and use of electronic devices using low frequency (LF) EMF.

Methods: We studied the use of common household electrical appliances and suspected risk of tumours in a multi-hospital-based case-control study. In total, 316 patients were included in the final analysis.

Results: The study results showed a below unity risk for most of the devices. A slight increased risk of tumour was observed for computer screen use OR: 1.13 (95% CI: 0.43-3.02) and use of microwave oven OR: 1.21 (95% CI: 0.36-4.04). We also had chance to investigate ELF MFs exposure association with tumour. Where we observed elevated odd ratios in individuals living near electricity transformer stations, with a statistically significant risk OR: 2.16 (95% CI: 1.30-3.59). However, the risk was below unity (OR: 0.98) in individuals residing close to powerlines.

Conclusion: The current study serves as a pilot study of primary data and will be helpful in future epidemiological research studies on the topic in the region.

Open access paper: https://www.mdpi.com/2076-3271/13/2/36

--

Magnetoreception and the ruling hypothesis

Vortman Y, Fitak R, Natan E. Magnetoreception and the ruling hypothesis. J Exp Biol. 2025 Apr 1;228(7):jeb250252. doi: 10.1242/jeb.250252.

Abstract

Whereas science is written by humans and cannot escape emotions intervening with scientific thought, the scientific community should be on guard against unnoticeably adopting a favorite hypothesis. When adopting a favorite hypothesis, scientists tend to review their work in favor of this hypothesis and reject contradictory data. In 1890, Thomas Chrowder Chamberlin first described this phenomenon as when 'the search for facts, and their interpretation are dominated by affection for the favored theory until it appears to its advocate to have been overwhelmingly established'. The favorite hypothesis can then quickly transition into a ruling hypothesis, leading to an unconscious bias in favor of supporting evidence and neglect of contradictory observations. This is especially problematic when a scientific field adopts a favorite hypothesis. In this Commentary, we suggest that the field of animal magnetoreception - in particular mechanisms based on radical-pair chemistry and cryptochrome proteins - may be under the reign of a ruling hypothesis. We argue that repeatedly, conclusions are unfounded or otherwise not consistent with the results presented. We use the case of magnetoreception - the only sense that remains without a clearly described receptor - to raise general awareness of the phenomenon of a ruling

hypothesis in the scientific community. We emphasize the distinction between the scientist and the scientific community suffering from a hypothesis regime, and further highlight suggestions to mitigate the risk of working under a ruling hypothesis.

https://pubmed.ncbi.nlm.nih.gov/40207401/

--

Effect of electromagnetic field radiation on transcriptomic profile and DNA methylation level in pig conceptuses during the peri-implantation period

Franczak A, Zmijewska A, Drzewiecka EM, Kozlowska W, Wydorski P, Paukszto L, Brevini TL. Effect of electromagnetic field radiation on transcriptomic profile and DNA methylation level in pig conceptuses during the peri-implantation period. Sci Rep. 2025 Apr 23;15(1):14025. doi: 10.1038/s41598-025-98918-9.

Abstract

Extremely low-frequency electromagnetic field (ELF-EMF) radiation alters the steroidogenic activity of porcine conceptuses during the peri-implantation period. This study investigated whether short exposure (2 h) to ELF-EMF radiation may induce changes in the transcriptomic profile of conceptuses and their DNA methylation levels, with a focus on the promoter regions of differentially expressed genes of interest. Porcine conceptuses were collected on days 15-16 of pregnancy, preincubated, and then exposed in vitro to ELF-EMF at a frequency of 50 Hz. ELF-EMF treatment affected the expression of 21 protein-coding transcripts, including solute carrier family 34 member 1 (SLC34A1), hydroxysteroid 17-beta dehydrogenase 2 (HSD17B2), apolipoprotein M (APOM), regucalcin (RGN), heat shock protein family A (Hsp70) member 6 (HSPA6), fibrinogen beta chain, and fibrinogen gamma chain (FGG). There were evaluated 21 GO annotations for biological process terms, nine GO annotations for cellular component terms and two KEGG pathways. The ELF-EMF exposure increased approximately 16 times genomic DNA methylation in conceptuses. ELF-EMF-induced changes in methylation were observed in the promoter regions of APOM, HSD17B2, FGG, and SLC34A1. The results determined 116 predicted single nucleotide variant substitutions within RNA editing sites. In conclusion, ELF-EMF radiation appears to interact with DNA methylation levels and to affect the expression of genes involved mainly in cellular homeostasis, conceptuses development, and attachment, suggesting the need for further investigation to elucidate better the impact of ELF-EMF exposure on the physiology of early conceptuses.

Open access paper: https://www.nature.com/articles/s41598-025-98918-9

--

Potential influence of geomagnetic activity on blood pressure statistical fluctuations at mid-magnetic latitudes

He P, Li C, Xu M, Guo R, Degeling AW, Pitkänen T, Bu Y, Zheng X, Zhang Y, Jia X, Tian A, Han C, Wang S, Chen T, Fang J, Sun S, Liu W, Cao J, Quan K, Cong Z, Ma D, Zong Q, Fu S, Yao S, Zhang H, Shi Q. Potential influence of geomagnetic activity on blood pressure statistical fluctuations at mid-magnetic latitudes. Commun Med (Lond). 2025 Apr 28;5(1):143. doi: 10.1038/s43856-025-00822-w.

Abstract

Background: Solar activity and the consequent geomagnetic activity (GMA) profoundly influence human biological rhythms and cardiovascular system functions. Although the response of blood pressure (BP) to GMA has attracted considerable attention, it is unclear whether the GMA can have an influence alone and how it occurs.

Methods: In this six-year time series analysis, we collated over 500,000 BP measurements from two representative cities (Qingdao and Weihai) at mid-magnetic latitudes in China. Using various statistical methods, we analyzed the correlation between BP and the GMA (represented by Ap index) and their quasi-periodic fluctuations. Additionally, we conducted a comparative analysis of the influence of other environmental factors (air temperature and PM2.5) on BP.

Results: The statistical BP level fluctuations correlate with the GMA. Both BP and the GMA index exhibit similar annual bimodal patterns and multiple periodicities, including 12-month and 6-month cycles, and an intermittent 3-month cycle. In contrast, other known environmental factors influencing BP such as air temperature and PM2.5 do not exhibit similar periodicities, particularly they lack 3-month cycles. In years with higher GMA levels, the BP shows stronger correlations with the Ap index and responds on a shorter timescale. Additionally, BP in females appears to be more strongly correlated with GMA.

Conclusions: Our findings highlight potential risks to individuals with hypertension with elevated GMA levels, deepen our understanding of GMA's role in human health, and offer insights for healthcare policymakers on the clinical significance of the geomagnetic environment.

Plain language summary

Human blood pressure fluctuates constantly and is influenced by various factors such as hormone levels in the body and environmental conditions. Geomagnetic activity is change in the Earth’s magnetic field as a consequence of the sun’ s activity, such as the impact of solar winds. We used six years of data to investigate whether changes in geomagnetic activity have an impact on blood pressure. Our results show a relationship between geomagnetic activity and blood pressure, particularly during periods of high geomagnetic activity. Understanding this relationship could enhance our knowledge of the geomagnetic environment’s clinical significance and potentially contribute to better hypertension risk assessment and management.

Open access paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC12038039/

--

Earth's magnetic field and its relationship to the origin of life, evolution and planetary habitability

Tarduno JA, Zhou T, Huang W, Jodder J. Earth's magnetic field and its relationship to the origin of life, evolution and planetary habitability. Natl Sci Rev. 2025 Mar 3;12(5):nwaf082. doi: 10.1093/nsr/nwaf082.

Abstract

Earth's magnetic field history can provide insight into why life was able to originate and evolve on our planet, and how habitability has been maintained. The magnetism of minute magnetic inclusions in zircons indicates that the geomagnetic field is at least 4.2 billion years old, corresponding with genetic estimates for the age of the last universal common ancestor. The early establishment of the field would have provided shielding from solar and cosmic radiation, fostering environments for life to develop. The field was also likely important for preserving Earth's water, essential for life as we know it. Between 3.9 and ca. 3.4 billion years ago, zircon magnetism suggests latitudinal stasis of different ancestral terrains, and stagnant lid tectonics. These data also indicate that the solid Earth was stable with respect to the spin axis, consistent with the absence of plate tectonic driving forces. Moreover, these data point to the existence of low-latitude continental nuclei with equable climate locales that could have supported early life. Near the end of the Precambrian (0.591 to 0.565 billion years ago), the dynamo nearly collapsed, but growth of the inner core during earliest Cambrian times renewed the magnetic field and shielding, helping to prevent drying of the planet. Before this renewal, the ultra-weak magnetic shielding may have had an unexpected effect on evolution. The extremely weak field could have allowed enhanced hydrogen escape to space, leading to increased oxygenation of the atmosphere and oceans. In this way, Earth's magnetic field may have assisted the radiation of the macroscopic and mobile animals of the Ediacara fauna. Whether the Ediacara fauna are genetically related to modern life is a matter of debate, but if so, magnetospheric control on atmospheric composition may have led to an acceleration in evolution that ultimately resulted in the emergence of intelligent life.

Open access paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC11980988/

--

The effects of radiofrequency radiation on male reproductive health and potential mechanisms (Review)

Bektas H, Dasdag S. The effects of radiofrequency radiation on male reproductive health and potential mechanisms. Electromagn Biol Med. 2025 Mar 19:1-26. doi: 10.1080/15368378.2025.2480664.

Abstract

Recent studies have demonstrated that radiofrequency (RF) radiation emanating from devices such as mobile phones and Wi-Fi may have adverse effects on male reproductive health. This radiation can elevate testicular temperature, potentially compromising sperm quality and DNA integrity, and influence the specific absorption rate (SAR) across different body regions, leading to detrimental reproductive outcomes. Furthermore, exposure to RF radiation has been linked to conditions that could affect male reproductive function, such as oxidative stress, alterations in ion transitions across cell membranes, and inflammation. The article reviews research conducted on both humans and animal models regarding the effects of electromagnetic radiation on sperm quality, DNA damage, oxidative stress, hormone levels, and testicular function, suggesting that exposure to electromagnetic radiation could have harmful implications for male reproductive health. However, further research is necessary to fully understand the mechanisms and implications of non-ionizing electromagnetic radiation on male infertility.

Plain-language Summary

In this review, the effects of RF on male reproduction and the mechanisms related to them were examined and the results were discussed. “The effects of RF on sperm, reproductive hormones and testis, occupational exposure and male reproductive health, mechanisms of RF effects on reproductive system” were discussed here. In conclusion, further research is necessary to fully understand the mechanisms and implications of non-ionizing electromagnetic radiation on male infertility.

Excerpts

A total of 90 articles were evaluated in this review.

The effects of RF radiation on sperm

In conclusion, despite an incomplete understanding of the precise mechanisms underlying the impact of non-thermal RF radiation on sperm and testicular function, research across animal and human studies consistently suggests a connection between increased levels of ROS and/or DNA damage and negative impacts on fertility factors. This indicates that the negative effects of RF radiation on sperm parameters may primarily be linked to the induction of ROS. Furthermore, the variability in study outcomes underscores the complexity of RF radiation’s effects, which may be influenced by factors such as exposure duration, frequency, and individual susceptibility. Therefore, limiting mobile phone use may be advisable to mitigate the incidence of male infertility. However, further research is warranted to explore the long-term effects of mobile phone radiation on male fertility and the generalizability of these findings (). Future studies should aim to clarify the specific biological pathways affected by RF radiation and standardize experimental protocols to resolve inconsistencies in the literature.

The effects of RF radiation on DNA

In conclusion, these studies highlight the harmful effects of RF radiation on male reproductive health, including DNA damage, decreased sperm mobility and vitality, increased DNA fragmentation, and oxidative stress. The variability in the findings, driven by differences in study design, exposure parameters, and biological models, suggests that the biological impact of RF radiation may be multifactorial, involving complex interactions between ROS production, DNA repair mechanisms, and cellular apoptosis pathways. It is advisable to limit RF exposure by keeping mobile phones away from the pelvic region and minimizing the use of RF-emitting electronic devices. However, given the inconsistent outcomes across various studies, future research should focus on standardizing experimental conditions and exploring the dose-response relationship to better understand the threshold levels at which RF radiation becomes detrimental to reproductive health. Further research is necessary to fully understand the long-term implications of RF radiation on human reproductive health.

The effects of RF radiation on reproductive hormones and testis

In conclusion, the research discussed in this section indicates that RF radiation exposure could negatively affect male reproductive health, leading to changes in hormone levels, testicular morphology, and sperm count and viability (). Prolonged exposure to wireless internet and mobile phones has been associated with alterations such as increased abnormalities in sperm heads and reduced weight of reproductive organs. Additionally, RF radiation can impact testicular parenchyma in rats, resulting in irregularly shaped seminiferous tubules with epithelial cell abnormalities. The complexity of these findings suggests that RF radiation may induce a cascade of biological responses, including oxidative stress, hormonal imbalances, and apoptotic processes, which collectively contribute to testicular

damage and compromised reproductive health. The observed effects appear to be dose-dependent, with longer durations and higher intensities of RF exposure correlating with more severe reproductive outcomes. These findings emphasize the importance of avoiding long-term exposure to RF radiation emissions, particularly from wireless technologies, to prevent potential harm to male reproductive health. Future research should focus on elucidating the precise mechanisms by which RF radiation affects the reproductive system and identifying potential protective strategies to mitigate these effects.

Occupational exposure and male reproductive health

In conclusion, the diverse findings from these studies shed light on the complex relationship between occupational exposure, particularly to radiofrequency radiation, and male reproductive health outcomes (). The variability in study outcomes highlights the challenges in establishing a clear causal link, suggesting that individual susceptibility, exposure duration, and the specific characteristics of RF equipment used in different occupational settings may play significant roles. Further research is warranted to elucidate the underlying mechanisms and to inform occupational health guidelines and practices. Future studies should focus on standardizing exposure assessment methods and consider long-term monitoring of workers in high-risk occupations to better understand the cumulative effects of RF radiation. Additionally, exploring potential protective strategies, such as shielding technologies and exposure limits, could be critical in mitigating the reproductive risks associated with occupational RF exposure.

Conclusion

This study comprehensively examined the mechanisms by which RF radiation may impact male reproductive health, focusing on both thermal and non-thermal pathways. The findings indicate that RF exposure, particularly through increased oxidative stress, ionic imbalances, and inflammation, can disrupt spermatogenesis and impair sperm quality. While thermal effects highlight the role of testicular temperature elevation, non-thermal mechanisms such as ROS generation and ionic dysregulation further emphasize the multifaceted nature of RF-induced reproductive toxicity. Despite the robust body of evidence, inconsistencies across studies warrant standardized research protocols to resolve methodological discrepancies. Future investigations should prioritize long-term exposure assessments and explore protective strategies to mitigate the risks associated with RF radiation. These efforts are critical for informing public health guidelines and ensuring reproductive health safety in the era of pervasive wireless communication technologies. Findings underline the importance of revisiting current regulatory standards to better protect reproductive health.

https://pubmed.ncbi.nlm.nih.gov/40108785/

--

Use of Mobile and Cordless Phones and the Association with Prostate Cancer

Hardell L, Carlberg M. Use of Mobile and Cordless Phones and The Association with Prostate Cancer. Fortune Journal of Health Sciences, 8 (2025): 267-273. doi: 10.26502/fjhs.273

Abstract

Exposure to radiofrequency (RF) radiation in the frequency range 30 kHz–300 GHz was in 2011 evaluated by the International Agency for Research on Cancer (IARC) at the World Health Organization

(WHO) to be a 'possible' human carcinogen, Group 2B. This was based on epidemiological results on increased risk for glioma and acoustic neuroma. Results on other cancer types are sparse. An increased incidence in male rats of proliferative lesions in the prostate gland induced by RF radiation was found in the US NTP study. Thus, it was pertinent to study an association with prostate cancer in human studies. We analyzed data in two of our previous studies, one on brain tumors (only deceased subjects; those who died from prostate cancer were defined as cases) and another on prostate cancer (living subjects) that included similar questions on use of mobile phones or cordless phones. The pooled analysis gave for mobile phone use OR = 1.8, 95 % CI = 1.01-3.1, increasing in the >10 year latency group to OR = 2.8, 95 % CI = 1.5-5.3. Also, use of the cordless phone gave increased risk, although not statistically significant. Dose-response analysis gave highest risk for >2,000 h use of the mobile phone with OR = 2.4, 95 % CI = 1.2-5.1. The cordless phone yielded highest risk in the group 1001-2000 h with OR = 2.3, 95 % CI = 1.01-5.4. Lower OR was seen for use > 2,000 h but based on low numbers. Higher risk was seen in cases with more aggressive cancer based on Gleason score, PSA, and high risk profile, and among subjects with heredity for prostate cancer.

Open access paper: https://www.fortunejournals.com/articles/use-of-mobile-and-cordless-phones-and-the-association-with-prostate-cancer.html

--

Disease burden, risk factors, and trends of primary central nervous system (CNS) cancer: A global study of registries data

My note: Because it can take up to three decades for a solid tumor to be diagnosed and reported in tumor registries, it is important to model the latency when examining tumor incidence data. Nonetheless, it is tricky trying to interpret ecological studies. Moreover, trends over time in tumor incidence can be difficult to interpret due to changes in screening, diagnostic and reporting procedures.

Huang J, Chan SC, Lok V, Zhang L, Lin X, Lucero-Prisno DE, Xu W, Zheng ZJ, Elcarte E, Withers M, Wong MCS; NCD Global Health Research Group; Association of Pacific Rim Universities (APRU). Disease burden, risk factors, and trends of primary central nervous system (CNS) cancer: A global study of registries data. Neuro Oncol. 2023 May 4;25(5):995-1005. doi: 10.1093/neuonc/noac213.

Key Points

- Brain cancer burden was higher in more developed countries and male population.
- Brain cancer was related to HDI, GDP, brain injuries, carcinogens, and phone use.
- There was an increasing trend of brain cancer in the younger male population.

Abstract

Background: This study aimed to evaluate the global incidence, mortality, associated risk factors, and temporal trends of central nervous system (CNS) cancer by sex, age, and country.

Methods: We extracted incidence and mortality of CNS cancer from the GLOBOCAN (2020), Cancer Incidence in Five Continents series I-X, WHO mortality database, the Nordic Cancer Registries, and the Surveillance, Epidemiology, and End Results Program. We searched the Global Health data exchanges for

the prevalence of its associated risk factors. We tested the trends by Average Annual Percentage Change (AAPC) from Joinpoint regression analysis with 95% confidence intervals in different age groups.

Results: The age-standardized rates (ASRs) of CNS cancer incidence and mortality were 3.5 and 2.8 per 100,000 globally. Southern Europe (ASR = 6.0) and Western Asia (ASR = 4.2) had the highest incidence and mortality, respectively. The incidence was associated with Human Development Index [HDI], Gross Domestics Products per capita [GDP], prevalence of traumatic brain injuries, occupational carcinogens exposure, and mobile phone use at the country level. There was an overall stable and mixed trend in the CNS cancer burden. However, increasing incidence was observed in younger male population from five countries, with Slovakia (AAPC = 5.40; 95% CI 1.88, 9.04; P = .007) reporting the largest increase.

Conclusions: While the overall global trends of cancer have been largely stable, significant increasing trends were found in the younger male population. The presence of some higher-HDI countries with increasing mortality suggested an ample scope for further research and exploration of the reasons behind these epidemiological trends.

Excerpts

"There are inconsistent results reporting by different settings of studies regarding the associations between mobile phone use and CNS cancer. Although some studies show mobile phone use was not associated with an increased risk of brain cancer, 45–47 they may have suffered from poor exposure assessment that likely contributed to exposure mis-classification. 48 A meta-analysis of 11 studies found a significant positive association between long-term ipsilateral mobile phone use and the risk of glioma (OR = 1.46, 95% CI 1.12–1.92). 49 Another meta-analysis showed a similar significant 1.33 times increase in risk. 11 A more recent meta-analysis of 46 studies found increased CNS cancer incidence with cumulative call time of 1000 or more hours. 50"

"Higher CNS cancer incidence was associated with a higher-HDI (β_{male} = 0.87, CI 0.66–1.08, P < .001; β_{female} = 0.59, CI 0.45–0.73, P < .001), GDP per capita (β_{male} = 0.49, CI 0.32–0.66, P < .001; β_{female} = 0.30, CI 0.18–0.42, P < .001), and higher prevalence of traumatic brain injuries (β_{male} = 2.92, CI 2.47–3.37, P < .001; β_{female} = 2.79, CI 2.35–3.24, P < .001), occupational carcinogens (β_{male} = 1.39, CI 0.13–2.66, P = .031; β_{female} = 0.78, CI 0.06–1.49, P = .034), and **mobile [phone] use (β_{male} = 2.07, CI 0.83–3.30, P = .001; β_{female} = 1.43, CI 0.61–2.24, P = .001)** but not with exposure to unsafe water (P > .05 for both sexes; Figure 2)."

"Higher CNS cancer mortality was associated with a higher-HDI (β_{male} = 0.58, CI 0.41–0.74, P < .001; β_{female} = 0.37, CI 0.26–0.47, P < .001), GDP per capita (β_{male} = 0.29, CI 0.16–0.42, P < .001; β_{female} = 0.18, CI 0.09–0.26, P < .001), and higher prevalence of traumatic brain injuries (β_{male} = 1.96, CI 1.60–2.32, P < .001; β_{female} = 1.64, CI 1.28–2.00, P < .001), occupational carcinogens (β_{male} = 1.06, CI 0.12–2.00, P = .027; β_{female} = 0.63, CI 0.12–1.14, P = .016), **and mobile [phone] use (β_{male} = 1.34, CI 0.42–2.25, P = .005; β_{female} = 0.81, CI 0.22–1.39, P = .007)** but not with the exposure to unsafe water (P > .05 for both sexes; Figure 3)."

"The largest incidence and mortality of CNS cancer were found in populations with very high HDI. The high incidence could be attributable to the ample resources in early detection with advanced diagnostic techniques and regular health check-ups while the high mortality could be ascribed to higher exposure to its related risk factors, such as traumatic brain injuries, occupational carcinogens exposure, and mobile phone use. While the overall global trends of cancer have been largely stable, significant

increasing trends were found in the younger male population. The presence of some higher-HDI countries with increasing mortality suggested an ample scope for further research and exploration of the reasons behind these epidemiological trends."

Open access paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC10158137/

--

Could electrohypersensitivity be a specific form of high sensory processing sensitivity?

Bordarie J, Ledent M, Dieudonné M, Choisay F, De Clercq E. Could electrohypersensitivity be a specific form of high sensory processing sensitivity? Front Public Health. 2025 Feb 28;13:1550427. doi: 10.3389/fpubh.2025.1550427.

Abstract

Introduction: Electrohypersensitivity (EHS) refers to a syndrome in which individuals claim to suffer from a variety of symptoms that they attribute to electromagnetic fields. The characteristics of this specific hypersensitivity, particularly in terms of symptoms, are similar to those associated with high sensory processing sensitivity (HSPS). This article raises the question of the superposition of these two types of sensitivity and investigates the existence of a link between the two.

Methods: Participants (n = 100) completed a questionnaire measuring EHS and HSPS, as well as absorption, risk perception and avoidance strategies related to electromagnetic fields, and anxiety and depressive disorders.

Results: They showed an overrepresentation of highly sensitive people within the electrohypersensitive group. Furthermore, the results showed differences in terms of anxiety-depressive symptomatology and cognitive strategies (risk perception and avoidance strategies).

Discussion: The article discusses these results in the light of the literature and suggests avenues for future research and ways to help highly sensitive people, whether wor not this condition is considered to be caused by electromagnetic radiation.

Open access paper: https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1550427/full

---

Self-diagnosing electromagnetic hypersensitivity—A case study

Ashton D. Self-diagnosing electromagnetic hypersensitivity—A case study. Front Public Health 2025; 13: 1535513

No abstract

Open access paper: https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1535513/full

--

A Novel Method for Achieving Precision and Reproducibility in a 1.8 GHz Radiofrequency Exposure System That Modulates Intracellular ROS as a Function of Signal Amplitude in Human Cell Cultures

Dahon C, Aguida B, Lebon Y, Le Guen P, Dangremont A, Meyer O, Citerne JM, Pooam M, Raad H, Thoradit T, Jourdan N, Bertagna F, Ahmad M. A Novel Method for Achieving Precision and Reproducibility in a 1.8 GHz Radiofrequency Exposure System That Modulates Intracellular ROS as a Function of Signal Amplitude in Human Cell Cultures. Bioengineering (Basel). 2025 Mar 4;12(3):257. doi: 10.3390/bioengineering12030257.

Abstract

Radiofrequency fields in the 1-28 GHz range are ubiquitous in the modern world, giving rise to numerous studies of potential health risks such as cancer, neurological conditions, reproductive risks and electromagnetic hypersensitivity. However, results are inconsistent due to a lack of precision in exposure conditions and vastly differing experimental models, whereas measured RF effects are often indirect and occur over many hours or even days. Here, we present a simplified RF exposure protocol providing a single 1.8 GHz carrier frequency to human HEK293 cell monolayer cultures. A custom-built exposure box and antenna maintained in a fully shielded anechoic chamber emits discrete RF signals which can be precisely characterized and modelled. The chosen amplitudes are non-thermal and fall within the range of modern telecommunication devices. A critical feature of the protocol is that cell cultures are exposed to only a single, short (15 min) RF exposure period, followed by detection of immediate, rapid changes in gene expression. In this way, we show that modulation of genes implicated in oxidative stress and ROS signaling is among the earliest cellular responses to RF exposure. Moreover, these genes respond in complex ways to varying RF signal amplitudes consistent with a hormetic, receptor-driven biological mechanism. We conclude that induction of mild cellular stress and reactive oxygen species (ROS) is a primary response of human cells to RF signals, and that these responses occur at RF signal amplitudes within the range of normal telecommunications devices. We suggest that this method may help provide a guideline for greater reliability and reproducibility of research results between labs, and thereby help resolve existing controversy on underlying mechanisms and outcomes of RF exposure in the general population.

Excerpts

Concluding Remarks and Future Perspectives

We here describe an experimental RF exposure device and protocol that present fully characterized, defined RF signals to human cells in culture. Their effects are consistent with a biological receptor-driven mechanism whereby RF exposure modulates intracellular ROS and ROS signaling pathways. This provides a testable hypothesis for the many and varied effects of RF described in the literature.

These cellular responses occur at RF signal amplitudes that are orders of magnitude below those needed to achieve thermal effects, and lie within the signal range of personal electronic devices and mobile phones. Because this human cell response to RF is not linear as a function of the RF signal amplitude, the relation between RF exposure conditions and a physiological outcome is not readily deducible; indeed a robust gene expression response may occur at one amplitude but be undetected at another signal amplitude, or even undergo the opposite response entirely (e.g., opposite expression of the same gene at different signal amplitudes). It is therefore necessary to assess physiological response to RF signal

exposure at multiple signal amplitudes and wavelengths, and preferably by using a readout assay that is rapid and direct. This may help explain existing confusion and contradictions in the literature, as well as stimulate future studies on the nature of the biological reception mechanisms.

Finally, although RF exposure from cell phones and telecommunications devices has not been proven harmful in any way, there is a definite physiological response in human beings to this signal range. Risk factors may therefore exist for susceptibility to RF exposure, for instance in individuals with reduced tolerance to oxidative stress and/or who are exposed to excessive stressors in their daily life. These additive or synergistic effects may contribute to certain poorly defined syndromes such as electromagnetic hypersensitivity (EHS) that have been linked to RF exposure in rare individuals in the past [29].

Open access paper: https://www.mdpi.com/2306-5354/12/3/257

--

RF-EMF exposure assessment with add-on uplink exposure sensor in different microenvironments in seven European countries

Bladel HV, Stroobandt B, Veludo AF, Deprez K, Röösli M, Tognola G, Parazzini M, Thuróczy G, Polańska K, Politański P, Wiart J, Guxens M, Joseph W. RF-EMF exposure assessment with add-on uplink exposure sensor in different microenvironments in seven European countries. Environment International, Volume 197, 2025, doi: 10.1016/j.envint.2025.109368.

Abstract

Introduction Several devices have been developed to assess exposure to radiofrequency electromagnetic field (RF-EMF). Since the existing solutions to measure the personal exposure induced by emerging 5G New Radio (NR) are expensive, complex, and bulky, a new cost efficient and low-complexity sensor is developed, that aims to measure RF-EMF exposure in different scenarios of data transmission within different areas.

Methods With this novel sensor, activity-based microenvironmental surveys were conducted across seven European countries: Belgium, Hungary, Italy, Poland, Switzerland, the Netherlands, and the United Kingdom. The device is attached to a smartphone to quantify the auto-induced uplink (a-UL) transmission component of the total exposure for a broadband frequency range from 100 MHz to 6000 MHz and is thus denoted as add-on sensor. In-situ measurements were performed for three usage scenarios, namely non-user (i.e., environmental exposure), maximum downlink (max DL), and maximum uplink (max UL) scenarios, in a large city, a secondary city, and three rural villages a priori selected within each country.

Results Power levels were lowest in non-user scenarios (median: −2.64 dBm or 0.54mW), increasing by a factor of 5.00 dB in maximum downlink scenarios and by a factor of 14.15 dB in maximum uplink scenarios. In the maximum uplink scenarios, the highest median a-UL power of 18.68 dBm (= 73.79 mW) was recorded in The Netherlands, while the lowest median a-UL power of 4.77 dBm (= 3 mW) was observed in the UK. The analysis of the measured data showed a prominent trend of a 2.72 dB lower power in the cities compared to the villages. Further comparisons were made based on microenvironment groups, where the lowest a-UL power levels (median: 12.35 dBm) were measured in

outdoor areas, with an increase of 1.78 dB and 1.91 dB in power was measured compared to public transport and public places, respectively.

Conclusion This study compares RF-EMF power levels between different countries, urbanization settings, and usage scenarios, which is important for future epidemiological studies.

Conclusions

An add-on RF-EMF sensor was used in activity-based microenvironmental surveys. The low-cost broadband add-on sensor was used to map RF-EMF exposure in Belgium, Hungary, Italy, Poland, Switzerland, The Netherlands and The UK in different microenvironments for three network usage scenarios, i.e., non-user, maximum downlink and maximum uplink. Median powers were used to determine any underlying general trends. When examining the measured data, the most prominent trend is found with the difference of 2.27 dB between the cities and villages and the different city areas. Lowest powers are obtained during the non-user scenarios and increase 5.00 dB and 14.15 dB for the maximum downlink and maximum uplink scenarios, respectively.

Future work entails the comparison of the measurement results obtained by this add-on sensor with the results of other measurement devices and phone applications, namely QualiPoc and the ExpoM – RF4, that were used simultaneously to verify the discussed trends. Including the GPS data creates the possibility of investigating present base stations on the different routes and locations. Furthermore, the temporal behavior will be investigated as this activity-based measurement campaign will be repeated after two years, to investigate the influence of the 5G deployment.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412025001199

--

The Effect of Proximity Sensor & Grip Sensor Use on Specific Absorption Rate (SAR) in Smartphones

Aydin E. The Effect of Proximity Sensor & Grip Sensor Use on Specific Absorption Rate (SAR) in Smartphones. Gazi University Journal of Science Part A: Engineering and Innovation. 2025, 12(1):292-306, 26.03.2025. doi: 10.54287/gujsa.1616086.

Abstract

Today, with the contribution of the new generation of communication technologies, many smart devices are produced. Almost every electronic device, including smart phones, smart watches, wireless headphones, tablets, emits some form of radiation. While most of this electromagnetic radiation is harmless, some of it can have potential health effects, depending on the frequency of use over long periods of time and in close usage. Specific Absorption Rate is a measure of how much human body tissue absorbs energy when the body is exposed to radiation. This measurement helps determine whether a device is safe for regular use. The SAR value may vary depending on the antenna and schematic design of the smartphone. To support high band requirements for 5G smartphones, more RF antennas required to be added in PCB design. When designing smartphones, designers also need to design proximity-grip sensors that accurately meet the industry's Specific Absorption Rate (SAR) requirements. In this study, the effects of proximity and grip sensors used in smartphones on LTE and 5G NR SAR values are investigated. During these measurements, a combination of Grip and Proximity

Sensors were alternately turned on and off. Although the proximity sensor and grip sensor are not mainly used to optimize SAR values, it is foreseen that they may have indirect effects on SAR. In this context, SAR measurements were made in 3D environment for different frequencies. As a result of this study, it was observed that the grip-proximity sensors used in smartphones significantly reduce the SAR value and transfer less energy to the users in close range use. The effect of using the proximity sensor on the SAR rate was measured to be approximately 8%, while the effect of using the Grip Sensor was observed to be approximately 10%.

Excerpt

In this study, the effects of the use of the grip sensor and the proximity sensor on the SAR values are realized in 4 different scenarios for different communication technologies. Both grip sensors and proximity sensors contribute indirectly to reducing SAR in mobile phones. However, their effect is primarily through optimized power management. Specifically, proximity sensing can actively reduce transmission power when the phone is near the head of the user, directly reducing SAR. Grip sensors, while helpful in managing the behavior of the phone, have a less direct impact on SAR, but they can contribute to power adjustment based on the way the device is being held. As shown in Figure 7, we achieved best case with both sensors are in ON state, provides from 5% to 11% improvement on SAR values with different frequencies. It is important to use these sensors in future smartphones as a standard hardware component.

Open access paper: https://dergipark.org.tr/en/download/article-file/4504630

--

A Decision Support System for Managing Health Symptoms of Living Near Mobile Phone Base Stations

Parsaei H, Faraz M, Mortazavi SMJ (2024). A Decision Support System for Managing Health Symptoms of Living Near Mobile Phone Base Stations. Journal of Biomedical Physics and Engineering, (), -. doi: 10.31661/jbpe.v0i0.2310-1667.

Abstract

Background: The rapid increase in the number of Mobile Phone Base Stations (MPBS) has raised global concerns about the potential adverse health effects of exposure to Radiofrequency Electromagnetic Fields (RF-EMF). The application of machine learning techniques can enable healthcare professionals and policymakers to proactively address concerns surrounding RF-EMF exposure near MPBS.

Objective: The current study aimed to investigate the potential of machine learning models for the prediction of health symptoms associated with RF-EMF exposure in individuals residing near MPBS.

Material and Methods: This analytical study utilized Support Vector Machine (SVM) and Random Forest (RF) algorithms, incorporating 11 predictors related to participants' living conditions. A total of 699 adults participated in the study, and model performance was assessed using sensitivity, specificity, accuracy, and the Area Under Curve (AUC).

Results: The SVM-based model demonstrated strong performance, with accuracies of 85.3%, 82%, 84%,

82.4%, and 65.1% for headache, sleep disturbance, dizziness, vertigo, and fatigue, respectively. The corresponding AUC values were 0.99, 0.98, 0.920, 0.89, and 0.81. Compared to the RF model and a previously developed model, the SVM-based model exhibited higher sensitivity, particularly for fatigue, with sensitivities of 70.0%, 83.4%, 85.3%, 73.0%, and 69.0% for these five health symptoms. Particularly for predicting fatigue, sensitivity and AUC were significantly improved (70% vs. 8% and 11.1% for SVM, Multilayer Perceptron Neural Network (MLPNN), and RF, respectively, and 0.81 vs. 0.62 and 0.64, for SVM, MLPNN, and RF, respectively).

Conclusion: Machine learning methods, specifically SVM, hold promise in effectively managing health symptoms in individuals residing near or planning to settle in the vicinity of MPBS.

Excerpts

"Figure 1 highlights three variables as the most important predictors of health symptoms: the distance from the mobile base station, the age of the participant, and the duration of residence in the area."

"From a broader standpoint, our findings are in line with studies that have reported that while there is an increasing concern regarding the potential negative health consequences of RF-EMF exposures from mobile phone base stations, the health complaints of individuals living near these base stations cannot be fully explained by these concerns alone [31]. No- tably, previous large population-based studies have shown that residents who were concerned about or attributed detrimental biological effects of RF-EMF generated by mobile phone base stations, as well as those living closer to the base station (e.g., <500 m), had more health complaints compared to others [31]. Furthermore, our results support reports showing the presence of sleep disturbances, headaches, dizziness, irritability, concentration difficulties, and hypertension in the majority of people residing near mobile phone base stations [32]. Additionally, the obtained results align with reports indicating a higher risk of developing neuropsychiatric problems in individuals living in the vicinity of mobile phone base stations. Headache, memory changes, dizziness, tremors, depressive symptoms, and sleep disturbance have been reported to be significantly higher in individuals living around mobile phone base stations [33]."

Open access paper: https://jbpe.sums.ac.ir/article_49803.html

--

Brain Disease-Modifying Effects of Radiofrequency as a Non-Contact Neuronal Stimulation Technology (Review)

Sun S, Bok J, Jang Y, Seo H. Brain Disease-Modifying Effects of Radiofrequency as a Non-Contact Neuronal Stimulation Technology. Int J Mol Sci. 2025 Mar 4;26(5):2268. doi: 10.3390/ijms26052268.

Abstract

Non-invasive, non-contact, and painless methods of electrical stimulation to enhance neural function have been widely studied in recent years, particularly in the context of neurodegenerative diseases such as Alzheimer's disease (AD) and related dementias, which cause cognitive decline and other neurological symptoms. Radiofrequency (RF), which is a rate of oscillation in the range of 3 kHz to 300 GHz (3 THz), has been suggested as one potential non-contact neuronal stimulation (NCNS) technique for improving brain function. A new type of electrical stimulation uses a radiofrequency electromagnetic field (RF-

EMF). RF exposure has been shown to modulate neural stimulation and influence various brain activities in in vitro and in vivo models. Recent studies have explored the effects of RF-EMF on human physiology, particularly in areas such as brain activity, cognition, and sleep behavior. In this review, we summarize recent findings about the effects of non-contact stimulations in in vitro studies, in vivo animal models, and human clinical cases.

Conclusions

In conclusion, RF exposure has the potential to affect neural stimulation and influence various brain activities in in vitro and in vivo models. The in vitro/in vivo effects of RF-EMF exposure are summarized in Figure 1. RF-EMF exposure therapy might improve cognitive performance in optimized conditions. Cognitive dysfunction caused by increased reactive oxygen species and oxidative stress may be improved after RF-EMF exposure through cellular mechanisms such as mitochondrial restoration, gene expression regulation, and cytoskeletal trafficking, etc. Recent studies have explored the influence of EMF on human physiology, particularly brain activity, cognition, sleep, behavior, and sensory functions. Research data vary depending on the RF stimulation conditions, highlighting the need for more clinical trials to clarify its potential effects. However, research is still very limited, and conflicting results can arise depending on the exposure conditions and individual variations. Moreover, although extensive research has been conducted to assess the effects of RF exposure, current data remain insufficient to understand its biological impact. Therefore, careful consideration is needed before clinically applying RF exposure. Additionally, further investigation into the effects of RF and their mechanisms is essential, as many researchers strive to establish reliable safety and efficacy data.

Open access paper: https://www.mdpi.com/1422-0067/26/5/2268

--

Repeated Head Exposures to a 5G-3.5 GHz Signal Do Not Alter Behavior but Modify Intracortical Gene Expression in Adult Male Mice

Lameth J, Royer J, Martin A, Marie C, Arnaud-Cormos D, Lévêque P, Poirier R, Edeline JM, Mallat M. Repeated Head Exposures to a 5G-3.5 GHz Signal Do Not Alter Behavior but Modify Intracortical Gene Expression in Adult Male Mice. Int J Mol Sci. 2025 Mar 10;26(6):2459. doi: 10.3390/ijms26062459. PMID: 40141104; PMCID: PMC11941837.

Abstract

The fifth generation (5G) of mobile communications promotes human exposure to electromagnetic fields exploiting the 3.5 GHz frequency band. We analyzed behaviors, cognitive functions, and gene expression in mice submitted to asymmetrical head exposure to a 5G-modulated 3.5 GHz signal. The exposures were applied for 1 h daily, 5 days per week over a six-week period, at a specific absorption rate (SAR) averaging 0.19 W/kg over the brain. Locomotor activities in an open field, object location, and object recognition memories were assessed repeatedly after four weeks of exposure and did not reveal any significant effect on the locomotion/exploration, anxiety level, or memory processes. mRNA profiling was performed at the end of the exposure period in two symmetrical areas of the right and left cerebral cortex, in which the SAR values were 0.43 and 0.14 W/kg, respectively. We found significant changes in the expression of less than 1% of the expressed genes, with over-representations of genes related to glutamatergic synapses. The right cortical area differed from the left one by an over-

representation of responsive genes encoded by the mitochondrial genome. Our data show that repeated head exposures to a 5G-3.5 GHz signal can trigger mild transcriptome alterations without changes in memory capacities or emotional state.

Exposure System and Exposure Protocol to 5G-3.5 GHz Signals

Head-only exposures to a 5G-3.5 GHz signal were performed in awake restrained mice. Each mouse was habituated to the exposure conditions by progressively increasing the time during which the mouse was head-fixed (from 15 min to 60 min over a week) in a red plastic tube (internal diameter: 32 mm) with the hook screwed to a small plastic post (see Figure 1A,B). Each day, a dipole antenna (SID3500, MVG, Plouzané, France) was positioned 5 mm from the animal's head in a fixed/standardized position (Figure 1B) for one hour in the vicinity of the right temporal cortex.

The exposure system was similar to the one described in a previous study [35], replacing the frequency emitted by the radiofrequency generator to generate a 5G-3.5 GHz signal corresponding to 5G NR (release 15, Digital Standards SMBVB-K444; Rohde & Schwarz, Munich, Germany) with FDD duplexing, QPSK modulation, and a 100 MHz channel bandwidth. Briefly, a radiofrequency generator emitting a 5G-3.5 GHz electromagnetic field (SMBV100B, Rohde & Schwarz) was connected to a power amplifier (ZHL-4W-422+, Mini-Circuits, Brooklin, NY, USA), a circulator (Pasternack, PE83CR1005, Irvine, CA, USA), a bidirectional coupler (Mini-circuits, ZGBDC30-372HP+, Brooklin, NY, USA) and a four-way power divider (Mini-circuits ZB4PD-462W-N+, Brooklin, NY, USA), allowing potential simultaneous exposure of four animals. A power meter (E4417A and E9323A, EPM-P Series Power Meter, Agilent, Santa Clara, CA, USA) connected to the bidirectional coupler allowed continuous measurements and monitoring of incident and reflected powers within the setup.

Each exposed mouse was matched with a PSD-exposed mouse that was in head-fixed restrained conditions next to it (at about 30 cm, Figure 1A). An antenna was also placed 5 mm from the head of the PSD-exposed mouse, but this antenna was not connected to the 5G-3.5 GHz generator.

Conclusions and Limitations

Altogether, our results show that 1h of daily head exposure to a 5G-3.5 GHz signal over a six-week period does not alter emotional state and memory performance but triggers significant modification of expression in a limited set of genes, which can potentially affect glutamatergic synapses and mitochondrial activities. We acknowledge the limitations of our study. We cannot exclude that prolongations of the head exposures beyond 6 weeks could ultimately affect the emotional state or memory abilities of the exposed mice. While the classical OF tests used in our study did not show 5G-induced change in the animals' behavior, we cannot rule out subtle alterations in emotional state that might be revealed by applying other behavioral tests, such as the elevated plus maze or emergence tests [55]. The experiments were carried out with male mice because of the impossibility of separately housing male and female mice during 5G exposure and behavioral testing. Further investigation will be required to evaluate whether female and male mice could be differentially affected by chronic exposure to 5G-3.5 GHz. In addition, our RNA-seq analyses were performed at a single time point, e.g., 24 h after the last exposure to the 5G-3.5 GHz signal. The kinetics and reversibility of the reported changes in gene expression are undetermined, and further studies are needed to assess whether and to what extent the

changes in transcript levels translate into proteomic or functional alterations in mitochondria and in glutamatergic synapses. The 5G-triggered transcriptome modifications were observed in cortical areas where the average SAR levels range around 0.43 or 0.14 W/kg. These SAR levels may be considered in light of the European safety guidelines for human head exposures [56], which set the upper SAR limit to 2 W/kg. This value is higher than the corresponding SAR levels reached in the brain due to the energy absorption in the surrounding skull tissues. Recent dosimetric analyses of human exposures to downlink RF-EMF from base stations show intracortical SAR levels attributed to environmental 5G-3.5 GHz that are much lower than the values applied in our study, being less than 1 mW/kg [9]. Further investigations are needed to specify the levels of SAR reached in the human cerebral cortex when mobile phones emitting a 5G-3.5 GHz signal are held close to the ear of the mobile-phone user.

Open access paper: https://www.mdpi.com/1422-0067/26/6/2459

--

5G Radiofrequency Exposure Reduces *PRDM16* and *C/EBP β* mRNA Expression, Two Key Biomarkers for Brown Adipogenesis

Seewooruttun C, Bouguila B, Corona A, Delanaud S, Bodin R, Bach V, Desailloud R, Pelletier A. 5G Radiofrequency Exposure Reduces *PRDM16* and *C/EBP β* mRNA Expression, Two Key Biomarkers for Brown Adipogenesis. Int J Mol Sci. 2025 Mar 20;26(6):2792. doi: 10.3390/ijms26062792.

Abstract

The widespread use of wireless technologies has raised public health concerns about the biological effects of radiofrequency (RF) exposure. Children have a higher specific absorption rate (SAR) of radiation energy compared to adults. Furthermore, brown adipose tissue (BAT) is more prevalent in infants and tends to decrease with age. Previous animal studies demonstrated a cold sensation in rats exposed to 900 MHz (second generation, 2G). UCP1-dependent thermogenesis and BAT hyperplasia are two fundamental adaptive mechanisms initiated in response to cold. This study investigated the impact of short-term exposure to 2G and fifth generation (5G) on key thermogenic and adipogenic markers related to these mechanisms while considering age and exposure duration. Juvenile and young adult Wistar rats were randomized into three subgroups: a 5G group (3.5 GHz), 2G group (900 MHz), and a control group (SHAM). They were exposed to their respective continuous-wave RF signals for 1 or 2 weeks at an intensity of 1.5 V/m, with two exposure sessions of 1 h per day. After the exposure period, a RT-qPCR was carried out to evaluate the genetic markers involved in BAT thermogenesis and adipogenesis. Two adipogenic biomarkers were affected; a fold change reduction of 49% and 32% was detected for *PRDM16* ($p = 0.016$) and *C/EBP β* ($p = 0.0002$), respectively, after 5G exposure, regardless of age and exposure duration. No significant RF effect was found on UCP1-dependent thermogenesis at a transcriptional level. These findings suggest that exposure to a 5G radiofrequency may partially disrupt brown adipocyte differentiation and thermogenic function by downregulating *PRDM16* and *C/EBP β*, possibly leading to higher cold sensitivity.

RF Exposure System

The 5G group was exposed to a **continuous-wave RF signal** set at 3.5 GHz and the 2G group to one at 900 MHz, with an exposure period of one or two weeks. The two one-hour RF exposure sessions per day were applied at an intensity of 1.5 V/m. These sessions were scheduled randomly; one in the morning

and one in the afternoon. According to the French national frequency agency (ANFR), this intensity level reflects our current environmental exposure when using wireless network technologies and mobile phones [27].

A generator (AnaPico ASPIN4010—9 kHz–4000 MHz, Glattbrugg, Switzerland), located outside the climatic chambers, was set to produce a 3.5 GHz band signal for 5G exposure. It was coupled with an amplifier RFPA (RF26003800-4x0.5W) connected to one antenna (antenna Laird multi-band CFS60383) inside the chambers. For 2G exposure, the same generator was set to 900 MHz, paired with another amplifier RFPA (RFS7002500-6x0.5) capable of emitting this RF band, and connected to two antennas (Kathrein 800-10465, Rosenheim, Germany).

The antennas were aligned horizontally in the climatic chamber, 80 cm above the exposed rats' boxes, thus at a height larger than 2.4λ and 9λ, at 900 MHz and 3.5 GHz, respectively, with λ representing the wavelength. The position of the antennas was adjusted to minimize the variation in the field amplitude within each cage and between cages.

Using an electric field probe EP600 (Narda Safety Test Solutions, Cisano sul Neva, Italy), we measured the intensity level in five different positions in each cage. An electric field of 1.6 ± 0.4 V/m was measured under 5G exposure and 1.6 ± 0.5 V/m under 2G exposure. These data were recorded on WinEP600 (Narda Safety Test Solutions, Cisano sul Neva, Italy). The transmitting device did not generate a static magnetic field. For the non-exposed groups (controls), the antennas in the adjacent climatic chamber remained unconnected to the generator.

Using these data, the mean intensity of the RF signal per cage was used to estimate the mean whole-body SAR during the experiment. The mean whole-body SAR was calculated to be 0.07 mW/kg for the 5G exposure and 0.24 mW/kg for the 2G exposure, following the method described by Mai et al. [4].

Conclusions

Our research showed a fold change decrease of 49% for PRDM16 and 32% for C/EBP β in terms of their mRNA levels after exposure to 5G. As mentioned previously, these adipogenic markers are important in the differentiation and maturation of brown adipocytes from BAT precursors, as well as the maintenance of their thermogenic capacity. In contrast, UCP1-dependent thermogenesis was not impacted by RF exposure at the transcriptional level. Most of the studied thermogenic markers showed no age-related or exposure duration-related effects associated with RF exposure, except for PPAR α and ADRβ3. This study provides new insights into the potential impact of 5G exposure on brown adipogenesis. The disrupted differentiation and thermogenic capacity of brown adipocytes through PRDM16 and C/EBP β downregulation may affect the development and characteristics of BAT, potentially leading to increased cold sensitivity after RF exposure. These findings could partially explain the physiological events related to cold stress seen after RF exposures. However, it is important to investigate the peripheral tail temperature in rats in order to confirm our hypothesis and explain the observed biological effects. Furthermore, it may provide valuable insights to help us better understand the impact of low-intensity 2G and 5G RF signals on vasomotor responses. Most studies on low-level radiofrequency exposure have predominantly focused on "non-thermal" biological effects, such as oxidative stress, genetic instability, and reproductive health, although these findings remain

heterogeneous [77]. The impact of low-intensity RFs on thermoregulation remains largely an uncharted aspect of environmental health and safety. This research addresses this gap and can help to raise public awareness about the potential health risks posed by radiofrequency electromagnetic radiation, particularly 5G, with the rise in wireless technologies.

Open access paper: https://www.mdpi.com/1422-0067/26/6/2792

--

The CB1R of mPFC is involved in anxiety-like behavior induced by 0.8/2.65 GHz dual-frequency electromagnetic radiation

Sun B, Xue T, Gao AN, Wang XY, Wu S, Liu XM, Zhang LH, Li MH, Zou DF, Gao Y, Wang CZ. The CB1R of mPFC is involved in anxiety-like behavior induced by 0.8/2.65 GHz dual-frequency electromagnetic radiation. Front Mol Neurosci. 2025 Mar 12;18:1534324. doi: 10.3389/fnmol.2025.1534324.

Abstract

As mobile phones and communication base stations become more widespread, concerns have arisen regarding the potential risks of environmental exposure to multi-frequency electromagnetic radiation (EMR) and its effects on mental health. To address these concerns, our study established a dual-frequency EMR mouse model at 0.8/2.65 GHz to explore potential molecular mechanisms and intervention targets. Our results revealed that exposure to this dual-frequency EMR significantly induced anxiety-like behavior in mice. Molecular experiments further showed a significant decrease in cannabinoid receptor type 1 (CB1R) levels in the medial prefrontal cortex (mPFC) of the mice, along with a notable reduction in the endogenous cannabinoids 2-arachidonoylglycerol and anandamide. This led to a downregulation of the entire endocannabinoid system (ECS). Additional confirmation was obtained by overexpressing and knocking down CB1R in the mPFC. We found that increasing mPFC CB1R levels could effectively reduce anxiety-like behavior, while decreasing mPFC CB1R levels exacerbated it. Furthermore, we found dual-frequency EMR induced the change of ECS in the basolateral amygdala (BLA). Notably, female mice exhibited similar behavioral phenotypes and molecular mechanisms in response to dual-frequency EMR. In summary, our study demonstrates that anxiety induced by dual-frequency EMR is closely linked to the function of the ECS in the mPFC and BLA, and that CB1R expression in the mPFC plays a significant role in modulating emotional behavior in mice.

EMR exposure equipment

The electromagnetic reverberation chamber (RC) utilized in this experiment was developed by Wu Tongning's team at the Department of Environment and Security, China Institute of Information and Communication Technology (Li et al., 2016). Constructed with reinforced concrete, the RC is a large shielded enclosure featuring highly conductive reflective walls and multiple mechanical stirrers. The stirrers' rotation alters the chamber's boundary conditions, creating a statistically uniform, isotropic, and randomly polarized electromagnetic environment. Key components of the RC include signal generators, power amplifiers, and shielding structures. The chamber can produce electromagnetic waves within a frequency range of 0 to 3 GHz. In this study, 0.8 and 2.65 GHz frequencies were applied at a dose of 4 W/kg. The electric field intensity was calculated based on the mice's average body weight, with specific experimental parameters detailed in Table 1.

Conclusions

In summary, this study demonstrates that anxiety-like behavior induced by dual-frequency EMR is closely associated with the ECS in the mPFC and BLA. Moreover, overexpression of CB1R in the mPFC significantly alleviates anxiety-like behavior in mice, while knockdown of CB1R in the mPFC exacerbates negative emotional responses (Figure 7). This research offers new insights into potential strategies for the treatment or prevention of the effects of dual-frequency EMR.

Open access paper: https://www.frontiersin.org/journals/molecular-neuroscience/articles/10.3389/fnmol.2025.1534324/full

--

Mitigating Heat-Induced Sperm Damage and Testicular Tissue Abnormalities: The Protective Role of Radiofrequency Radiation from Wi-Fi Routers in Rodent Models

Mahmoudi R, Karbalay-Doust S, Masoudi E, Jafari-Barmak M, Ghanbri A, Nikseresht M, Mortazavi SMJ, Mortazavi SA. (2024). Mitigating Heat-Induced Sperm Damage and Testicular Tissue Abnormalities: The Protective Role of Radiofrequency Radiation from Wi-Fi Routers in Rodent Models. *Journal of Biomedical Physics and Engineering*, (), -. doi: 10.31661/jbpe.v0i0.2405-1759

Abstract

Background: Radiofrequency electromagnetic fields (RF-EMF) have raised concerns due to their potential adverse effects on reproductive health. However, emerging evidence indicates that exposure to low-level RF-EMF may induce adaptive responses, rendering cells or organisms more resilient to subsequent stressors.

Objective: To investigate whether exposure to 2.45 GHz Wi-Fi radiation could mitigate heat-induced damage in the reproductive system of male rats.

Material and Methods: In this factorial experimental study, 32 adult male Wistar rats were divided into four groups: control, RF-EMF alone, heat stress alone, and RF-EMF combined with heat stress. Rats in the RF-EMF group were exposed to RF-EMF for 2 hours daily over 52 days, while those in the heat group experienced 10 minutes of heat stress per day over the same period. The 'RF-EMF + heat' group received both RF-EMF and heat exposure. After 52 days, the testes and sperm parameters were assessed.

Results: Animals exposed to 'RF-EMF + heat' combined with heat showed significant improvements in testis volume, tubular epithelium, interstitium, cell counts, sperm quality, and Leydig cells compared to those exposed to heat alone ($P<0.05$).

Conclusion: As far as we know, this is the first study to explore the potential protective effects of RF-EMF exposure against heat-induced structural abnormalities in the testes of male rats. Our findings suggest that RF-EMF exposure may mitigate heat-induced damage, possibly through the induction of adaptive responses. These results have implications for various fields, including reproductive biology, environmental health, and occupational safety, highlighting the need for further research to elucidate the underlying mechanisms.

Excerpts

The Wi-Fi router operated on power level of 1W and the Specific Absorption Rate at the distance of 30 cm in animals' head level, as reported in another publication by our team, was 0.091 W/kg [11]

The results suggest that RF-EMF exposure may lead to alterations in testicular weight, volume, and sperm parameters, consistent with previous research highlighting the potential negative effects on male reproductive health.

The observed decline in sperm parameters and germinal cell count in response to RF-EMF exposure and heat align with existing literature, indicating a potential link between environmental factors and male fertility. The increase in Leydig cells following exposure to heat underscores the intricate cellular responses to thermal stress.

Furthermore, the concept of adaptive response (AR) emerges as a potential mechanism through which cells may develop resistance to subsequent exposures to damaging agents. The induction of AR by RF-EMF and heat, as suggested by the study results, presents a fascinating avenue for further exploration into the protective mechanisms activated in response to low-level injuries.

Open access paper: https://jbpe.sums.ac.ir/article_50328.html

--

Histomorphometry and Sperm Quality in Male Rats Exposed to 2.45 GHz Wi-Fi

Vijay S, Ibrahim SF, Osman K, Zulkefli AF, Mat Ros MF, Jamaludin N, Syed Taha SMA, Hairulazam A, Jaffar FHF. Histomorphometry and Sperm Quality in Male Rats Exposed to 2.45 GHz Wi-Fi. Reproduction. 2025 Apr 1:REP-25-0048. doi: 10.1530/REP-25-0048.

Abstract

Numerous studies have documented the effect of 2.45 GHz Wi-Fi exposure on the testes and sperm quality. Nevertheless, detailed histological alterations of other male reproductive organs are underexplored. Therefore, this study aimed to evaluate detailed histological alterations of the testes, epididymis, seminal vesicle, coagulating organ, and sperms parameters following 2.45GHz Wi-Fi exposure. Eighteen adult male Sprague Dawley rats (N=18) were equally divided into three groups (n=6): Control, 4-hour, and 24-hour groups. The groups were exposed to an active router daily for 4 or 24 hours, respectively. The Control group was sham-exposed using an inactive router. The exposure lasted for eight weeks at a 20cm distance, with a power density of 0.141W/m² and a specific absorption rate (SAR) of 0.41W/Kg. Histological findings revealed vacuolation in the testes and the corpus epididymis of the 4-hour and 24-hour groups. The seminal vesicle in both exposed groups exhibited multifocal atypical hyperplasia. Besides, the seminiferous tubule diameter decreased gradually in both exposed groups, with a substantial decrease in the 24-hour group. The spermatogenesis index in 4-hour and 24-hour groups also reduced significantly. The latter result was reflected in the sperm concentration, where both groups showed a significant reduction compared to the Control group. Sperm motility also decreased significantly in the 4-hour groups. Interestingly, there was a substantial increase in sperm viability in the 24-hour group. These findings indicate that 2.45 GHz Wi-Fi exposure causes changes in the histology and

histomorphometry measurement and impairs important sperm parameters. This highlights the consequences following Wi-Fi exposure on male reproductive health.

Open access paper: https://rep.bioscientifica.com/view/journals/rep/aop/rep-25-0048/rep-25-0048.xml

--

The Frequency of a Magnetic Field Determines the Behavior of Tumor and Non-Tumor Nerve Cell Models

López de Mingo I, Rivera González MX, Ramos Gómez M, Maestú Unturbe C. The Frequency of a Magnetic Field Determines the Behavior of Tumor and Non-Tumor Nerve Cell Models. Int J Mol Sci. 2025 Feb 26;26(5):2032. doi: 10.3390/ijms26052032.

Abstract

The involvement of magnetic fields in basic cellular processes has been studied for years. Most studies focus their results on a single frequency and intensity. Intensity has long been the central parameter in hypotheses of interaction between cells and magnetic fields; however, frequency has always played a secondary role. The main objective of this study was to obtain a specific frequency that allows a reduction in the viability and proliferation of glioblastoma (CT2A) and neuroblastoma (N2A) cell models. These were compared with an astrocyte cell model (C8D1A) (nontumor) to determine whether there is a specific frequency of response for each of the cell lines used. The CT2A, C8D1A, and N2A cell lines were exposed to a magnetic field of 100 µT and a variable frequency range between 20 and 100 Hz for 24, 48 and 72 h. The results fit a biological window model in which the viability and proliferation of N2A and CT2A cells decrease statistically significantly in a 50 Hz center of value window. In addition, the non-tumor cell model showed different behavior from tumor cell models depending on the applied frequency. These results are promising in the use of magnetic fields for therapeutic purposes.

Conclusions

It has been demonstrated that there are certain frequencies that can reduce the proliferation and viability of tumor cell models of nervous tissue. Exposure of cell models to different magnetic field frequency values produces a cellular response in viability and proliferation that responds to a “biological window” model centered at 50 Hz for the tumor models used. This window is cell-type specific. Astrocytes, the non-tumor cell line of comparison, show an increase in viability at 20 and 40 Hz. This allows us to consider the use of specific ’bioactive’ frequencies in therapeutic applications for different pathologies such as tumor development (with the consequent decrease in viability and proliferation) or neurodegenerative diseases (with the increase in viability and proliferation of the astrocytic network).

Open access paper: https://www.mdpi.com/1422-0067/26/5/2032

--

20 kHz Magnetic Field Emission of Induction Cooking Heaters

Kamata K, Haga A. 20kHz Magnetic Field Emission of Induction Cooking Heaters. EMC Europe 2004, International Symposium on Electromagnetic Compatibility, Eindhoven, The Netherlands, 2004, pp. 1-6, doi: 10.23919/EMC.2004.10805977.

Abstract

The distribution of the magnetic field leakage produced by an induction cooking heater (IH) is measured and analyzed using the finite element method. First, the dependence of the amplitude of leaked magnetic fields on the distance from the IH was measured with two S-type pans, one S-type pan, and one L-type pan and compared with the ICNIRP limit of general public exposure to 20 kHz magnetic fields. Next, the validity of the analysis is confirmed by comparing with measurement. The effects of pan size, etc. on magnetic field leakage are analyzed. The analysis makes clear the reason why the measured leaked magnetic fields from the IH are different with various types of pans. The data presented in this paper should be useful in understanding the levels of magnetic field produced by IHs, and also in estimating magnetic field exposure in homes and workplaces.

Conclusion

Four different lHs made by four manufacturers were surveyed for their magnetic field characteristics. The magnetic field levels of IH appliances as a function of distance were presented in graphical form. All sets of measurements were carried out at the fundamental frequency of 20kHz and narrow-band rms levels of magnetic flux densities expressed in µT. The magnetic field densities generated by the THs of manufacturers A, B, C and D are different. The maximum magnetic field measurement was 16 µT in the case of manufacturer A with two S-type pans at X = Z= 0, Y= -20 cm.

The analysis showed that the basic model (two S- type pans) had the most magnetic leakage followed by one S-type pan, and two L-type pans. The analysis result was in agreement with the measurement results.

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=10805977&isnumber=10805570

--

Extremely low frequency magnetic field distracts zebrafish from a visual cognitive task

Ziegenbalg L, Güntürkün O, Winklhofer M. Extremely low frequency magnetic field distracts zebrafish from a visual cognitive task. Sci Rep. 2025 Mar 12;15(1):8589. doi: 10.1038/s41598-025-90194-x.

Abstract

Electromagnetic fields emitted from overhead power lines and subsea cables are widely regarded to be a disruptive factor for animals using the natural magnetic field as orientation cue for guiding their directed movements. However, it is not known if anthropogenic electromagnetic fields also have the potential to disturb animals attending to information from other sensory modalities. To find out, we trained adult zebrafish (*Danio rerio*) individually to perform avoidance behavior in response to a visual signal (green LED light spot), which in the exposure group was presented simultaneously with a sinusoidally changing magnetic field (0.3 Hz, group A: 0.015 mT, group B: 0.06 mT). Despite the salience of the visual signal, which was both sufficient and necessary to elicit conditioned avoidance responses, the 0.06 mT magnetic condition had a negative impact on learning performance and response behavior. This suggests that extremely low frequency technical magnetic fields of Earth strength amplitude can act as cross-modal distractor that diverts the attention of animals away from environmentally relevant cues

based on nonmagnetic sensory modalities. Our research highlights the need to study the role of anthropogenic magnetic fields as sensory pollutant beyond the scope of magnetic orientation behavior.

Open access paper: https://www.nature.com/articles/s41598-025-90194-x

--

Learned magnetic map cues and two mechanisms of magnetoreception in turtles

Goforth KM, Lohmann CMF, Gavin A, Henning R, Harvey A, Hinton TL, Lim DS, Lohmann KJ. Learned magnetic map cues and two mechanisms of magnetoreception in turtles. Nature. 2025 Feb;638(8052):1015-1022. doi: 10.1038/s41586-024-08554-y.

Abstract

Growing evidence indicates that migratory animals exploit the magnetic field of the Earth for navigation, both as a compass to determine direction and as a map to determine geographical position1. It has long been proposed that, to navigate using a magnetic map, animals must learn the magnetic coordinates of the destination2,3, yet the pivotal hypothesis that animals can learn magnetic signatures of geographical areas has, to our knowledge, yet to be tested. Here we report that an iconic navigating species, the loggerhead turtle (Caretta caretta), can learn such information. When fed repeatedly in magnetic fields replicating those that exist in particular oceanic locations, juvenile turtles learned to distinguish magnetic fields in which they encountered food from magnetic fields that exist elsewhere, an ability that might underlie foraging site fidelity. Conditioned responses in this new magnetic map assay were unaffected by radiofrequency oscillating magnetic fields, a treatment expected to disrupt radical-pair-based chemical magnetoreception4-6, suggesting that the magnetic map sense of the turtle does not rely on this mechanism. By contrast, orientation behaviour that required use of the magnetic compass was disrupted by radiofrequency oscillating magnetic fields. The findings provide evidence that two different mechanisms of magnetoreception underlie the magnetic map and magnetic compass in sea turtles.

Open access paper: https://www.nature.com/articles/s41586-024-08554-y

https://www.nature.com/articles/s41586-024-08554-y?linkId=12930458

--

Effects of moderate static magnetic fields on voltage-gated potassium ion channels in sympathetic neuron-like PC12 cells

Kaneda E, Kawai T, Okamura Y, Miyagawa S. Effects of moderate static magnetic fields on voltage-gated potassium ion channels in sympathetic neuron-like PC12 cells. Physiol Rep. 2025 Mar;13(6):e70236. doi: 10.14814/phy2.70236.

Abstract

While exposure of moderate static magnetic fields (SMF) can alter neuronal excitability, the effects on sympathetic neurons remain underexplored. This study investigates the effects of moderate SMF on Kv channels in the plasma membrane of sympathetic neuron-like PC12 cells. The current density of Kv

channels was significantly lower in the 18-h magnet-exposed group, with effects persisting even after the magnet was removed before patch-clamp measurements. The current density of outward current in the presence of TEA was not different between the two groups, indicating that magnetic field affects TEA-sensitive Kv channels. To further explore these changes, RNA sequencing was performed on samples from both the Sham and 18-h magnet-exposed groups, identifying 37 moderate SMF-sensitive genes. Changes in mRNA expression levels and KEGG analysis suggested that pathways involved in the inhibition of neuronal excitability, such as GABAB receptor activation and Kir3 channel opening, may be more likely to be activated. In conclusion, moderate SMF is strongly associated with reduced current density in PC12 cells, particularly affecting Kv channels. The present study provides fundamental information on the influence of long-term SMF exposure on the excitability of sympathetic neurons.

Open access paper: https://physoc.onlinelibrary.wiley.com/doi/10.14814/phy2.70236

--

Enhancement Effect of Static Magnetic Field on Bactericidal Activity

Zhang M, Song Y, Wang J, Shi X, Chen Q, Ding R, Mou J, Fang H, Zhou Y, Chen R. Enhancement Effect of Static Magnetic Field on Bactericidal Activity. Small. 2025 Mar 22:e2412334. doi: 10.1002/smll.202412334.

Abstract

The biological effects of magnetic fields are pervasive in microorganisms, with significant attention given to alternating magnetic fields (AMFs). However, AMFs induce electrical and magnetothermal effects, which complicate the interpretation of magnetic field-induced biological effects and introduce uncertainties regarding cytotoxicity in practical applications. The static magnetic field (SMF) with few variables and high biocompatibility presents a promising alternative for both understanding biological mechanisms and ensuring safe applications, but has a remaining problem on weak interactions with microorganisms. Here we show that the combination of SMF with paramagnetic calcium-polypyrrole nanoparticles (Ca-PPy) remarkably enhances bactericidal activity. Our experiments indicate that the synergistic action of SMF and Ca-PPy significantly promotes the generation of reactive oxygen species (ROS), i.e., singlet oxygen and superoxide anion radicals, in Escherichia coli (E. coli) and Staphylococcus aureus (S. aureus), coupled with the physical disruption of bacterial membrane, exhibiting the extraordinary bactericidal performance (the bactericidal rate is over 94%). The mechanism disclosed by computations is that the singlet-to-triplet transition of radical pairs can be increased by the introduction of magnetic fields. These findings offer new insights into the biological effects of magnetic fields and pave the way for their safe, highly effective use in bactericidal applications.

https://pubmed.ncbi.nlm.nih.gov/40119825/

--

Magneto-oncology: a radical pair primer

Hore PJ. Magneto-oncology: a radical pair primer. Front Oncol. 2025 Mar 7;15:1539718. doi: 10.3389/fonc.2025.1539718

Abstract

There are few well-established biophysical mechanisms by which external magnetic fields can influence the biochemistry of molecules in living systems. The radical pair mechanism is arguably the most promising. In this mini-review I summarize the characteristics of radical pairs in a way that may be useful to those engaged in the field of magneto-oncology. The intention is to help researchers decide whether an observed biomedical magnetic field effect could have its origin in radical pair biochemistry. Armed with a physically plausible interaction mechanism, it may be possible to devise and refine a theoretical model and thereby iteratively optimise therapeutic protocols. Such an approach may also help identify experimental artefacts.

Open access paper:
https://www.frontiersin.org/journals/oncology/articles/10.3389/fonc.2025.1539718/full

--

Perspectives on terahertz honey bee sensing

Prokscha A, Sheikh F, Jalali M *et al.* Perspectives on terahertz honey bee sensing. *Sci Rep* 15, 10638 (2025). doi: 10.1038/s41598-025-91630-8.

Abstract

Terahertz (THz) technology provides precise monitoring capabilities in dynamic environments, offering unique insights into insect habitats. Our study focuses on environmental monitoring of European honey bees (Apis mellifera) through a combination of measurements and simulations. Initially, the dielectric material properties of honey bee body parts are characterized across the spectral range of 1–500 GHz to collect heterogeneous empirical data. To extend the study, honey bee mockups made from polyamide 12 (PA12) and epoxy resin are employed and validated as effective substitutes for real bees through comparative scattering analyses. The research further explores radar cross-section (RCS), imaging, and spectral properties using advanced THz technologies, including resonant tunneling diodes (RTDs) operating at 250 GHz and THz time-domain spectroscopy (THz-TDS) for frequencies exceeding 250 GHz. High-resolution imaging, utilizing a 450 GHz bandwidth, captures intricate anatomical features of both real and 3D-printed bees, showcasing the potential of THz technology for detailed environmental monitoring. Finally, simulations at 300 GHz assess the dosimetry and feasibility of non-invasive, continuous monitoring approaches based on the heterogeneous honey bee model.

Open access paper: https://www.nature.com/articles/s41598-025-91630-8

--

Sensation of electric fields in the Drosophila melanogaster larva

Tadres D, Riedl J, Eden A, Bontempo AE, Lin J, Reid SF, Roehrich B, Williams K, Sepunaru L, Louis M. Sensation of electric fields in the Drosophila melanogaster larva. Curr Biol. 2025 Mar 28:S0960-9822(25)00299-4. doi: 10.1016/j.cub.2025.03.014.

Abstract

Electrosensation has emerged as a crucial sensory modality for social communication, foraging, and predation across the animal kingdom. However, its presence and functional role as well as the neural basis of electric field perception in Drosophila and other invertebrates remain unclear. In environments with controlled electric fields, we identified electrosensation as a new sense in the Drosophila melanogaster larva. We found that the Drosophila larva performs robust electrotaxis: when exposed to a uniform electric field, larvae migrate toward the cathode (negatively charged elecrode) and quickly respond to changes in the orientation of the field to maintain cathodal movement. Through a behavioral screen, we identified a subset of sensory neurons located at the tip of the larval head that are necessary for electrotaxis. Calcium imaging revealed that a pair of Gr66a-positive sensory neurons (one on each side of the head) encodes the strength and orientation of the electric field. Our results indicate that electric fields elicit robust behavioral and neural responses in the Drosophila larva, providing new evidence for the significance of electrosensation in invertebrates.

Open access paper: https://www.cell.com/current-biology/fulltext/S0960-9822(25)00299-4

--

Research of the Biomedical Effects of Electromagnetic Field in Russia over 130 Years: Main Stages of Advances in Scientific Knowledge

Grigoriev OA, Ushakov IB, Alexseeva, VA. Research of the Biomedical Effects of Electromagnetic Field in Russia over 130 Years: Main Stages of Advances in Scientific Knowledge. Biology Bulletin, 2024, 51(12):1-13. doi: 10.1134/S1062359024701784.

Abstract

We have analyzed the key stages of advances in the scientific history of research into the biomedical effects of electromagnetic field (EMF) in Russia over 130 years and proved the continuity of this scientific knowledge and its passage from one generation of scientists to the next due to scientific schools, systematic research, and national scientific coordination. The article focuses mainly on the period in which planned EMF health research was formed, the basic methodological principles were founded, and scientific principles were established for understanding the nature of the biological effects of EMF and their impact on health. The basic concepts on the part of Russian scientists about the mechanism underlying the biological effects of EMF and principles of their hygienic regulation are summarized.

Excerpts

The data of radiobiological studies using complex irradiation modes suggest that microwave EMF with average EFD values ≤ 500 μW/cm2 (in continuous mode or in pulses) should be considered as a stimulus to the central nervous system. Therefore, it is fair to apply the laws of physiology on biological strength, adaptive response, and ways to enhance the biological significance of a weak stimulus [67, 68]....

According to current data, the Federal Statistics Agency Rosstat has recorded an increase in morbidity over 25 years for predicted disease groups in the population of adolescents aged 15–17 (malignant tumors, diseases of the nervous system, immunodefiency-related disorders, and diseases of the auditory and visual organs). Data from basic research, as well as assessment of the possible contribution of EMF

from cellular communications to the increase in morbidity, performed by the IARC methodology, show that the dependence of increased morbidity on the effect of EMF indeed exists and is not a random coincidence. However, it is difficult to directly determine risk due to the uncertainty of dosimetry data, the absence of modern comparison groups, and incomplete concept of acceptable risk for mass cellular technology [70]....

Conclusion

This review of the history of research of electro- magnetic biology, radiobiology, and hygiene of non-ionizing radiation in Russia cannot be considered exhaustive: the material accumulated over more than a century is vast. It does not include issues of applied research addressed by researchers of the biomedical effects of electromagnetism during that time: physiotherapy and clinical diagnostics, agriculture, including pest control and stimulation of plant growth; the influence on ecological systems, mitogenetic radiation, and the body's own EMF; solar–terrestrial electromagnetic coupling and problems of electromagnetic communication in nature; and anthropogenic electromagnetic pollution on the global natural background.

Continuity is evident in the evolution of scientific thought: analysis of the publications shows that many fundamental provisions on the biomedical effects of electromagnetism and their medical and environmental consequences were formulated before the 1960s, then confirmed in experiments, supplemented, and refined under different irradiation conditions in all frequency ranges from a constant EMF to EHF and modulated EMF. The fundamental results of planned research, which made it possible to predict effects, and the integration of electromagnetic topics into the system of radiobiology and hygiene made it possible to create a scientific theory and methodology for standardizing EMF in the 1970s–1980s. The continuity of knowledge is the main feature that indicates the continuous and consistent development of the scientific school in electromagnetic biology, radiobiology, and hygiene of nonionizing radiation from the first researchers in the Russian Empire to our contemporaries, scientists of the Russian Federation, over the past 130 years.

In our opinion, when discussing the problems of EMF biomedical effects, it is necessary to take into account that the discussion about the nature of electromagnetism, its physical basis, and the existence of "ether" or an equivalent physical concept has never ceased. The relativity of ideas about the nature of the electromagnetic factor allows for the relativity of knowledge about its interaction with living objects, especially since knowledge about the living cannot be considered complete either.

The scientific school of research into the biomedical effects of electromagnetism has been going on in the Russian Empire, Soviet Union, and Russian Federation for 130 years and has deep roots in national science. The number of individual publications on the subject of radiobiology and hygiene of nonionizing radiation in Russian has long exceeded 10 000, and the words of Prof. B.I. Davydov are currently becoming even more relevant: "a periodic systematization of the accumulated material, establishment of a unified terminology, and standardization of research methods and analysis of the resulting data are required."

--

A Critical Analysis of the World Health Organization (WHO) Systematic Review 2024 on Radiofrequency Radiation Exposure and Cancer Risks (Response to Karipidis et al. 2024)

Hardell L, Nilsson M. A Critical Analysis of the World Health Organization (WHO) Systematic Review 2024 on Radiofrequency Radiation Exposure and Cancer Risks. Journal of Cancer Science and Clinical Therapeutics. 9 (2025): 09-26. doi: 10.26502/jcsct.5079261.

Abstract

Radiofrequency (RF) radiation was in 2011 classified as a possible human carcinogen by the International Agency for Research on Cancer (IARC) at the WHO. Currently the WHO undertakes a systematic review of human studies on the cancer risks. In a publication by Karipidis et al (2024), commissioned by the WHO, it was argued that based on all available studies there would be “moderate certainty evidence” that mobile phone use “likely does not increase the risk of glioma, meningioma, acoustic neuroma, pituitary tumours, and salivary gland tumours in adults, or of paediatric brain tumours.” However, the authors have overlooked results showing increased risks for brain tumours in the most exposed groups, the most exposed part of the head, and longest latency time from first exposure to tumour diagnosis. The authors also claimed that there would be “moderate certainty evidence” that transmitters and mobile phone base stations do not increase the risk of pediatric leukaemia. These conclusions are based on selective inclusion of very few and low exposure studies. This WHO evaluation is contradicted by scientific results that show increased risks of cancer from exposure to RF-radiation from mobile and cordless phones, transmitters, and base stations. Other scientists have concluded, after reviewing the available evidence, that RF-radiation may increase the risk of cancer. This article analysis the Karipidis et al review and highlights several errors, omissions, and conflicts of interests that may explain the conclusions of no cancer risk. The flawed evaluation of scientific facts should lead to retraction of the article.

Conclusion

The Karipidis group’s conclusions on no cancer risks from use of mobile and cordless phones, or exposure to RF radiation from transmitters and base stations, are based on several errors in their interpretation of scientific results, omission of facts contradicting the conclusions and inherent conflicts of interest. Further, most of the results on which the authors base their conclusions are based on very low exposure levels not representative for the public’s exposure today and the authors have excluded or ignored results based on highest exposure categories. The conclusions by the authors of various grades of “certainty” that RF-EMF exposures do not cause cancer are unscientific and unjustified in view of the available scientific evidence. Evaluations of health risks from RF radiation should be performed by scientists without ties to ICNIRP or industry. Industrial direct or indirect ties may compromise objective and sound scientific evaluation. The serious scientific malpractice by Karipidis et al [1] with fatally flawed evaluation of radiofrequency radiation and cancer risks, as outlined in this review, should lead to retraction of the article.

--

Exposure to radiofrequency electromagnetic fields and risk of cancer: Epidemiology is not enough! (Response to Karipidis et al. 2024)

Di Ciaula A, Petronio MG, Bersani F, Belpoggi F. Exposure to radiofrequency electromagnetic fields and risk of cancer: Epidemiology is not enough! Environment International. Volume 196, 2025. doi: 10.1016/j.envint.2025.109275.

No abstract

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412025000261

--

Modern health worries and exposure perceptions of individuals reporting varying levels of sensitivity to electromagnetic fields: results of two successive surveys

Ledent M, Vatovez B, Roelandt P, Bordarie J, Dieudonné M, De Waegeneer E, Kremer C, Boucher L, Bouland C, De Clercq EM. Modern health worries and exposure perceptions of individuals reporting varying levels of sensitivity to electromagnetic fields: results of two successive surveys. Front Public Health. 2025 Feb 19;13:1536167. doi: 10.3389/fpubh.2025.1536167.

Abstract

Introduction: Individuals who claim to be affected by idiopathic environmental intolerance attributed to electromagnetic fields (EMFs) report symptoms linked to EMF exposure. Uncertainties about the causes of these symptoms often leave them seeking their own care solutions. In our connected societies, they may limit their exposure, leading to a spiral of avoidance that negatively impacts overall health. Our objective is to gain insights into the characteristics of people who report sensitivity to EMFs in an attempt to provide care guidance. This study focuses on modern health worries (MHW), behaviours, and exposure perceptions of people reporting various sensitivity levels to EMFs during the COVID-19 lockdowns, which altered habits and increased telecommunication device use.

Methods: We conducted two surveys during relaxed lockdown periods in Belgium (June/July 2020 and February/March 2021). A total of 97 and 285 participants, respectively, answered a questionnaire on sensitivity to EMFs, MHW, exposure perception, and strategies to limit EMF exposure. We applied nonparametric descriptive and multivariate statistical analyses.

Results: Higher sensitivity to EMFs correlates with greater MHW regarding EMF sources and more strategies to limit EMF exposure. However, these strategies were inconclusive, as many still felt highly exposed to EMFs.

Discussion: Given the high distress, social isolation, and professional difficulties faced by some EMF sensitive individuals, the relevance of exposure avoidance strategies is questionable. People who

perceive high sensitivity to EMFs report worries and avoidance behaviors, yet still feel highly exposed. The findings suggest exploring new care avenues.

Open access paper:

https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1536167/full

--

The protective effects of melatonin against electromagnetic waves of cell phones in animal models: A systematic review

Amiri M, Khazaie H, Mohammadi M. The protective effects of melatonin against electromagnetic waves of cell phones in animal models: A systematic review. Animal Model Exp Med. 2025 Feb 24. doi: 10.1002/ame2.12552.

Abstract

Background: Due to the widespread use of cell phone devices today, numerous research studies have focused on the adverse effects of electromagnetic radiation on human neuropsychological and reproductive systems. In most studies, oxidative stress has been identified as the primary pathophysiological mechanism underlying the harmful effects of electromagnetic waves. This paper aims to provide a holistic review of the protective effects of melatonin against cell phone-induced electromagnetic waves on various organs.

Methods: This study is a systematic review of articles chosen by searching Google Scholar, PubMed, Embase, Scopus, Web of Science, and Science Direct using the keywords 'melatonin', 'cell phone radiation', and 'animal model'. The search focused on articles written in English, which were reviewed and evaluated. The PRISMA process was used to review the articles chosen for the study, and the JBI checklist was used to check the quality of the reviewed articles.

Results: In the final review of 11 valid quality-checked articles, the effects of melatonin in the intervention group, the effects of electromagnetic waves in the case group, and the amount of melatonin in the chosen organ, i.e. brain, skin, eyes, testis and the kidney were thoroughly examined. The review showed that electromagnetic waves increase cellular anti-oxidative activity in different tissues such as the brain, the skin, the eyes, the testis, and the kidneys. Melatonin can considerably augment the anti-oxidative system of cells and protect tissues; these measurements were significantly increased in control groups. Electromagnetic waves can induce tissue atrophy and cell death in various organs including the brain and the skin and this effect was highly decreased by melatonin.

Conclusion: Our review confirms that melatonin effectively protects the organs of animal models against electromagnetic waves. In light of this conclusion and the current world-wide use of melatonin, future studies should advance to the stages of human clinical trials. We also recommend that more research in the field of melatonin physiology is conducted in order to protect exposed cells from dying and that melatonin should be considered as a pharmaceutical option for treating the complications resulting from electromagnetic waves in humans.

Conclusion

Electromagnetic radiation emitted from cell phones is proven to be harmful to various organs in animal models and the results of the current literature survey suggest that additional research should be done to achieve the best possible use of melatonin as a therapeutic agent against electromagnetic waves. Melatonin is one of the safest drugs, used for many years to help people with sleep disturbances. In order to take this research to clinical levels, it will be important to assess the primary pathophysiological theory of damage and also the mechanisms by which melatonin may protect against the specific threat, among which the most likely is by increasing the amount of antioxidants in various organs. Future reviews and interventional studies in this field should focus on determining the optimal dose and timing of melatonin administration to enhance understanding of its protective effects. Additionally, efforts should be made to develop and improve clinical trials to evaluate the benefits of melatonin in humans, enabling a deeper investigation into its potential applications and advantages in clinical settings.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/ame2.12552

--

Hypothesis: ultrasonography can document dynamic *in vivo* rouleaux formation due to mobile phone exposure

Brown RR, Biebrich B. Hypothesis: ultrasonography can document dynamic *in vivo* rouleaux formation due to mobile phone exposure. Front. Cardiovasc. Med. 10 February 2025. Volume 12. https://doi.org/10.3389/fcvm.2025.1499499

Abstract

Carrying a cellphone against the body has become commonplace in our world replete with smartphones. Acute and chronic health effects caused by these devices emitting radiofrequency radiation from multiple antennas have not been well evaluated. In this study, the popliteal vein of a healthy volunteer was imaged with ultrasonography prior to and following the placement of an idle, but active smartphone against her knee for 5 min. Pre-exposure longitudinal sonographic images demonstrate a normal anechoic lumen to the popliteal vein. Images obtained 5 min after direct skin exposure to the smartphone demonstrate a dramatic change in the acoustic appearance of the vessel. The interior of the vessel became coarsely hypoechoic with sluggish flow seen in real-time images, a typical sonographic appearance for rouleaux formation. A follow up examination performed 5 min after the subject walked around yielded continued rouleaux formation in the popliteal vein, albeit less dramatic than that observed immediately post exposure. This revolutionary *in vivo* method to assess radiofrequency radiation induced rouleaux formation should be further pursued in the general population to determine its prevalence and if its occurrence provides a unique biomarker of exposure that may predict morbidity

Excerpt

In summary, we present a subject in whom 5 min of exposure to radiofrequency radiation emitted by a smartphone causes abnormal erythrocyte aggregation “rouleaux formation” *in vivo*. This abnormality is associated with sluggish venous flow, as documented on diagnostic ultrasound obtained in real time. Although rouleaux formation is recognized as a transient phenomenon, we hypothesize that habitual

cellphone usage, common in the population, would re-expose individuals over and over again to this abnormal hematologic state. Chronic, long-term exposure to radiofrequency radiation may therefore lead to recurrent, chronic RBC aggregation and increased blood viscosity, potentially causing significant morbidity in certain patient populations, particularly diabetics and those with hypertension, ischemic heart disease, cerebrovascular insufficiency, prethrombotic states, and peripheral vascular disease. Recognizing the potential for red blood cell aggregation from radiofrequency radiation to occur in the general population is crucial. Further studies are needed to assess the incidence of this occurrence, in addition to defining which power densities and frequencies put individuals at risk.

Open access paper:

https://www.frontiersin.org/journals/cardiovascular-medicine/articles/10.3389/fcvm.2025.1499499/full

--

Three-dimensional monitoring of RBC sedimentation in external magnetic fields

Gholampour K, Moradi AR. Three-dimensional monitoring of RBC sedimentation in external magnetic fields. Biomed Opt Express. 2025 Jan 29;16(2):736-747. doi: 10.1364/BOE.545723.

Abstract

The external magnetic fields resulting from electronic devices around humans have become more prevalent nowadays, and studying their influence on living matter is a required task. Here, we experimentally model the movement of RBCs in veins under an external magnetic field; we monitor the sedimentation of multiple RBCs at different distances from a surrounding wall. The monitoring is performed in 3D by incorporation of digital holographic microscopy (DHM). DHM not only provides a 3D quantitative phase image of an RBC but also, through its numerical refocusing feature, 3D trajectories of several cells in the field of view can be obtained. Our results show that the magnetic field facilitates the sedimentation of cells, and the effect is higher in proximity to the walls. This influence is attributed to the presence of magnetic field-sensitive materials included in RBCs.

https://pmc.ncbi.nlm.nih.gov/articles/PMC11828455/

--

Investigations Into the Impact of Static Magnetic Fields on Blood Flow

Mayrovitz HN. Investigations Into the Impact of Static Magnetic Fields on Blood Flow. Cureus. 2025 Jan 26;17(1):e78007. doi: 10.7759/cureus.78007.

Abstract

Many claims are made regarding the impacts of static magnetic fields (SMFs) on biological and physiological processes. Some of these are based on scientific underpinnings, and others appear to have less evidence to support them. The present report focuses on the evidence regarding SMF's effects on blood flow. Fortuitously, the author has direct experimental experience in this area. The approach for this review was to search three major databases (Web of Science, PubMed, and Embase) for peer-reviewed articles written in English in which an SMF was used in humans or other animals and measurements of parameters related to blood flow or velocity before SMF application and either during

or after application were reported. After screening the initial 1,954 articles, 108 studies were retrieved and evaluated for relevancy. Of these, 23 were found to satisfy the inclusion criteria and be relevant. This included 10 studies on humans and 13 studies on other animals. The methods employed in many of these studies are illustrated in this review to enhance understanding of the findings. With regard to human studies, none showed an increase in blood flow, and one showed a decrease in flow. With regard to the animal studies, one showed a transient post-exposure increase that was later explained as due to an actual reduction during SMF exposure. Four studies showed a decrease, four showed no change or difference from sham-exposed animals, and four reported an increase. Of these four, two were from the same author using a method that may not have reflected a blood flow change. Based on these findings, it is concluded that claims of an SMF providing an increase in blood flow or circulation are not supported by human studies and not well supported by animal studies. However, this does not close the door to a possible effect for at least four considerations or limitations that may have impacted the absence of a positive finding in human studies: (1) the number of subjects included is relatively small, which affects the study power; (2) the duration of the SMF application of most studies was relatively short; (3) most studies were done on healthy individuals; and (4) the SMF was delivered perpendicular to the body surface, so the effects of tangential field directions are unknown. Although these provisos may impact the detection of a possible SMF effect, they do not alter the current findings, as no reviewed human study has demonstrated a statistically significant increase in blood circulation attributable to an SMF. Thus, the clinical use of an SMF to improve blood circulation is not supported by experimental evidence.

Open access paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC11859515/

--

Central Causation of Autism/ASDs via Excessive [Ca^{2+}]i Impacting Six Mechanisms Controlling Synaptogenesis during the Perinatal Period: The Role of Electromagnetic Fields and Chemicals and the NO/ONOO(-) Cycle, as Well as Specific Mutations

Pall ML. Central Causation of Autism/ASDs via Excessive [Ca^{2+}]i Impacting Six Mechanisms Controlling Synaptogenesis during the Perinatal Period: The Role of Electromagnetic Fields and Chemicals and the NO/ONOO(-) Cycle, as Well as Specific Mutations. *Brain Sciences*. 2024; 14(5):454. https://doi.org/10.3390/brainsci14050454

Abstract

The roles of perinatal development, intracellular calcium [Ca^{2+}]i, and synaptogenesis disruption are not novel in the autism/ASD literature. The focus on six mechanisms controlling synaptogenesis, each regulated by [Ca^{2+}]i, and each aberrant in ASDs is novel. The model presented here predicts that autism epidemic causation involves central roles of both electromagnetic fields (EMFs) and chemicals. EMFs act via voltage-gated calcium channel (VGCC) activation and [Ca^{2+}]i elevation. A total of 15 autism-implicated chemical classes each act to produce [Ca^{2+}]i elevation, 12 acting via NMDA receptor activation, and three acting via other mechanisms. The chronic nature of ASDs is explained via NO/ONOO(-) vicious cycle elevation and MeCP2 epigenetic dysfunction. Genetic causation often also involves [Ca^{2+}]i elevation or other impacts on synaptogenesis. The literature examining each of these steps is systematically examined and found to be consistent with predictions. Approaches that may be sed for ASD prevention or treatment are discussed in connection with this special issue: The current situation and prospects for children with ASDs. Such approaches include EMF, chemical avoidance, and

using nutrients and other agents to raise the levels of Nrf2. An enriched environment, vitamin D, magnesium, and omega-3s in fish oil may also be helpful.

Open access paper: https://www.mdpi.com/2076-3425/14/5/454

--

Impact of Microwave Exposure on Cynomolgus Monkeys: EEG and ECG Analysis

Ma L, Qiao N, Zou Y, Wang H, Wang Y, Zhi W, Zhao X, Xu X, Zhang M, Lin Z, Hu X, Wang L. Impact of Microwave Exposure on Cynomolgus Monkeys: EEG and ECG Analysis. Bioelectromagnetics. 2025 Feb;46(2):e70000. doi: 10.1002/bem.70000.

Abstract

The annual increase of microwave exposure in human environments continues to fuel debates regarding its potential health impacts. This study monitored the EEG and ECG responses of three Cynomolgus monkeys before and at 0, 3, 7, 14, and 30 days after exposure to 50 mW/cm² microwave radiation for 15 min. The findings revealed no significant differences in the power spectral densities (PSDs) of the whole brain, frontal, and temporal lobes across various frequency bands (δ, θ, α, β, low-γ, and high-γ) immediately and up to 30 days postexposure. Notable alterations were observed primarily at 14 days in the PSDs of the parietal lobe, prefrontal cortex, central zone, and occipital lobe, particularly in the θ and α bands. By Day 30, these values returned to normal ranges. ECG alterations were characterized by changes in T-wave shape and amplitude. One monkey exhibited bidirectional spikes at 7 and 14 days that normalized by Day 30. Another showed similar patterns with reduced amplitude, and a third monkey displayed a towering forward wave at 14 days that persisted at 30 days. In conclusion, the administration of L-band microwave radiation at the specified dose did not result in immediate alterations to EEG and ECG, but it induced transient modifications in brain electrical activity and normalized after 30 days, which contributed to evaluate the health implications of microwave exposure in humans.

https://pubmed.ncbi.nlm.nih.gov/39912376/

--

--

Evaluation of Neuronal PARP-1 and Caspase-3 Levels in the Brain Tissue of Female Rats Exposed to Electromagnetic Fields at Different Gestational Stages

Tüfekci KK, Tatar M, Elamin AAE, Kaplan S. An Evaluation of Neuronal PARP-1 and Caspase-3 Levels in the Brain Tissue of Female Rats Exposed to Electromagnetic Fields at Different Gestational Stages. Int J Dev Neurosci. 2025 Feb;85(1):e70010. doi: 10.1002/jdn.70010.

Abstract

Fetal exposure to electromagnetic fields (EMFs) may cause marked neurocognitive impairment, although the mechanisms involved are still unclear. EMF induces region-specific neuronal and astroglial

death in the rat hippocampus. Poly (ADP-ribose) polymerase-1 (PARP-1) regulates necrosis, apoptosis and other cellular processes occurring following injury. This study, therefore, investigated whether PARP-1 also regulates neuronal responses in the hippocampus of rats subjected to EMF radiation during different developmental periods. Male and female rats were first allowed to mate in separate cages. Rats identified as pregnant were then divided into four groups. A 900-MHz EMF was applied for 2 h daily on gestational days (GD) 1-7, GD 8-14 and GD 15-21. The female offspring were sacrificed at the end of the 28th postnatal day, and PARP-1 and Caspase-3 expressions in the hippocampus were then evaluated. No special treatment was applied to the control group. In the EMF-exposed group, pyramidal neurons in the cornu ammonis (CA) region appeared normal after exposure on GD 1-7 but were darkly stained and shrunken after exposure on GD 15-21, while the majority of granular cells exhibited a normal appearance during all GDs. The group exposed to EMF on GD 15-21 exhibited strong PARP-1 and Caspase-3 immune reactivity in CA and dentate gyrus (DG) cells. Higher H-scores were also observed in the EMF-exposed group following GD 15-21 irradiation. As a result, a 900-MHz EMF application at GD 15-21, which coincides with hippocampal neurogenesis, triggered hippocampal neuron cell death by activating PARP-1 and Caspase-3.

https://pubmed.ncbi.nlm.nih.gov/39964245/

--

The Behavior of a Population of Protozoan Microorganisms under Electromagnetic Radiation from Cell Phones

Morozova, L.A., Savel'ev, S.V. The Behavior of a Population of Protozoan Microorganisms under Electromagnetic Radiation from Cell Phones. *Biophysics* 69, 955–960 (2024). doi: 10.1134/S0006350924701021.

Abstract

For the first time, the effect of a significant change in the resistance of the aquatic medium of a population of protozoan microorganisms under the influence of electromagnetic radiation from cell phones has been experimentally investigated. The results of the experiment demonstrated the biological nature of the changes occurring in the water–protozoan population system, which indicated the effect of radiation directly on the cells. This conclusion was confirmed by the same time for information exchange between individual participants of the population in the calculation and in the experiment. An extremely low concentration of protozoan cells in the experiment was established, equal to 1000 cells/L, at which the specified effect was observed. It was concluded that the method for detecting changes in the physical properties of the aquatic medium of a population of protozoa under the action of electromagnetic fields can be used to identify the mechanisms of interaction of biological systems with electromagnetic radiation.

Open access paper: https://link.springer.com/article/10.1134/S0006350924701021

--

Changes in dopamine, serotonin, their precursors and derivatives in prefrontal cortex of young male rats under chronic exposure to low-intensity Wi-Fi (in Russian)

Chueshova NV, Shchemelev VM, Vismont FI, Cheshik IA. Changes in the content of dopamine, serotonin, their precursors and derivatives in the prefrontal cortex of young male rats under chronic exposure to a low-intensity electromagnetic field. Reports of the National Academy of Sciences of Belarus. 68:5; 2024. https://doi.org/10.29235/1561-8323-2024-68-5-381-389.

Abstract

The content of a number of biogenic amines, their precursors and derivatives in the prefrontal cortex (PFC) of the brain of male Wistar rats chronically exposed to a low-intensity electromagnetic field of a Wi-Fi device (2.45 GHz, PPE_{max} – 5.83 μW/cm2 ′ $PPE_{\bar{\chi}}$ – 0.46 ± 0.37 μW/cm2 ′ 24 h/day, 7 days a week) was studied during their early postnatal development. Changes in the content of dopamine, serotonin, their precursors and metabolites in the PFC of the brain were established, which determines the development of neurotransmitter disorders in the central nervous system, and subsequently the occurrence of behavioral disorders, deterioration of social adaptation and cognitive abilities. The revealed changes in the dopaminergic system of the PFC of the brain in rats at the early stages of their postnatal development indicate the need for further hygienic assessment of the safety of long-term exposure to electromagnetic radiation on the body, especially on the brain. Further research in this direction will allow us to determine the neurotransmitter mechanisms underlying the adverse effects of low-intensity electromagnetic fields on the central nervous system, assess their danger and correct the existing maximum permissible levels of electromagnetic radiation taking into account the reaction of the neurotransmitter systems of the brain structures to this type of exposure.

Open access paper: https://doklady.belnauka.by/jour/article/view/1213

--

A Novel Method for Achieving Precision and Reproducibility in a 1.8 GHz Radiofrequency Exposure System That Modulates Intracellular ROS as a Function of Signal Amplitude in Human Cell Cultures

Dahon C, Aguida B, Lebon Y, Le Guen P, Dangremont A, Meyer O, Citerne J-M, Pooam M, Raad H, Thoradit T, et al. A Novel Method for Achieving Precision and Reproducibility in a 1.8 GHz Radiofrequency Exposure System That Modulates Intracellular ROS as a Function of Signal Amplitude in Human Cell Cultures. *Bioengineering*. 2025; 12(3):257. https://doi.org/10.3390/bioengineering12030257.

Abstract

Radiofrequency fields in the 1–28 GHz range are ubiquitous in the modern world, giving rise to numerous studies of potential health risks such as cancer, neurological conditions, reproductive risks and electromagnetic hypersensitivity. However, results are inconsistent due to a lack of precision in exposure conditions and vastly differing experimental models, whereas measured RF effects are often indirect and occur over many hours or even days. Here, we present a simplified RF exposure protocol providing a single 1.8 GHz carrier frequency to human HEK293 cell monolayer cultures. A custom-built exposure box and antenna maintained in a fully shielded anechoic chamber emits discrete RF signals which can be precisely characterized and modelled. The chosen amplitudes are non-thermal and fall within the range of modern telecommunication devices. A critical feature of the protocol is that cell cultures are exposed to only a single, short (15 min) RF exposure period, followed by detection of immediate, rapid changes in gene expression. In this way, we show that modulation of genes implicated

in oxidative stress and ROS signaling is among the earliest cellular responses to RF exposure. Moreover, these genes respond in complex ways to varying RF signal amplitudes consistent with a hormetic, receptor-driven biological mechanism. We conclude that induction of mild cellular stress and reactive oxygen species (ROS) is a primary response of human cells to RF signals, and that these responses occur at RF signal amplitudes within the range of normal telecommunications devices. We suggest that this method may help provide a guideline for greater reliability and reproducibility of research results between labs, thereby help resolve existing controversy on underlying mechanisms and outcomes of RF exposure in the general population.

Concluding Remarks and Future Perspectives

We here describe an experimental RF exposure device and protocol that present fully characterized, defined RF signals to human cells in culture. Their effects are consistent with a biological receptor-driven mechanism whereby RF exposure modulates intracellular ROS and ROS signaling pathways. This provides a testable hypothesis for the many and varied effects of RF described in the literature.

These cellular responses occur at RF signal amplitudes that are orders of magnitude below those needed to achieve thermal effects, and lie within the signal range of personal electronic devices and mobile phones. Because this human cell response to RF is not linear as a function of the RF signal amplitude, the relation between RF exposure conditions and a physiological outcome is not readily deducible; indeed a robust gene expression response may occur at one amplitude but be undetected at another signal amplitude, or even undergo the opposite response entirely (e.g., opposite expression of the same gene at different signal amplitudes). It is therefore necessary to assess physiological response to RF signal exposure at multiple signal amplitudes and wavelengths, and preferably by using a readout assay that is rapid and direct. This may help explain existing confusion and contradictions in the literature, as well as stimulate future studies on the nature of the biological reception mechanisms.

Finally, although RF exposure from cell phones and telecommunications devices has not been proven harmful in any way, there is a definite physiological response in human beings to this signal range. Risk factors may therefore exist for susceptibility to RF exposure, for instance in individuals with reduced tolerance to oxidative stress and/or who are exposed to excessive stressors in their daily life. These additive or synergistic effects may contribute to certain poorly defined syndromes such as electromagnetic hypersensitivity (EHS) that have been linked to RF exposure in rare individuals in the past [29].

Open access paper: https://www.mdpi.com/2306-5354/12/3/257

--

Oxidative stress and energy metabolism in male reproductive damage from single and combined high-power microwave exposure at 1.5 and 4.3GHz

Li Y, Yao B, Men J, Pang Y, Gao J, Bai Y, Wang H, Zhang J, Zhao L, Xu X, Dong J, Li C, Peng R. Oxidative stress and energy metabolism in male reproductive damage from single and combined high-power microwave exposure at 1.5 and 4.3GHz. Reprod Toxicol. 2024 Nov 29;132:108759. doi: 10.1016/j.reprotox.2024.108759.

Abstract

The effect of multi-frequency electromagnetic environments on male reproduction has attracted the medical community's interest. Studies have investigated the effects and mechanisms of single-frequency microwave exposure on male reproduction, but comparative research on high-power microwave (HPM) composite and single exposure remains scarce. This study aimed to examine the effects and mechanisms of combined 1.5 GHz and 4.3 GHz microwave exposure on male reproduction. Male Wistar rats were exposed to 1.5 GHz (L-band) and 4.3 GHz (C-band) electromagnetic radiation for 15 minutes. The four groups were: sham, 10 mW/cm² L-band, 10 mW/cm² C-band, and 5 mW/cm² L-band and 5 mW/cm² C-band compound. Assessments were made on the pathological structures of testes, sperm viability, serum sex hormones, oxidative stress, and energy metabolism levels after radiation. Exposure to 1.5 GHz and 4.3 GHz microwaves individually resulted in testicular tissue damage and reduced sperm quality. There was little difference between the damage caused by HPM composite and single exposure. The exposed groups showed histological and ultrastructural changes, with reduced spermatozoa viability, motility parameters, and serum testosterone, luteinizing hormone, follicle-stimulating hormone, and serum inhibin-B on days 1 and 7 after exposure. These tended to recover partially by day 14. Adenosine triphosphate content and lactate dehydrogenase and succinate dehydrogenase activities in the exposed testicular tissue decreased, corresponding to decreased superoxide dismutase activity and increased malondialdehyde content. Both single and combined exposure to L- and C-band HPM affect the male reproductive system. Exposure to single and compound HPM shows no significant difference in risks, with oxidative stress and energy metabolism disturbances playing key roles.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0890623824002260

--

Electromagnetic pulse exposure induces neuroinflammation and blood-brain barrier disruption by activating the NLRP3 inflammasome/NF-κB signaling pathway in mice

Lin Y, Lang H, Gao P, Miao X, Guo Q, Hao Y, Ai T, Li J, Zhang J, Guo G. Electromagnetic pulse exposure induces neuroinflammation and blood-brain barrier disruption by activating the NLRP3 inflammasome/NF-κB signaling pathway in mice. Ecotoxicol Environ Saf. 2025 Feb 27;292:117972. doi: 10.1016/j.ecoenv.2025.117972.

Abstract

The electromagnetic pulse (EMP) is a widespread electromagnetic disturbance that can disrupt electronic systems and pose health risks to personnel in operational areas. The biological effects of EMP radiation, especially on the central nervous system (CNS), are not yet fully understood but are gaining attention. This study examines the impact of EMP on the CNS using microglial cells as a model system. We found that mice exposed to a field strength of 600 kV/m with 1000 pulses per day for two weeks exhibited increased levels of oxidative stress. This exposure induced a microglia polarization to the M1 state, leading to neuroinflammation and disruption of the blood-brain barrier (BBB) by the pro-inflammatory response of microglia. Further analysis revealed that the NLRP3 inflammasome/NF-κB signaling pathway modulates the pro-inflammatory mechanisms of EMP irradiation. In conclusion, our findings show that EMP irradiation triggers neuroinflammation and BBB damage via NLRP3 inflammasome/NF-κB activation. This research highlights the effects of EMP radiation on the CNS and offers valuable insights into the potential targets for biomedical protection against it.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0147651325003082?via%3Dihub

--

Influence of Super-Low-Intensity Microwave Radiation on Mesenchymal Stem Cells

Artamonov MY, Pyatakovich FA, Minenko IA. Influence of Super-Low-Intensity Microwave Radiation on Mesenchymal Stem Cells. *International Journal of Molecular Sciences*. 2025; 26(4):1705. https://doi.org/10.3390/ijms26041705

Abstract

Mesenchymal stem cells (MSCs) have emerged as a promising tool for regenerative medicine due to their multipotency and immunomodulatory properties. According to recent research, exposing MSCs to super-low-intensity microwave radiation can have a significant impact on how they behave and operate. This review provides an overview of the most recent studies on the effects of microwave radiation on MSCs with power densities that are much below thermal values. Studies repeatedly show that non-thermal mechanisms affecting calcium signaling, membrane transport, mitochondrial activity, along ion channel activation may increase MSC proliferation, differentiation along mesodermal lineages, paracrine factor secretion, and immunomodulatory capabilities during brief, regulated microwave exposures. These bioeffects greatly enhance MSC regeneration capability in preclinical models of myocardial infarction, osteoarthritis, brain damage, and other diseases. Additional study to understand microwave treatment settings, biological processes, and safety assessments will aid in the translation of this unique, non-invasive strategy of activating MSCs with microwave radiation to improve cell engraftment, survival, and tissue healing results. Microwave-enhanced MSC treatment, if shown safe and successful, might have broad relevance as a novel cell-based approach for a variety of regenerative medicine applications.

Open access paper: https://www.mdpi.com/1422-0067/26/4/1705

--

Magnetically Stimulated Myogenesis Recruits a CRY2-TRPC1 Photosensitive Signaling Axis

Iversen JN, Tai YK, Wu KY, Wong CJK, Lim HY, Franco-Obregón A. Magnetically Stimulated Myogenesis Recruits a CRY2-TRPC1 Photosensitive Signaling Axis. Cells. 2025 Feb 6;14(3):231. doi: 10.3390/cells14030231.

Abstract

The cryptochromes are flavoproteins that either individually or synergistically respond to light and magnetic field directionality as well as are implicated in circadian rhythm entrainment and development. Single brief exposures (10 min) to low energy (1.5 mT) pulsed electromagnetic fields (PEMFs) were previously shown to enhance myogenesis by stimulating transient receptor potential canonical 1 (TRPC1)-mediated Ca2+ entry, whereby downwardly directed fields produced greater myogenic enhancement than upwardly directed fields. Here, we show that growth in the dark results in myoblasts losing their sensitivity to both magnetic field exposure and directionality. By contrast, overexpressing or silencing cryptochrome circadian regulator 2 (CRY2) in myoblasts enhances or reduces PEMF responses, respectively, under conditions of ambient light. Reducing cellular flavin adenine

dinucleotide (FAD) content by silencing riboflavin kinase (RFK) attenuated responsiveness to PEMFs and inhibited selectivity for magnetic field direction. The upregulation of TRPC1 and cell cycle regulatory proteins typically observed in response to PEMF exposure was instead attenuated by upwardly directed magnetic fields, growth in the darkness, magnetic shielding, or the silencing of CRY2 or RFK. A physical interaction between CRY2 and TRPC1 was detected using coimmunoprecipitation and immunofluorescence, revealing their co-translocation into the nucleus after PEMF exposure. These results implicate CRY2 in an identified TRPC1-dependent magnetotransduction myogenic cascade.

Open access paper: https://www.mdpi.com/2073-4409/14/3/231

--

Smart Electronic Device-Based Monitoring of SAR and Temperature Variations in Indoor Human Tissue Interaction

Laganà F, Bibbò L, Calcagno S, De Carlo D, Pullano SA, Pratticò D, Angiulli G. Smart Electronic Device-Based Monitoring of SAR and Temperature Variations in Indoor Human Tissue Interaction. *Applied Sciences*. 2025; 15(5):2439. doi:10.3390/app15052439.

Abstract

The daily use of devices generating electric and magnetic fields has led to potential human overexposure in home and work environments. This paper assesses the possible effects of electric fields on human health at low and high frequencies. It presents an electronic monitoring device that captures the incidence of specific absorption rate (SAR) and temperature variation (ΔT) on the human body. The system transmits data to a cloud platform, where a feedforward neural network (FFNN) processes the received information. SAR and surface temperature values are detected in an indoor environment, monitoring stationary and moving subjects. The results effectively assess temperature distribution due to electromagnetic fields. The prototype detected temperature peaks and high SAR values when the subjects remained motionless. Predictive analysis confirms the need for workplaces with materials shielding external electromagnetic signals and attenuating internal sources. Moderate mobile phone use could lower SAR and temperature values.

Open access paper: https://www.mdpi.com/2076-3417/15/5/2439

--

Population Density and Downlink EMF Exposure Levels by Region in Korea (3G - 5G)

Lee AK, Jeon S, Wang S, Wiart J, Choi HD, Moon JI. Population Density and DL EMF Exposure Levels by Region in Korea. *2024 IEEE Asia-Pacific Microwave Conference (APMC)*, Bali, Indonesia, 2024, pp. 781-783, doi: 10.1109/APMC60911.2024.10867416.

Abstract

In 2023, the electric field strength within mobile communication bands was measured in the largest city (Seoul), a small city (Gwangju, Gyeonggi province), and a rural area (Yangpyeong, Gyeonggi province) in South Korea. The three measurement regions were selected based on population density; The population densities of Seoul, Gwangju, and Yangpyeong are about 15,550/km2, 856/km2, and 133/km2, respectively. Measurements were performed by mounting the SRM3006 antenna on the roof

of a vehicle and driving for approximately 40 km in each region. In this paper, the authors report the results of analyzing downlink RF-EMF levels in mobile communication networks currently in operation by frequency, time, and region.

https://ieeexplore.ieee.org/document/10867416

The median of total DL E-field strength for all generations was 1.75 V/m (Seoul), 0.75 V/m (Gwangju), and 0.61 V/m (Yangpyeong); The difference in DL EMF between the small city and rural area was not large, but that in Seoul was significantly higher.

Discussion and Conclusions

For most users, 5G terminals will spend a significant portion of time not connected to the base station. Therefore, regardless of whether the terminal is connected to the 5G network, it is necessary to distinguish the level of radiated EMF caused by the SSB beam(s), which is (are) the only always-on signal in 5G NR, from the 5G base station.

It is impossible to distinguish between the always-on signal radiated from 5G base stations and signals generated by the use of 5G UEs using the measurement method reported in this paper. In a future study, we plan to estimate the average SSB EMF level radiated from the 5G base station with the Synchronization Signal Reference Signal Received Power (SS-RSRP) received by the terminal. We also seek to compare the total exposure level to DL EMF by all mobile networks with previously measured levels in terms of brain dose [3].

In Korea, CDMA2000 (2G) was terminated in early 2021, and WCDMA, LTE, and 5G NR are currently in service. At the time of measurement, i.e. around November 2023, the ratio of 3G subscriptions is about 2.6% of total mobile communication subscriptions, with 4G and 5G accounting for the majority; About 58.6% and 38.8% respectively. The DL EMF level by service technology generation did not match the corresponding subscription rate.

Currently, the number of 4G mobile communication base stations installed in Seoul is estimated at about 70% of the total, and those of 3G and 5G are analyzed to have similar ratios of 10 to 20% each. Compared to the ratio of subscriptions to the 5G service, the number of base stations installed is still very low. Therefore, the DL EMF exposure will continue to change in the coming years, so continuous monitoring is necessary. Additionally, since the base station installation information can be a good parameter related to the DL EMF level, it seems important to obtain data on this.

--

Progress in the study of the effects of electromagnetic radiation on the mood and rhythm (review)

Zou DF, Li ZH, Liu YB, Wang CZ. Progress in the study of the effects of electromagnetic radiation on the mood and rhythm. Electromagn Biol Med. 2025 Feb 18:1-16. doi: 10.1080/15368378.2025.2460971.

Abstract

The ever-expanding use of a large number of electrical appliances and mobile communication systems, which outnumber the global population, emit electromagnetic radiation through mobile telephones, power stations, transmission lines, radar, microwave ovens, televisions, refrigerators, therapeutic and other electronic devices. Electromagnetic radiation has been classified by the International Agency for

Research on Cancer (IARC) as possibly carcinogenic to humans (Group 2B). A large number of research results show that short-term and long-term exposure to electromagnetic radiation can lead to anxiety, depression, decreased learning ability, memory loss, sleep rhythm disorders and other adverse effects. Sleep rhythm disorders affect many people worldwide and may be associated with psychiatric disorders such as anxiety and depression. In this review, we summarise key experiments related to the effects of electric field exposure on mood and rhythms in animal and cellular studies over the past decade, describe the effects of electromagnetic radiation on emotional behaviors and circadian rhythms in humans and mammals, and explore the relationship between electromagnetic radiation，mood and rhythms as well as its underlying mechanisms of action. Most animal studies suggest that electromagnetic radiation may affect the physiological organization and functioning of the brain, influence neurotransmitters and receptors, interfere with neuronal formation and structure, or alter associated endocrine hormones and free radicals, which may lead to the unfavorable development of psychiatric disorders and sleep rhythm disorders. This summary may provide researchers with better clues and ideas to develop therapeutic solutions with sleep disorders and depressive psychiatric disorders.

Plain Language Summary

In this paper, we summarize the effects of electromagnetic radiation on body mood and rhythm and its mechanism through the recent research status at home and abroad. Published work has investigated the effects of different substances (including neurotransmitters, neurons, endocrine hormones, and free radicals) on the body's mood and rhythm. In many studies, the thermal effects of EMR have not been fully controlled or considered, and most studies have focused on the common frequency bands used in power lines and wireless communication, while the biological effects of other frequencies and complex electromagnetic environments have been less discussed. In conclusion, our understanding of EMR's role in body mood and rhythm is quite limited, and further research is needed to answer unanswered questions.

https://pubmed.ncbi.nlm.nih.gov/39964745/

--

Modern health worries and exposure perceptions of individuals reporting varying levels of sensitivity to electromagnetic fields: results of two successive surveys

Ledent M, Vatovez B, Roelandt P, Bordarie J, Dieudonné M, De Waegeneer E, Kremer C, Boucher L, Bouland C, De Clercq EM. Modern health worries and exposure perceptions of individuals reporting varying levels of sensitivity to electromagnetic fields: results of two successive surveys. Frontiers in Public Health, 13, 2025. doi: 10.3389/fpubh.2025.1536167,

Abstract

Introduction: Individuals who claim to be affected by idiopathic environmental intolerance attributed to electromagnetic fields (EMFs) report symptoms linked to EMF exposure. Uncertainties about the causes of these symptoms often leave them seeking their own care solutions. In our connected societies, they may limit their exposure, leading to a spiral of avoidance that negatively impacts overall health. Our

objective is to gain insights into the characteristics of people who report sensitivity to EMFs in an attempt to provide care guidance. This study focuses on modern health worries (MHW), behaviours, and exposure perceptions of people reporting various sensitivity levels to EMFs during the COVID-19 lockdowns, which altered habits and increased telecommunication device use.

Methods: We conducted two surveys during relaxed lockdown periods in Belgium (June/July 2020 and February/March 2021). A total of 97 and 285 participants, respectively, answered a questionnaire on sensitivity to EMFs, MHW, exposure perception, and strategies to limit EMF exposure. We applied nonparametric descriptive and multivariate statistical analyses.

Results: Higher sensitivity to EMFs correlates with greater MHW regarding EMF sources and more strategies to limit EMF exposure. However, these strategies were inconclusive, as many still felt highly exposed to EMFs.

Discussion: Given the high distress, social isolation, and professional difficulties faced by some EMF sensitive individuals, the relevance of exposure avoidance strategies is questionable. People who perceive high sensitivity to EMFs report worries and avoidance behaviors, yet still feel highly exposed. The findings suggest exploring new care avenues.

Open access paper:

https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2025.1536167

--

Wide-band metamaterial absorber for sub-6 GHz 5G applications: Reducing specific absorption rate

Routray P, Ghosh D. Wide-band metamaterial absorber for sub-6 GHz 5G applications: Reducing specific absorption rate. AEU - International Journal of Electronics and Communications, 2025. doi: 10.1016/j.aeue.2025.155709.

Abstract

Wide coverage area, less sensitivity to noise, high data transceiving, low latency, IOT-based mass application, and increased connectivity make sub-6 GHz, a popularly used frequency band for 5G. However, the health issues due to the high amount of harmful electromagnetic radiation generated from the growing usage of the sub-6 GHz band have drawn the attention of researchers. This article presents a wide-band, polarization, and oblique angle insensitive metamaterial absorber with an absorption bandwidth of 1.9 GHz stretching from 3.1 to 5 GHz, covering bands of n77, n78, and n79 allocated for 5G. The design consists of combination of rings, split rings, and meander-crossed dipoles loaded with lumped resistors. The equivalent circuit is designed using a quasi-static approach and the design is fabricated and measured. The proposed absorber is demonstrated for the reduction of electromagnetic radiation and specific absorption rate in the entire 5G band allocated in the sub-6 GHz range.

Conclusion

The growing demands of 5G, especially in the sub-6 GHz band, can generate a large amount of EM radiation, which is harmful to living beings. This article presents a resistive loaded wideband metamaterial absorber based on the combination of rings, split rings, and meander dipoles. The

absorption bandwidth of the absorber is 1.9 GHz ranging from 3.1 to 5 GHz, which includes the bands of n77 (3.3 to 4.2 GHz), n78 (3.3 to 3.8 GHz), and n79 (4.4 to 5 GHz). The absorber is independent of polarization angle and exhibits an absorptivity of over 80% up to an oblique angle of 60◦. The equivalence of the circuit is confirmed by comparing the absorptivity plots obtained from EM simulation and circuit simulation. The agreement between measured and simulation results veri fies that the structure can be used in a practical environment. The demonstration of the proposed absorber for the reduction of EM radiation and SAR generated in the 5G band is demonstrated to be up to 95%. Hence, the proposed absorber can be used to protect living beings and also can help to solve the concerns regarding health hazards due to the widespread use of 5G in the sub-6 GHz band.

https://www.sciencedirect.com/science/article/pii/S1434841125000500

--

Network-Scale Impact of Vegetation Loss on Coverage and Exposure for 5G Networks

Schampheleer J, Huss A, Deruyck M. Network-Scale Impact of Vegetation Loss on Coverage and Exposure for 5G Networks. *IEEE Access*, vol. 13: 23902-23912, 2025, doi: 10.1109/ACCESS.2025.3538054.

Abstract

This study investigates the effects of vegetation on 5G network performance, with a particular focus on coverage, user exposure, and base station deployment strategies in an urban environment (Utrecht, The Netherlands). This is the first study to perform network planning simulations that account for vegetation and building-induced propagation challenges on a city-wide scale, providing understanding of their effects on 5G network performance and exposure. The study also explores the influence of user height, examining how vegetation's blocking and shielding effects vary with user height. By evaluating both sub-6 GHz and mmWave networks under various simulated scenarios, the research qualifies the dual role of vegetation as both a coverage barrier and a mitigator of user exposure. Key findings include a significant 14.71% reduction in coverage for sub-6 GHz networks in the presence of vegetation and a 42.98% decrease in downlink whole-body SAR in mmWave networks due to vegetation's shielding effects. Flexible base station placement is shown to effectively counteract coverage losses while maintaining stable exposure metrics, but mmWave networks remain highly sensitive to environmental obstructions. These findings emphasize the importance of incorporating vegetation and other environmental factors into network planning, especially for high-frequency 5G networks, to ensure optimal performance and limit user exposure.

Open access paper: https://ieeexplore.ieee.org/document/10870177

--

Study on field strength prediction using different models on time series from urban continuous RF-EMF monitoring

Song X, Feng W, Yang C, Djuric N, Kljajic D, Djuric S. Study on field strength prediction using different models on time series from urban continuous RF-EMF monitoring. Expert Systems with Applications. 2025. doi: 10.1016/j.eswa.2025.126963.

Abstract

State-of-the-art electromagnetic field (EMF) monitoring networks, such as the latest one – the Serbian EMF RATEL system, are able to provide continuous and daily monitoring of radio-frequency (RF) EMF levels, which is especially important for urban areas where people may spend many hours and, consequently, experience increased sensitivity to RF-EMF exposure. By generating considerable time series sets of RF-EMF data, these monitoring networks are initiating a new research topic – near-future RF-EMF prediction, which is valuable for a number of public health activities, from supplementing EMF monitoring in high-risk areas, to proactively reducing exposure times, and towards advancing pre-testing of EMF compliance. This paper investigates the impact of different models on the prediction of field strength in urban environments, where Seasonal Auto-Regressive Integrated Moving Average (SARIMA), Convolutional Neural Networks (CNN), Long Short-Term Memory (LSTM), Extreme Learning Machine (ELM), Partial Least Squares Regression (PLS) and Transformer models are considered. The prediction performance of each model is analyzed on a case study of EMF-sensitive areas in the Serbian city of Novi Sad, i.e., two kindergartens and an elementary school; however, the established framework has strong potential for generalization to other urban environments. Based on two-year long monitoring data sets, a comprehensive comparison of the six models on prediction accuracy, performance degradation rate, extreme value prediction accuracy, and training time is made, showing that the PLS model outperforms other models in predicting EMF exposure. This preliminary study may be a valuable reference for large-scale deployment in real-time monitoring systems for public health protection and may trigger additional research on this ultimate EMF topic.

https://www.sciencedirect.com/science/article/pii/S0957417425005858

--

A Survey of IEEE 802.11ax WLAN Temporal Duty Cycle for the Assessment of RF Electromagnetic Exposure (Wi-Fi 6)

Yang Y, Vermeeren G, Verloock L, Guxens M, Joseph W. A Survey of IEEE 802.11ax WLAN Temporal Duty Cycle for the Assessment of RF Electromagnetic Exposure. *Applied Sciences*. 2025; 15(5):2858. https://doi.org/10.3390/app15052858

Abstract

The increasing deployment of IEEE 802.11ax (Wi-Fi 6) networks necessitates an accurate assessment of radiofrequency electromagnetic field (RF-EMF) exposure under realistic usage scenarios. This study investigates the duty cycle (DC) and corresponding exposure levels of Wi-Fi 6 in controlled laboratory conditions, focusing on bandwidth variations, multi-user scenarios, and application types. DC measurements reveal significant variability across internet services, with FTP upload exhibiting the highest mean DC (94.3%) under 20 MHz bandwidth, while YouTube 4K video streaming showed bursts with a maximum DC of 89.2%. Under poor radio conditions, DC increased by up to 5× for certain applications, emphasizing the influence of degraded signal-to-noise ratio (SNR) on retransmissions and modulation. Weighted exposure results indicate a reduction in average electric-field strength by up to

10× when incorporating DC, with maximum weighted exposure at 4.2 V/m (6.9% of ICNIRP limits) during multi-user scenarios. These findings highlight the critical role of realistic DC assessments in refining exposure evaluations, ensuring regulatory compliance, and advancing the understanding of Wi-Fi 6's EMF exposure implications.

Open access paper: https://www.mdpi.com/2076-3417/15/5/2858

--

The Numerical Assessment of RF Human Exposure to Microwave Ovens with Contact-Type Doors

Tian R, Wei J-C, Lu M. The Numerical Assessment of RF Human Exposure to Microwave Ovens with Contact-Type Doors. *Electronics*. 2025; 14(5):873. https://doi.org/10.3390/electronics14050873

Abstract

In complex electromagnetic environments, cardiac pacemakers may be interfered with easily. Microwave ovens, as common household appliances, may display electromagnetic leakage, which may pose risks to pacemaker wearers. This work evaluates the electromagnetic exposure of pacemaker wearers under various conditions. One involves different distances from the microwave oven to the human body, and the other involves a distinct oven door gap. This work uses COMSOL Multiphysics to establish a human thoracic cavity model with a heart and unipolar pacemaker, as well as a model of a microwave oven with contact-type doors. The results show that the specific absorption rate (SAR_{10g}) and temperature increase in the thoracic cavity and heart tissue are inversely proportional to the distance from the microwave source. They are directly proportional to the oven door gap size. The induced electric field intensity, the temperature increase, and the induced voltage in the pacemaker show the same trend. When the human body is closest to the microwave oven with the largest door gap (D = 100 mm, d = 0.3 mm), the SAR_{10g} and temperature increase of the thoracic cavity and heart tissue reach their maximum values, which are significantly below the safety standards recommended by ICNIRP. Similarly, the maximum value of the temperature increase and the induced electric field intensity in the pacemaker are below the safety standard recommended by ISO 14708-3 (+2 °C) and IEC 60601-1-2 (28 V/m). The maximum induced voltage at the pacemaker electrode is 5.322 mV, which exceeds the sensing sensitivity setting recommended by ISO 14117 (2 mV) for unipolar pacemakers. These findings demonstrate that microwave ovens with contact-type doors electromagnetic radiation do not threaten human health under normal usage conditions. However, the maximum value of the induced voltage exceeds the sensing sensitivity of some unipolar pacemakers, which may affect the operation of the unipolar pacemaker. This phenomenon requires attention from clinicians and patients. We still recommend that pacemaker wearers keep a distance from microwave ovens when using them.

Conclusions

Through the simulation of the induced electric field intensity, the SAR_{10g} distribution, the temperature increase after 30 min of exposure of the human thoracic cavity, the heart tissue, and the cardiac pacemaker, and the induced voltage at the electrode under different distances and oven door gap sizes, the following conclusions were drawn:

- Human thoracic tissue: The SAR_{10g} and temperature increase in the thoracic cavity gradually decreased as the distance between the human thoracic cavity and the microwave source

increased, and they increased with the size of the oven door gap. Under a condition of a 100 mm distance from the microwave source and a 0.3 mm oven door gap, the SAR_{10g} and temperature increase in the thoracic cavity reached their maximum values, which were 0.295 W/kg and 0.008687 °C, respectively. Nevertheless, the SAR_{10g} value and temperature increase remained below the safety standard recommended by ICNIRP.

- Heart tissue: The SAR_{10g} value and temperature increase in the heart tissue were lower than those of the thoracic tissue, and the change trends in the heart tissue were similar to those in the thoracic tissue. Under the most unfavorable conditions (distance of 100 mm and a 0.3 mm oven door gap), the maximum SAR_{10g} value and temperature increase in the heart tissue were 0.000789 W/kg and 0.00001156 °C, respectively, which were far below the safety standard recommended by ICNIRP. Although the SAR_{10g} of the human thoracic cavity and heart tissue for a single exposure was below the safety standard recommended by ICNIRP, long-term exposure could have led to electromagnetic energy accumulation in the tissue, potentially causing health risks [31].

- Electromagnetic compatibility of cardiac pacemaker: The temperature increase, the induced electric field intensity, and the induced voltage of the cardiac pacemaker were also inversely proportional to the distance between the human thoracic cavity and the microwave source, and directly proportional to the size of the microwave oven door gap. The induced electric field intensity, temperature increase, and induced voltage of the cardiac pacemaker reached their highest values when the distance was 100 mm and the oven door gap was 0.3 mm, which were 27.7 V/m, 0.000892 °C and 5.322 mV, respectively. However, the induced electric field intensity and temperature increase remained below the immunity recommended by IEC 60601-1-2 (28 V/m) and the temperature rise standard recommended by the ISO 14708-3 standard (+2 °C). ISO 14117 recommends setting the sensing sensitivity of the unipolar pacemaker to 2 mV. Some manufacturers set the sensing sensitivity according to the characteristics of their products at the factory. For example, Medtronic's unipolar pacemaker is set to 2.8 mV [32], and Biotronik's unipolar pacemaker is set to 2.5 mV [33]. The maximum induced voltage used in this work was 5.322 mV. This value already exceeded the sensitivity of the aforementioned pacemakers. We hope that this conclusion is helpful to those involved in research.

To sum up, under the conditions set in this work, the electromagnetic radiation of the microwave oven with contact-type doors had a minimal impact on the human body, remaining within the safe range. However, when the human body is within 300 mm of the microwave oven, it may affect the operation of the unipolar pacemaker. It was recommended that patients implanted with medical devices, especially those with pacemakers, maintain a safe distance or replace aging microwave ovens, while also minimizing the time spent near an operating microwave. In addition, patients are advised to opt for bipolar pacemakers whenever possible to reduce potential electromagnetic exposure risks.

Open access paper: https://www.mdpi.com/2079-9292/14/5/873

--

Bio-Electromagnetic Safety Assessment of Wireless Charging Environment for Electric Vehicles

Zheng H, Zhao X. Bio-Electromagnetic Safety Assessment of Wireless Charging Environment for Electric Vehicles. *Progress In Electromagnetics Research Letters*, Vol. 122, 87-92, 2024. doi:10.2528/PIERL24081306.

Abstract

Wireless power charging technology has been developed rapidly and is extensively utilized for electric vehicle wireless charging due to its numerous over plug-in charging. The electromagnetic bio-safety of the human body in charging environment has become a significant public concern. To address this issue, this paper employs the finite element analysis method to assess the electromagnetic safety of crucial organs in a typical charging environment. Firstly, human-vehicle models in various typical postures were constructed in COMSOL, and the spatial distribution of electromagnetic fields in the critical organs was calculated in a 7.7 kW, 85 kHz charging environment. Subsequently, the electromagnetic radiation dose of each organ was calculated and compared with the ICNIRP standards. The results indicated that the electromagnetic radiation dose received by different organs is influenced by both the electromagnetic parameters and position of the organs. When the human body is positioned flat in the car, the electromagnetic radiation exposure to various organs is at its highest. Additionally, the maximum radiation dose for each organ is significantly below ICNIRP standard in a low-power wireless charging environment, supporting the commercial adoption of wireless charging technology for electric vehicles.

Open access paper: https://www.jpier.org/issues/volume.html?paper=24081306

--

Electromagnetic exposure level of pure electric vehicle inverter to human body in different seating positions

Dong X, Ren Y, Lu M. Electromagnetic exposure level of pure electric vehicle inverter to human body in different seating positions. *Radiation Protection Dosimetry*, 2025. https://doi.org/10.1093/rpd/ncaf013

The market share of pure electric vehicle (PEV) as a green transportation steadily increases as the global demand for renewable energy sources and environmentally friendly mobility continues to increase. However, during PEV operation, the inverter system, as the key power conversion device, generates strong electromagnetic field in the local space. Long-term exposure to such electromagnetic environments may have potential effects on human body. In this study, the electromagnetic environment model of the PEV body, human body and simplified inverter is established. The finite element software, COMSOL Multiphysics, is used to calculate and analyse the variations in the induction field in different tissues of the driver and rear passenger, caused by the electromagnetic field generated by the inverter system operating at maximum power. The electromagnetic exposure level of the driver and rear passenger is assessed. Results show significant differences in the electromagnetic exposure levels of different seating positions in the vehicles. The electromagnetic exposure level in the driver's body is higher than that of the rear passenger, but it does not exceed the exposure limits defined by the International Commission for Non-ionizing Radiation Protection. This finding effectively complements the study on evaluating the safety of the electromagnetic environment of PEV and improves public awareness.

https://doi.org/10.1093/rpd/ncaf013

--

Brain Disease-Modifying Effects of Radiofrequency as a Non-Contact Neuronal Stimulation Technology

Sun S, Bok J, Jang Y, Seo H. Brain Disease-Modifying Effects of Radiofrequency as a Non-Contact Neuronal Stimulation Technology. *International Journal of Molecular Sciences*. 2025; 26(5):2268. https://doi.org/10.3390/ijms26052268

Abstract

Non-invasive, non-contact, and painless methods of electrical stimulation to enhance neural function have been widely studied in recent years, particularly in the context of neurodegenerative diseases such as Alzheimer's disease (AD) and related dementias, which cause cognitive decline and other neurological symptoms. Radiofrequency (RF), which is a rate of oscillation in the range of 3 kHz to 300 GHz (3 THz), has been suggested as one potential non-contact neuronal stimulation (NCNS) technique for improving brain function. A new type of electrical stimulation uses a radiofrequency electromagnetic field (RF-EMF). RF exposure has been shown to modulate neural stimulation and influence various brain activities in in vitro and in vivo models. Recent studies have explored the effects of RF-EMF on human physiology, particularly in areas such as brain activity, cognition, and sleep behavior. In this review, we summarize recent findings about the effects of non-contact stimulations in in vitro studies, in vivo animal models, and human clinical cases.

Open access paper: https://www.mdpi.com/1422-0067/26/5/2268

--

Learned magnetic map cues and two mechanisms of magnetoreception in turtles

Goforth KM, Lohmann CMF, Gavin A, Henning R, Harvey A, Hinton TL, Lim DS, Lohmann KJ. Learned magnetic map cues and two mechanisms of magnetoreception in turtles. Nature. 2025 Feb 12. doi: 10.1038/s41586-024-08554-y.

Abstract

Growing evidence indicates that migratory animals exploit the magnetic field of the Earth for navigation, both as a compass to determine direction and as a map to determine geographical position[1]. It has long been proposed that, to navigate using a magnetic map, animals must learn the magnetic coordinates of the destination[2,3], yet the pivotal hypothesis that animals can learn magnetic signatures of geographical areas has, to our knowledge, yet to be tested. Here we report that an iconic navigating species, the loggerhead turtle (Caretta caretta), can learn such information. When fed repeatedly in magnetic fields replicating those that exist in particular oceanic locations, juvenile turtles learned to distinguish magnetic fields in which they encountered food from magnetic fields that exist elsewhere, an ability that might underlie foraging site fidelity. Conditioned responses in this new magnetic map assay were unaffected by radiofrequency oscillating magnetic fields, a treatment expected to disrupt radical-pair-based chemical magnetoreception[4-6], suggesting that the magnetic map sense of the turtle does not rely on this mechanism. By contrast, orientation behaviour that required use of the magnetic compass was disrupted by radiofrequency oscillating magnetic fields. The findings provide evidence that two different mechanisms of magnetoreception underlie the magnetic map and magnetic compass in sea turtles.

https://pubmed.ncbi.nlm.nih.gov/39939776/

--

Effects of electromagnetic radiation from offshore wind power on the physiology and behavior of two marine fishes

Xu P, Wang B, Wang Z, Jin R, Ahmad M, Shang Y, Hu M, Chen F, Khalil MF, Huang W, Wang Y. Effects of electromagnetic radiation from offshore wind power on the physiology and behavior of two marine fishes. Mar Pollut Bull. 2025 Feb 7;213:117633. doi: 10.1016/j.marpolbul.2025.117633.

Abstract

With the widespread promotion of wind power, its potential ecological impacts on marine ecosystems have raised concerns, among which electromagnetic radiation is one of the significant impacts. It is well known that the effects of electromagnetic fields on different marine organisms vary greatly, but little is known about their effects on commercially important fish species. We studied the toxicity differences between the large yellow croaker (Larimichthys crocea) and the black sea bream (Acanthopagrus schlegelii) when exposed to magnetic field strengths of 0mT, 0.5mT, 1.0mT, 1.5mT and 2.0mT. We found that under the 1.5mT and 2.0mT, the swimming velocity of L. crocea and A. schlegelii was reduced and the antioxidant enzymes in the body had a protective function on the body. Under the 2.0mT, the stress response of A. schlegelii was higher and the immune systems of both fish species were activated. The electromagnetic intensity below 2.0mT was more sensitive to L. crocea and A. schlegelii. However, electromagnetic radiation seems to have no significant effect on the nutrient absorption capacity of the organisms. After several days of recovery, all affected markers showed signs of reversibility. In summary, the experiment could provide valuable data for developing early warning systems and preventive measures to mitigate potential threats to marine life from offshore wind farms.

https://pubmed.ncbi.nlm.nih.gov/39921984/

--

The Collision between Wireless and Biology

Héroux P. The Collision between Wireless and Biology. *Heliyon*, 2025. doi: 10.1016/j.heliyon.2025.e42267.

Abstract

This article examines the historical development of the concept of energy of activation, which has been used in the past to frame the belief that non-ionizing electromagnetic radiation is harmless at non-thermal levels. The power and telecommunications industries have used two arguments to support their view that human exposures to non-thermal non-ionizing radiation is inoffensive. First, the radiation is non-ionizing. Second, the energy quanta of the radiation are too weak to overcome the competing energy of thermal motion. Those arguments rest on the Arrhenius equation (1889) and on the concept of energy of activation. Later scientific developments such as the Eyring equation (1935) and the Bennett-Chandler (1977-1978) equation on reaction rates, and the Second Law of Thermodynamics, recognized in living systems by Schrödinger as “negentropy”, all undermine these arguments. The

Second Law of Thermodynamics supplies the ionization claimed to be lacking, while some biological processes are independent of thermal motion. We contribute a new view of the physics and biology of non-ionizing electromagnetic radiation hygiene which explains many epidemiological, toxicological and scientific observations. The health effects of non-thermal non-ionizing radiation presently observed are in fact completely supported by physics. Furthermore, the expansion of wireless data rates is in direct conflict with preservation of a healthy environment.

Conclusion

Modeling living tissues as sugar/water/salt, or as separate bio-molecular components such as protein, biopolymers and solvated biomolecules excludes the most essential element of the living process, which is the continuous traffic of electrons and protons maintained by metabolism.

These currents are required by the Second Law of Thermodynamics, and their vulnerability to NTER [non-thermal electromagnetic radiation] is enhanced by the anatomy of OXPHOS [oxidative phosphorylation].

In a more complete model, and from the principle of superposition, oscillating electric or magnetic fields penetrating from the environment will alter the tissue's electron and proton currents. These can be far more easily disturbed by external electromagnetic fields than ionic species. To be biologically active, NTER does not need to produce ionized targets or electronic currents itself, but only to disturb those already provided by biology.

The present environmental perspective is sad given that a substantial engineering toolbox is available to mitigate NTER exposures: optical fiber, wire twisting, earthing practices, DC power. But a reversal of exposure tendencies will be difficult, as industry has long enjoyed the support of a critical actor, the military.

The same wireless communication techniques used to enable maneuverability in war theaters can also be used for high data rate access in civilian environments. Innovations such as 5G are useful to the military. Consequently, the military supports these developments with the goal of achieving superiority over their adversaries in a theater of war (Department of Defense 2020).

The alliance between industry and the military certainly has some practical advantages, but it has one major drawback.

If your environment is made to mimic a theater of war, it will not be a theater of health.

This paper addresses aspects of electron and proton dynamics in living systems. While it raises the subject of telecommunications signals crest factors, it ignores characteristics of common carrier modulations such as GSM and LTE, believed to be of importance in determining health impacts (NTP 2019). Refinement of modulations may minimize biological impacts, with the ultimate goal of controlling enough of those impacts to uncover new modulation schemes capable of reducing major health effects.

Open access paper: https://www.sciencedirect.com/science/article/pii/S2405844025006474

--

Two letters attacking WHO-sponsored Systematic Review on RF-EMF Exposure and Cancer by Karipidis et al. (2024) and Authors' (Inadequate) Responses

Karipidis K, Baaken D, Loney T, Blettner M, Brzozek C, Elwood M, Narh C, Orsini N, Röösli M, Paulo MS, Lagorio S. The effect of exposure to radiofrequency fields on cancer risk in the general and working population: A systematic review of human observational studies - Part I: Most researched outcomes. Environ Int. 2024 Sep;191:108983. doi: 10.1016/j.envint.2024.108983. https://www.sciencedirect.com/science/article/pii/S0160412024005695

Frank JW, Moskowitz JM, Melnick RL, Hardell L, Philips A, Héroux P, Kelley E. The Systematic Review on RF-EMF Exposure and Cancer by Karipidis et al. (2024) has Serious Flaws that Undermine the Validity of the Study's Conclusions. Environment International, 2024. doi: 10.1016/j.envint.2024.109200. Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024007876

Karipidis K, Baaken D, Loney T, Blettner M, Mate R, Brzozek C, Elwood M, Narh C, Orsini N, Röösli M, Paulo MS, Lagorio S. Response to the letter from members of the ICBE-EMF. Environment International, 2024. doi: 10.1016/j.envint.2024.109201. Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024007888

ICBE-EMF. Scientific rebuttal to the misleading responses from Karipidis et al. on the ICBE-EMF critique of their systematic review on exposure to RF-EMF and human cancer. Jan. 15, 2025. Open access response: https://icbe-emf.org/scientific-response-to-the-rebuttal-of-karipidis-et-al-to-the-icbe-emfs-criticisms-of-the-who-cell-phone-radiation-cancer-review.

Di Ciaula A, Petronio MG, Bersani F, Belpoggi F. Exposure to radiofrequency electromagnetic fields and risk of cancer: Epidemiology is not enough! Environment International, 2025. doi: 10.1016/j.envint.2025.109275. Open access response: https://www.sciencedirect.com/science/article/pii/S0160412025000261

Karipidis K, Baaken D, Loney T, Blettner M, Mate R, Brzozek C, Elwood M, Narh C, Orsini N, Röösli M, Paulo MS, Lagorio S. Response to the letter from Di Ciaula et al. Environment International, 2025. doi: 10.1016/j.envint.2025.109276. https://www.sciencedirect.com/science/article/pii/S0160412025000273

--

The effect of exposure to radiofrequency fields on cancer risk in the general and working population: A systematic review of human observational studies – Part II: Less researched outcomes

Karipidis K, Baaken D, Loney T, Blettner M, Mate R, Brzozek C, Elwood M, Narh C, Orsini N, Röösli M, Paulo MS, Lagorio S. The effect of exposure to radiofrequency fields on cancer risk in the general and working population: A systematic review of human observational studies – Part II: Less researched outcomes. Environment International, 2025. doi: 10.1016/j.envint.2025.109274.

Abstract

Background

In the framework of the World Health Organization assessment of health effects of exposure to radiofrequency electromagnetic fields (RF-EMF), we have conducted a systematic review of human observational studies on the association between exposure to RF-EMF and risk of neoplastic diseases.

Due to the extremely large number of included exposure types/settings and neoplasm combinations, we decided to present the review findings in two separate papers. In the first one we addressed the most investigated exposure-outcome pairs (e.g. glioma, meningioma, acoustic neuroma in relation to mobile phone use, or risk childhood leukemia in relation to environmental exposure from fixed-site transmitters) (Karipidis et al., 2024). Here, we report on less researched neoplasms, which include lymphohematopoietic system tumours, thyroid cancer and oral cavity/pharynx cancer, in relation to wireless phone use, or occupational RF exposure.

Methods

Eligibility criteria: We included cohort and case-control studies of neoplasia risks in relation to three types of exposure to RF-EMF: 1. exposure from wireless phone use; 2. environmental exposure from fixed-site transmitters; 3. occupational exposures. In the current paper, we focus on less researched neoplasms including leukaemia, non-Hodgkin's lymphoma and thyroid cancer in mobile phone users; lymphohematopoietic system tumours and oral cavity/pharynx cancer in exposed workers. We focussed on investigations of specific neoplasms in relation to specific exposure sources (termed exposure-outcome pair, abbreviated E-O pairs), noting that a single article may address multiple E-O pairs. Information sources: Eligible studies were identified by predefined literature searches through Medline, Embase, and EMF-Portal. Risk-of-bias (RoB) assessment: We used a tailored version of the Office of Health Assessment and Translation (OHAT) RoB tool to evaluate each study's internal validity. Then, the studies were classified into three tiers according to their overall potential for bias (low, moderate and high) in selected, predefined and relevant bias domains. Data synthesis: We synthesized the study results using random effects restricted maximum likelihood (REML) models. Evidence assessment: Confidence in evidence was assessed according to the Grading of Recommendations, Assessment, Development and Evaluations (GRADE) approach.

Results

We included 26 articles, which were published between 1988 and 2019, with participants from 10 countries, reporting on 143 different E-O pairs, including 65 different types of neoplasms. Of these, 19 E-O pairs satisfied the criteria for inclusion in quantitative syntheses of the evidence regarding the risks of leukaemia, non-Hodgkin's lymphoma or thyroid cancer in relation to mobile phone use, and the risks of lymphohematopoietic system tumours or oral cavity/pharynx cancer following occupational exposure to RF-EMF. RF-EMF exposure from mobile phones (ever or regular use vs no or non-regular use) was not associated with an increased risk of leukaemia [meta-estimate of the relative risk (mRR) = 0.99, 95 % CI 0.91–1.07, 4 studies), non-Hodgkin's lymphoma (mRR = 0.99, 95 % CI = 0.92–1.06, 5 studies), or thyroid cancer (mRR = 1.05, 95 % CI = 0.88–1.26, 3 studies). Long-term (10 + years) mobile phone use was also not associated with risk of leukaemia (mRR = 1.03, 95 % CI 0.85–1.24, 3 studies), non-Hodgkin lymphoma (mRR = 0.99, 95 % CI 0.86–1.15, 3 studies), or thyroid cancer (no pooled estimate given the small number of studies). There were not sufficient studies of any specific neoplasms to perform dose–response meta-analyses for either cumulative call time or cumulative number of calls; individual studies did not show statistically significant associations between lifetime intensity of mobile phone use and any specific neoplasm. Occupational RF-EMF exposure (exposed vs unexposed) was not associated with an increased risk of lymphohematopoietic system tumours (mRR = 1.03, 95 % CI = 0.87–1.28, 4 studies) or oral cavity/pharynx cancer (mRR = 0.68, 95 % CI 0.42–1.11, 3 studies). There were not sufficient studies of any specific neoplasms to perform meta-analysis on the intensity or duration of occupational RF-EMF

exposure; individual studies did not show statistically significant associations with either of those exposure metrics and any specific neoplasms. The small number of studies, and of exposed cases in some instances, hampered the assessment of the statistical heterogeneity in findings across studies in the meta-analyses. Based on the summary risk of bias, most studies included in the quantitative evidence syntheses were classified at moderate risk of bias. The most critical issue was exposure information bias, especially for occupational studies where the exposure characterization was rated at high risk of bias for all included studies. Outcome information bias was an issue in mortality-based occupational cohort studies investigating non-rapidly fatal neoplasms. Further, the healthy subscriber effect, and (at a lesser extent) the healthy worker effect, were identified as plausible explanations of the decreased risks observed in some studies. The association of RF-EMF exposure from wireless phone use, or workplace equipment/devices, with other important neoplasms was reported by only one or two studies per tumour, so no quantitative evidence syntheses were conducted on these outcomes. It is noted that there were generally no statistically significant exposure-outcome associations for any combinations, independently of the exposure metric and level, with a few studies reporting decreased risks (especially for smoking-related cancers). There was only one study which assessed the effect of RF-EMF exposure from fixed-site transmitters on less researched neoplasms and it reported no statistically significant associations between exposure from base stations and risk of lymphomas overall, lymphoma subtypes, or chronic lymphatic leukaemia in adults.

Conclusions

For near field RF-EMF exposure to the head from mobile phones, there was low certainty of evidence that it does not increase the risk of leukaemia, non-Hodgkin's lymphoma or thyroid cancer. For occupational RF-EMF exposure, there was very low certainty of evidence that it does not increase the risk of lymphohematopoietic system tumours or oral cavity/pharynx cancer. There was not sufficient evidence to assess the effect of whole-body far-field RF-EMF exposure from fixed-site transmitters (broadcasting antennas or base stations), or the effect of RF-EMF from any source on any other important neoplasms.

Other

This project was commissioned and partially funded by the World Health Organization (WHO). Co-financing was provided by the New Zealand Ministry of Health; the Istituto Superiore di Sanità in its capacity as a WHO Collaborating Centre for Radiation and Health; and ARPANSA as a WHO Collaborating Centre for Radiation Protection. Registration: PROSPERO CRD42021236798. Published protocol: [(Lagorio et al., 2021) DOI https://doi.org/10.1016/j.envint.2021.106828].

Excerpts

"Our conclusive statements, formulated in accordance with the GRADE guidelines 26 (Santesso et al., 2020), are provided below.

- For near field RF-EMF exposure to the head from mobile phones, there was low certainty of evidence that it does not increase the risk of leukaemia, non-Hodgkin's lymphoma or thyroid cancer.
- For occupational RF-EMF exposure, there was very low certainty of evidence that it does not increase the risk of lymphohematopoietic system tumours or risk of oral cavity/pharynx cancer.

- There was not sufficient evidence to assess whole-body far-field RF-EMF exposure from fixed-site transmitters (broadcasting antennas or base stations) or the effect of exposure to RF-EMF (from any source) on any other *important* neoplasms."

"The main limitation in this second paper on findings from our systematic review was the small number of studies per tumour type (which in some ways is inherent to this paper, that is dedicated to less researched neoplasms). A formal synthesis of the evidence was only possible for a few types of neoplasms, mainly different types of lymphohematopoietic system tumours, as well as thyroid and oral cavity/pharynx cancers. There were 54 types of neoplasms which were investigated in only one or two studies and did not satisfy the criteria for a quantitative synthesis of the evidence, but none of these showed evidence of an effect of RF EMF."

"Looking at specific sources of RF-EMF exposure, the majority of the evidence was on mobile phone use. There was only sufficient evidence to assess the effect of ever (or regular) use *vs* no (or non-regular) use, as well as of long-term (10 + years) use. There was not sufficient evidence on lifetime intensity of mobile phone use, including cumulative call time and cumulative number of calls, so we could not conduct a dose–response analysis."

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Mark Elwood has given expert advice on topics in electromagnetic fields and health, and on the objective interpretation of epidemiological and other scientific information, over many years to individuals and groups, including government ministries, environmental regulators, community groups, commercial organisations, and formal inquiries by government and professional groups including parliamentary and legal proceedings. Some of this work has been financially supported, by universities, health care organisations, research bodies, or by government, professional or commercial groups. Some work has been reported 'blind', with the client being unidentified. Susanna Lagorio was principal investigator (April 2019 – March 2020) of the research project "BRiC 2018/06 – Systematic reviews of exposure to radiofrequency fields and cancer", supported by the Italian Workers' Compensation Authority, a public no-profit entity (grant code I85B19000120005). Her employment duties involved provision of advice on health hazards from exposure to RF-EMF to the Italian Ministry of Health and Higher Health Council (she retired on August 1st, 2023). Martin Röösli's research is entirely funded by public entities or not for profit foundations. He has served as advisor on potential health effects of exposure to non-ionizing radiation to several national and international public advisory and research steering groups, including the World Health Organization, the International Agency for Research on Cancer, the International Commission on Non-Ionizing Radiation Protection, the Swiss Government (member of the working group "Mobile phone and radiation" and chair of the expert group BERENIS), the German Radiation Protection Commission (member of the committee Non-ionizing Radiation (A6) and member of the working group 5G (A630)) and the Independent Expert Group of the Swedish Radiation Safety Authority. From 2011 to 2018, M.R. was an unpaid member of the foundation board of the Swiss Research Foundation for Electricity and Mobile Communication, a non-profit research foundation at ETH Zurich. Neither industry nor nongovernmental organizations are represented on the scientific board of the foundation. Chris Brzozek and Rohan Mate as part of their employment are involved in the provision of advice to the Australian Commonwealth Government, Australian States and

Territories and the general public on the risks and health effects of exposure to ionising and non-ionising radiation.

The other authors declare that they have no known conflicts of interest.

Open access paper: https://www.sciencedirect.com/science/article/pii/S016041202500025X

--

Histopathologic effects of mobile phone radiation exposure on the testes and sperm parameters: a systematic literature review of animal studies

Assefa EM, Abdu SM. Histopathologic effects of mobile phone radiation exposure on the testes and sperm parameters: a systematic literature review of animal studies. Front Reprod Health. 2025 Jan 17;6:1515166. doi: 10.3389/frph.2024.1515166.

Abstract

Introduction: Male infertility, often attributed to insufficient production of healthy and active sperm, can be exacerbated by electromagnetic radiation emitted from mobile phones, which disrupts normal spermatogenesis and leads to a notable decline in sperm quality. The main targets of mobile phone-induced damage in the testes are Leydig cells, seminiferous tubules, and sperm cells. The aim of this systematic literature review is to identify histopathological changes in the testes due to mobile phone radiation exposure and to examine its effects on sperm parameters in experimental animals.

Methods: In this systematic review, an extensive literature search was conducted across databases such as PubMed, ScienceDirect, Hinari, and Google scholar.

Results: A total of 752 studies were identified for screening, and 18 studies were deemed eligible for data extraction. Studies have identified histopathological alterations in testicular tissue caused by mobile phone radiation, such as reduced seminiferous tubule diameter, tunica albuginea and germinal epithelial thickness, Leydig cell hypoplasia, and increased intertubular space. Consistent exposure to mobile phone radiation has been shown to significantly reduce sperm count, motility, and viability, while also increasing abnormal sperm morphology in male rats, mice, and rabbits.

Conclusion: Animal studies indicate that electromagnetic radiation from mobile phones can negatively impact testicular tissue and sperm parameters, including sperm count, motility, viability, and morphology. As a precaution, preventive measures are recommended to minimize potential risks from mobile phone exposure, and further research is needed to fully understand its effects on human reproductive health.

Open access paper: https://www.frontiersin.org/journals/reproductive-health/articles/10.3389/frph.2024.1515166

--

Laptop and tablet use and their influence on total motile sperm count parameters: are laptops linked to infertility in Jamaican men?

Sterling L, Carroll K, Harris LR. Laptop and tablet use and their influence on total motile sperm count parameters: are laptops linked to infertility in Jamaican men? Rev Int Androl. 2024 Dec;22(4):25-32. doi: 10.22514/j.androl.2024.027.

Abstract

Background: This study aims to determine the impact of laptop and tablet use on total motile sperm count (TMSC) in men being investigated for assisted reproduction.

Methods: A cross-sectional study was conducted on 156 men attending a fertility clinic in Jamaica. Routine semen analyses were performed and parameters specific to TMSC assessed. All data analyses were performed using SPSS Version 26. Logistic regression analyses were performed to independently predict the impact of quantifiable measures of laptop and tablet use. The main outcome measures were the parameters associated with TMSC.

Results: Overall, 64% of the participants reported using laptops and 36% reported using tablets. There was a significant relationship seen with time spent on laptops and time trying to conceive ($p = 0.015$). Regression analyses showed that persons who used their laptops for 2 to 5 h daily were approximately 16 times (adjusted Odds Ratio (aOR) = 15.9; 95% Confidence Interval (CI), 2.5-103.3, $p = 0.004$) more likely to be diagnosed with low semen volume (hypospermia). Although no significant association was found between total motile sperm count (TMSC) and laptop use, a trend towards significance was observed with high laptop use ($p = 0.052$), suggesting potential implications for TMSC as a predictor of pregnancy outcomes.

Conclusions: Our findings highlight the need for clinicians to take into consideration wireless device usage in men undergoing fertility investigations.

Open access paper: https://files.intandro.com/files/article/20241230-28/pdf/RIA20240615001.pdf

--

A review of effects of electromagnetic fields on ageing and ageing dependent bioeffects of electromagnetic fields

Wei X, Huang Y, Sun C. A review of effects of electromagnetic fields on ageing and ageing dependent bioeffects of electromagnetic fields. Sci Total Environ. 2025 Jan 15;963:178491. doi: 10.1016/j.scitotenv.2025.178491.

Abstract

Thanks to the progress of science and technology, human life expectancy has dramatically increased in the past few decades, but accompanied by rapid ageing of population, resulting in increased burden on society. At the same time, the living environment, especially the electromagnetic environment, has also greatly changed due to science and technology advances. The effect of artificial electromagnetic fields (EMFs) emitted from power lines, mobile phones, wireless equipment, and other devices on ageing and

ageing-related diseases are receiving increasing attention. However, the information on the relationship between EMFs and ageing and ageing related susceptibility to EMFs is fragmentary, a review is needed. Only few studies directly investigate the effect of EMFs on ageing, and we reviewed the impact of EMFs on lifespan and cellular senescence to pry whether EMFs have an effect on ageing, and reviewed the age-dependent bioeffects and health impacts of EMFs to see whether ageing would affect biological susceptibility to EMFs. The results indicated that EMFs may have an effect on longevity and cellular senescence, but the results were inconsistent which may depend on EMF types (frequency, intensity, wave shape, etc.), species, and cell lines. Ageing has an impact on the biological or health effects of EMFs; however, the results differ depending on the EMF type and the endpoint or health outcome. Age-dependent changes in free radical metabolism, ion homeostasis, gene expression, enzyme activity, and tissue biophysical properties may be the reason; however, the underlying mechanisms are not fully elucidated.

Highlights

- Electromagnetic fields (EMF) exposure has an effect on lifespan, but the effect may vary with type of EMF and animal model.
- Extremely-low-frequency EMFs exposure can affect cellular senescence in a wave shape- and/or magnetic density-dependent manner.
- Radio-frequency EMFs promotes cellular senescence in a frequency- and intensity-dependent manner.
- The influence of EMFs on cellular, animal, and human health exhibits a degree of age-dependency.

https://pubmed.ncbi.nlm.nih.gov/39818160/

Conclusion

With the intensification of ageing and the variety of EMFs in the environment continues to increase, the effects of EMFs on ageing and ageing-dependent effects of EMFs will get constant attention. Although current research is insufficient to answer this question, some important information can be gleaned. The ageing state of the cell or body plays a role in determining the biological or health effects of EMF exposure, but the exact effect depends on the specific conditions, and further investigation of the underlying mechanism is needed.

6.1. The effect of EMFs on ageing

Ageing is a complex and expanding concept (López-Otín et al., 2023); therefore, understanding the effects of EMFs on ageing is not to be done in a short time, but we can gain insights into this mystery from the study of lifespan and cellular senescence. SMFs have an effect on delaying cellular senescence (Fig. 3) and extending the lifespan of C. elegans and mice (Fig. 2); however, SMFs at high flux intensity had negative effects on C. elegans (Fig. 2). RMFs have both beneficial effects on anti-cellular senescence and prolong the lifespan of C. elegans (Fig. 2, Fig. 3). The impact of ELF-EMFs on lifespan and cellular

senescence had been found to be inconsistent, potentially dependent on wave shape and/or magnetic flux density (Fig. 2, Fig. 3). Exposure to RF-EMFs in the wireless communication bands (1800 MHz–2480 MHz) appears to have no significant impact on lifespan (Fig. 2). However, these RF-EMFs may influence cellular senescence in a frequency- and intensity-dependent manner (Fig. 3).

6.2. Ageing-dependent effect of EMFs

With the global ageing of the population, it is very meaningful to understand the age-dependent health effects of EMFs. Although the current research is still in the preliminary stage, the biological effects of SMFs, ELF-EMFs, and RF-EMFs have been reported to be age dependent (Fig. 4). SMFs have an age-dependent effect on cell apoptosis, calcium signalling and tissue hydration. ELF-EMFs have an age-dependent effect on human-derived primary cells and brain oxidative stress in animals, and epidemiological studies have shown that the relationship between ELF-EMFs and the risk of neurodegenerative disease is greater in older individuals and that the risk of carcinogenesis is greater in young individuals. RF-EMFs have an age-dependent effect on immune cells, tissue development, and brain activity but do not interfere with sleep. Importantly, RF-EMFs are more strongly related to the risk of carcinogenesis in younger age groups.

6.3. Possible mechanisms for the age-dependent effect of EMFs

Cellular senescence is the basis of ageing, so the age-dependent effect may be fundamentally due to cellular senescence (van Deursen, 2014). A deeper understanding of the molecular mechanism underlying the interactions between EMFs and cells may lead to new findings. Free radical metabolism and calcium signalling have mostly been studied for their ability to mediate the bioeffects of EMFs; thus, age-related functional changes in these two aspects might be the cause of age-dependent effects (Falone et al., 2008; Selakovic et al., 2013). Existing studies have shown that other possible mechanisms, such as age-dependent Na+/K+ pump dysfunction (Deghoyan et al., 2014) and decreased expression levels of high-affinity ouabain receptors (Narinyan et al., 2013), play important roles in the age-dependent effect of SMFs on tissue hydration. The effect of EMFs is also dependent on the biophysical properties of tissues or cells, and ageing-related differences in the biophysical properties that result in different responses to EMFs may also be the cause of age-dependent effects.

--

Effects of radiofrequency electromagnetic radiation with a focus on hematology parameters: a brief review and future research needs

N. Žura, I. Žura Žaja*, P. Perić, K. Malarić, S. Milinković Tur, N. Poljičak Milas, V. Rimac, J. Pejaković Hlede, S. Kunštek, M. Pećin, M. Vili. Effects of radiofrequency electromagnetic radiation with a focus on haematology parameters: a brief review and future research needs. Veterinarska Stanica, 56(3):375-387. 2025. doi: 10.46419/vs.56.3.6.

Abstract

The use of radiofrequency electromagnetic radiation (RF-EMR) has steadily increased since the 1950s. RF-EMR is used in medicine, industry, household appliances, security and navigation, and especially in wireless tele- communications and animal husbandry. The widespread use of RF-EMR, especially with

the introduction of 5G networks, raises concerns about potential adverse effects on human and animal health. The effects and mechanisms of RF-EMR impacts of 5G network frequencies on human and animal health are still unknown or poorly understood. Current research findings include the biological effects of RF-EMR on genotoxicity, cell proliferation, gene expression, cell signalling, cell membrane function, and the function of immune, hematopoietic, and reproductive systems. Exposure of humans and laboratory animals to RF-EMR emitted from cell phones and many other electronic devices of 4G and older technologies has been shown to have detrimental effects on blood cells and to cause changes in the complete blood count. This depends on the type of organisms exposed, sources, frequency, electric field level and duration of exposure. There is sparse data in the available literature on the effects of RF-EMR on haematology indicators and erythrocyte morphometry in domestic animals. Therefore, the aim of this scientific review is to highlight the effects of RF-EMR on haematology indicators, erythrocyte morphometry, and platelet activation in humans and animals, taking into account the findings on the effects of 5G electromagnetic radiation on these indicators. Considering the ubiquitous electromagnetic pollution, it is important to gain knowledge about the effects of RF-EMR on human and animal health. In addition, it is necessary to determine the effects following in vitro exposure of blood to RF-EMR, especially due to the storage and use of blood and blood products in transfusion medicine.

Open access paper: https://hrcak.srce.hr/file/462530

--

Waveforms of 4G and 5G Radiofrequency Signals: Are Differences Relevant to Biology or Health?

My note: Studies of **4G** and **5G** exposure have found adverse effects as well as exposure to **millimeter waves**.

Foster KR, Maxson D, Zollman PM. Waveforms of 4G and 5G Radiofrequency Signals: Are Differences Relevant to Biology or Health?. Health Physics. November 20, 2024. doi: 10.1097/HP.0000000000001895.

Abstract

This Note briefly reviews, at a level that is intended to be accessible to non-specialists, the similarities and differences between waveforms of 4G Long-Term Evolution (4G LTE) and 5G New Radio (5G NR) transmitted by cellular base stations, as a resource for health physicists and others who are engaged in public communication about cellular telephone technologies. Despite the difference in levels of controversy presently existing between 4G LTE (introduced in 2008) and 5G NR (introduced in 2019), the differences in waveform as represented by the baseband waveform are minimal, although 5G NR offers system designers a much wider choice of parameters. Transmitted radiofrequency signals in both technologies appear noise-like, in a frequency range that is narrowly contained in the assigned channel. It is concluded that the modulation differences between 4G LTE and 5G NR are unlikely to have any biological relevance, but the greater range of accessible frequencies in 5G NR suggests the need for further health research, particularly in the 5G FR2 millimeter-wave band.

Conclusion

In closing, it is noted that the topic of 5G NR and health extends beyond the technical considerations concerning modulation, waveform, and frequency, to include population exposure assessment, risk

perception and risk communication, and other topics. One factor contributing to public concerns about 5G NR cell sites is the use of small wireless facilities ("small cells") along city streets, sometimes close to residences (network operators have deployed 4G LTE and 5G NR small cells in some localities). To address this broader range of issues, in 2022, the European Union funded a large set of studies called CLUE-H that combines efforts of "more than 70 European research organizations in four research consortia" (https://emf-health-cluster.eu) to study the risks and perceived risks of 5G technology.

As communications technology continues to develop, the authors hope that high-quality targeted research will resolve the remaining gaps in our knowledge.

https://journals.lww.com/health-physics/abstract/9900/waveforms_of_4g_and_5g_radiofrequency_signals__are.200.aspx

--

Auto-induced uplink 4G and 5G RF-EMF exposure assessment using a network monitoring application in different microenvironments across seven European countries

Stroobandt B, Van Bladel H, Veludo AF, Deprez K, Aerts S, Verloock L, Thuróczy G, Politanski P, Polanska K, Tognola G, Parazzini M, Wiart J, Guxens M, Röösli M, Joseph W. Auto-induced uplink 4G and 5G RF-EMF exposure assessment using a network monitoring application in different microenvironments across seven European countries. Environmental Research, 2025, doi: 10.1016/j.envres.2025.121029.

Highlights

Assessment of uplink 4G and 5G microenvironmental RF-EMF exposure in Europe

Novel methodology using network monitoring application QualiPoc for 4G and 5G

Transmit powers were higher in villages than in big cities, with 5G lower than 4G

Base station density was suggested as key predictor of auto-induced uplink exposure

Transmit powers are related to uplink duty cycles and frequency band usage

Abstract

The auto-induced uplink (a-UL) radio-frequency electromagnetic field (RF-EMF), often the dominant part of the total RF-EMF exposure, has not been included in previous microenvironmental studies. As 5G exposure depends more on mobile phone usage, monitoring typical transmit power levels is crucial towards more accurate personal exposure assessment. This study describes spatial differences in average mobile phone transmit power and investigates the influence of uplink duty cycles and frequency band usage. A novel methodology using the network monitoring application QualiPoc in fourth-generation (4G) and non-standalone fifth-generation (5G) networks was presented. For the first time, the assessment of 4G and 5G a-UL RF-EMF exposure was conducted simultaneously in a large-scale microenvironmental study in Europe. Measurements were performed along predefined routes in 282 different microenvironments (e.g., parks, residential areas) across seven European countries, during a maximum uplink usage scenario. The Netherlands had the highest average transmit powers per microenvironment (median 20.6 dBm). Transmit powers in villages were 0.6 to 2.1 dB higher than in big cities. The study suggested that base station density is a key predictor of a-UL exposure. Comparing

technologies and frequency bands, average transmit powers for 5G were about 3.3 dB lower than for 4G and lowest for frequency bands with a time division duplexing (TDD) scheme due to the low uplink duty cycle (below 20%). This study provides crucial measurement data for epidemiologists and governments to enhance the understanding of the a-UL component of personal RF-EMF exposure.

https://www.sciencedirect.com/science/article/pii/S0013935125002804

--

Electromagnetic Field-Aware Radio Resource Management for 5G and Beyond: A Survey (Review)

Salem MA, Lim HS, Diong KS, Alaghbari KA, Zarakovitis CC, Chien SF. Electromagnetic Field-Aware Radio Resource Management for 5G and Beyond: A Survey. *Computers*. 2025; 14(2):51. doi: 10.3390/computers14020051.

Abstract

The expansion of 5G infrastructure and the deployment of large antenna arrays are set to substantially influence electromagnetic field (EMF) exposure levels within mobile networks. As a result, the accurate measurement of EMF exposure and the integration of EMF exposure constraints into radio resource management are expected to become increasingly important in future mobile communication systems. This paper provides a comprehensive review of EMF exposure evaluation frameworks for 5G networks, considering the impacts of high-energy beams, the millimeter wave spectrum, network densification and reconfigurable intelligent surfaces (RISs), while also examining EMF-aware radio resource management strategies for 5G networks and beyond, with RIS technology as an assistive factor. Furthermore, challenges and open research topics in the EMF evaluation framework and EMF-aware resource management for 5G mobile networks and beyond are highlighted. Despite the growing importance of RIS technology in enhancing mobile networks, a research gap remains in addressing specific EMF exposure considerations associated with RIS deployments. Additionally, the impact of EMF-aware radio resource allocation approaches on RIS-assisted 5G networks is still not fully understood.

Open access paper: https://www.mdpi.com/2073-431X/14/2/51

--

Measurement of Ambient Millimeter Wave Exposure Levels around Small Base Stations

My note: Averaging exposure over time, 30 minutes in this study, ensures very low power densities for 5G millimeter waves (MMW) especially when few (or no) 5G cell phones are downloading files using the MMW band. Assessing MMW exposure from a 5G base station is both a costly and tricky endeavor especially if one is concerned about peak exposures: "... when a UE was used to attract the beam toward the measurement location, the maximum instantaneous exposure measured was -60 dBm (0.8% public MPE)." This was 381 times the 30-min average maximum exposure at that location.

Bushberg JT, Butcher MJ. Measurement of Ambient Millimeter Wave Exposure Levels around Small Base Stations. Health Phys. 2025 Jan 13. doi: 10.1097/HP.0000000000001935.

Abstract

This study investigated the implementation and impact of fifth-generation (5G) wireless millimeter wave (mmW) technology. 5G offers significant advancements over previous generations and supports additional frequency bands, including mmW, to enhance mobile broadband with ultra-reliable, low-latency communications, supporting a high volume of diverse communications. This technology is expected to enable billions of new connections in the Internet of Things (IoT), fostering innovations in various sectors including healthcare, manufacturing, and education. This research contributes to the understanding and safe implementation of this transformative technology. Global adoption of 5G is rapidly increasing, with over 1.5 billion subscriptions as of 2024, projected to reach 58% of all wireless subscriptions by 2029. Despite its benefits, 5G mmW installations have raised concerns regarding exposure to electromagnetic fields. This study was conducted using a dual-polarized horn antenna and relatively inexpensive spectrum analyzers to measure typical ambient mmW radiofrequency field power densities near operational radio base stations (RBS) in urban and suburban environments. The measurements were taken at various times of the day and in different weather conditions to ensure a comprehensive understanding of the ambient mmW exposure. The study's results provide reassuring evidence that the ambient mmW exposure from RBSs is significantly lower than the safety limits set by the Federal Communications Commission (FCC) and other international standards. The exposure levels ranged from 0.0003% to 0.0082% of the public maximum permissible exposure (MPE), with the highest levels being more than 25,000 times lower than the allowed continuous public exposure. This study concludes that typical mmW exposure from 5G RBSs is minimal and substantially below established safety limits.

https://pubmed.ncbi.nlm.nih.gov/39804717/

Excerpts

Table 1 - RF hardware.

Device	**Description**	**Manufacturer/Part Number**
Spectrum Analyzer #1	Spectrum Analyzer 24-43 GHz v.2	SAF/J0GSAP541B
Spectrum Analyzer #2	Spectrum Compact 24-40 GHz v.2	SAF/J0GSAP741B
Antenna	18.0-40.0GHz Dual Polarization Horn Antenna	A-Info/LB-SJ-180400-KF
RF Cable Assembly (x2)	Sucoflex 102 500 mm Cable with 2.92 mm Connectors	HUBER+SUHNER/SF102/11SK/11SK/500 mm
Right Angle Adapter (x2)	2.92 mm Plug to 2.92 mm Jack Right Angle Adapter	Dynawave/1101-9495-6200

Thirty-minute (30 min) measurements were made at three distance intervals (3-15, 15-30, and 35+ m) from the base of the RBS support structure as measured with a laser rangefinder (Fig. 3). These distance intervals were selected to highlight the fact that exposure levels can vary with distance and angle

relative to the BTS. Locations were selected for alignment with the RBS antenna panels, line of sight to the antenna, and critically in a location that allowed for 30-min uninterrupted measurements. All the criteria could not be met for all locations. As seen in Fig. 4, the middle and far locations were selected off-boresight to avoid vehicle traffic and achieve line-of-sight. Following the 30-min measurement at each location, four additional 2-min measurements were made with the horn oriented at 0°, 90°, 180°, and 270° relative to the direction of the 30-min measurement to assess the presence of reflections and mmW signals from other RBSs in the area. An example is shown in Fig. 5. The first 2 min are oriented at the RBS, and the remaining time is at the other azimuths before returning to the RBS.

At most of the measured RBSs, a 1-min sample acquisition was obtained with a mobile phone (UE) behind the horn antenna assembly. High-definition/large file-size video downloads were performed to attract the beam toward the measurement location. The increase in signal strength over the RBS signal bandwidth was used to confirm that the measurement equipment and RBS were operating as expected and to estimate the maximum exposure corresponding to the RBS operating close to full capacity. It was noted that some RBS operated at 800, 400, and 300 MHz, corresponding to eight, four, and three 100-MHz channels.

The spectrum analyzers display and record received signal power in units of decibels relative to a milliwatt (dBm). The analyzers required a little over 10 s to scan the entire 1 GHz spectrum, using overlapping 100 kHz bins spaced every 30 kHz. SAF settings are listed in Table 2. Before measurements, the spectrum analyzers, the horn antenna, and the feedlines were checked with an RF source at mmW frequencies and found to agree with the expected levels. Each frequency bin was averaged throughout the measurement, and the bin with the highest received averaged power was selected to represent the highest time-average exposure at the base station. Extrapolating from the highest received averaged power bin provides a very conservative assessment of power over the entire spectrum based on the measurements. The highest received averaged power bin was converted to power received over the operating bandwidth, considering the ratio of the bin bandwidth to the observed bandwidth of the signal and the loss of the transmission lines.

The power density for both polarizations was summed to determine the time-averaged total exposure and compared with the exposure limit. This result is presented for three distances for each RBS. RBSs in four geographic areas—Los Angeles, CA; Arlington, VA; Washington, DC; and Providence, RI—were selected. All have deployed different manufacturers' equipment by different operators, including T-Mobile and AT&T. Locations included urban commercial and residential areas.

Table 3 shows data of 30-min measurement results at different distance intervals, indicating the maximum average bin over that 30-min expressed as a percent of the General Public exposure limit. The ambient mmW measurements ranged from −77 dBm down to −94 dBm, which, when polarizations are combined, correspond to total exposure levels of 0.0082% to 0.0003% of the public maximum permissible exposure (MPE). The noise floor adjacent to the measured signal was −98 dBm (Fig. 6). By comparison, **when a UE was used to attract the beam toward the measurement location, the maximum instantaneous exposure measured was -60 dBm (0.8% public MPE) (Fig. 7)**. At the same location, the 30-min average maximum 100 kHz bin is −83 dBm (0.0021% GP). The figures show only one polarization at RBS CYLA4-7 and include the 100 MHz channel average.

Conclusions

One of the distinguishing characteristics of 5G mmW base stations is dynamic beamforming using a multi-element antenna. When the scanning beams detect a UE in its service environment, the RBS creates a narrow beam directed toward UEs requesting service. Even when an obstacle in its direct path blocks the beam, the system will use multiple directed beams for spatial multiplexing. Spectral analysis of mmW signals typically requires expensive portable spectrum analyzer equipment, costing upwards of $70,000. Substantially less expensive equipment (approximately $20,000) can be used. This report demonstrates that a cross-polarized horn and two spectrum analyzers can make power density measurements of orthogonal polarization and capture the total mmW exposure from an RBS.

RF surveys were conducted in the United States in various urban environments, using different frequency bands, operators, and equipment manufacturers. Measurements at several distances from operational radio base stations (RBS) indicate that the typical exposure from an mmW RBS at ground level was an exceedingly small fraction of the FCC public safety limit. The overall measurements of ambient mmW exposure ranged from 0.0003% to 0.0044% of the FCC public MPE. Even the highest exposure level was more than 25,000 times lower than allowed for continuous public exposure.

This work is made possible through the funding provided by the Mobile & Wireless Forum under the project grant number TBD. The authors alone are responsible for the content of this paper.

--

Explainable Machine Learning-Based Electric Field Strength Mapping for Urban Environmental Monitoring

Kiouvrekis Y, Psomadakis I, Vavouranakis K, Zikas S, Katis I, Tsilikas I, Panagiotakopoulos T, Filippopoulos I. Explainable Machine Learning-Based Electric Field Strength Mapping for Urban Environmental Monitoring: A Case Study in Paris Integrating Geographical Features and Explainable AI. *Electronics*. 2025; 14(2):254. https://doi.org/10.3390/electronics14020254

Abstract

The objective of this study is to determine the optimal machine learning model for constructing electric field strength maps across urban areas, advancing the field of environmental monitoring. These models are unique because they use a detailed dataset that goes beyond electromagnetic readings, incorporating information like population density, urbanization levels, and building characteristics. This novel approach, combined with explainable AI, helps identify the key factors affecting electromagnetic exposure. The models enable the creation of highly detailed and dynamic maps of electromagnetic pollution. These maps are not just static snapshots, they can track changes over time, evaluate the success of mitigation efforts, and provide deeper insights into how electromagnetic fields are distributed in urban areas. To construct a detailed electric field strength map, we conducted an extensive analysis using 410 machine learning models across the urban area of Paris, incorporating three fundamental approaches: k-nearest neighbors, neural networks, and decision trees. This comprehensive exploration allowed us to evaluate and optimize various model configurations, ensuring robust and accurate predictions of electric field strength across diverse urban environments. The kNN model exhibited the most consistent performance, with an RMSE of 1.63 and an SD of 0.20. The analysis indicates that kNN outperforms simple neural networks and decision trees in terms of both RMSE and performance stability. From the SHAP analysis, we conclude that the feature representing the total volume of

buildings in the area around each antenna (V) is the most significant in predicting electromagnetic field strength in the kNN regression model, consistently showing a high impact across predictions. The population density feature (POP) also demonstrates considerable influence.

Open access paper: https://www.mdpi.com/2079-9292/14/2/254

--

Effects of 5G mobile phone network electromagnetic field exposure on testicular endoplasmic reticulum stress and the protective role of coenzyme Q10 (CoQ10)

Yilmaz H, Tümkaya L, Mercantepe T, Yılmaz A, Gül F, Suzan ZT. Effects of 5G mobile phone network electromagnetic field exposure on testicular endoplasmic reticulum stress and the protective role of coenzyme Q10. Arch Med Res. 2025 Jan 6;56(4):103157. doi: 10.1016/j.arcmed.2024.103157.

Abstract

Aim: Nowadays, the electromagnetic field (EMF) has become an issue of electromagnetic pollution. This study aimed to determine the effect of 5 G Fr1 frequency band EMF waves on endoplasmic reticulum (ER) stress in testicular tissue and to demonstrate the efficacy of coenzyme Q10 (CoQ10) in suppressing the potential situation.

Materials and methods: Three groups of eight male Sprague-Dawley rats were established. In group I (control) received only saline solution by oral gavage. In group II (EMF), saline solution was administered by oral gavage and the rats were exposed to 5.9 GHz EMF for 2 h/d (09:00-11:00). In group III (EMF+ CoQ10), the EMF exposure was created in the same way as in group II and CoQ10 was given daily at 10 mg/kg (CoQ10), 2 mg/kg dissolved in corn oil and administered by oral gavage for the experimental period (30 days).

Results: Histopathological damage, including edematous regions, vascular congestion, and neutrophil inflammation accompanied by loss of spermatogenic cells, particularly spermatozoa, and spermatids, increased thiobarbituric acid (TBARS) and decreased total thiol (TT) levels were observed in the seminiferous tubules in the EMF group. Furthermore, germinal epithelial cells showing intense GRP-78 immunopositivity and spermatogenic cells indicating C/EBP homologous protein (CHOP) immunopositivity were increased in the seminiferous tubules. On the other hand, in the EMF+CoQ10 group, both histopathological and biochemical findings were observed in the opposite direction.

Conclusion: Due to its antioxidant properties and its ability to suppress oxidative stress and apoptosis, CoQ10 may be a promising candidate against 5 G EMF-induced testicular ER damage.

https://pubmed.ncbi.nlm.nih.gov/39765017/

--

Short-term exposure of 2.4 GHz electromagnetic radiation on cellular ROS generation and apoptosis in SH-SY5Y cell line and impact on developing chick embryo brain tissue

Deena K, Maadurshni GB, Manivannan J, Sivasamy R. Short-term exposure of 2.4 GHz electromagnetic radiation on cellular ROS generation and apoptosis in SH-SY5Y cell line and impact on developing chick embryo brain tissue. Mol Biol Rep. 2025 Jan 21;52(1):144. doi: 10.1007/s11033-025-10217-8.

Abstract

Background: Electromagnetic radiation (EMR) from wireless technology and mobile phones, operates at various frequencies. The present study analyses the major impact of short-term exposure to 2.4 GHz frequency EMR, using the two model systems chick embryos and SH-SY5Y cell lines. We hypothesized that exposure to this frequency would induce oxidative stress and apoptosis in neurons.

Methods and results: Chick embryos were exposed continuously to 2.4 GHz EMR for 4 h each day over a 5-day period, and comparisons were made with a control group. At the end of the exposure, brain tissues were dissected for histopathological analysis, antioxidant assays, and reactive oxygen species (ROS) detection. Additionally, SH-SY5Y cells were exposed to 2.4 GHz EMR to assess cell viability, DNA damage, and apoptosis. Our results showed that exposure to 2.4 GHz EMR induces oxidative stress in both chick embryos and the SH-SY5Y cells, though no significant tissue-level impact was observed. In SH-SY5Y cells, ROS production increased after 4 h of exposure, accompanied by moderate DNA damage and early markers of apoptosis, such as upregulation of the Bax gene. Furthermore, we observed that antioxidants, such as NAC and Mito-TEMPO, helped mitigate the cytotoxic effects of EMR in both the study models.

Conclusion: In conclusion, short-term exposure (4 h) to 2.4 GHz EMR induced moderate cellular and molecular changes, primarily oxidative stress. The oxidative stress was reduced by antioxidants, which suggests potential benefits in preventing EMR-induced cytotoxicity. Extended exposure to EMR beyond 4 h may pose adverse health risks to humans, endorsing further investigation.

https://pubmed.ncbi.nlm.nih.gov/39836269/

Conclusion

This study demonstrates the intricacy of the influential impacts of short-term exposure to 2.4 GHz electromagnetic radiation (EMR) at the cellular and tissue levels, thus shedding light on its potential biological implications. Even though there were no significant structural changes detected in histopathological studies of chick embryo brain tissues, results of scanning electron microscopy revealed micro-impacts after short-term exposure of 2.4 GHz, which induced moderate surface modifications. Similarly, antioxidant enzyme activities displayed differential responses where the levels of SOD increased, and on the other hand, CAT activity remained the same. These observations thus emphasise the tight regulation of oxidative stress responses to EMR exposure. In neuronal cell, SH-SY5Y, a marked increase in ROS generation was accompanied by DNA damage and upregulation of apoptotic markers, especially the Bax gene. These findings show that EMR causes oxidative stress and triggers the activation of apoptosis-related pathways stating that mitochondria have a role in these events. Interestingly, the antioxidant protection by NAC and Mito-TEMPO shows a potential area of intervention for the abrogation of oxidative stress due to EMR.

--

Compound exposure of 2.8 GHz and 9.3 GHz microwave causes learning and memory impairment in rats

Sun L, Wang X, Ren K, Yao C, Wang H, Xu X, Wang H, Dong J, Zhang J, Yao B, Wei X, Peng R, Zhao L. Compound exposure of 2.8 GHz and 9.3 GHz microwave causes learning and memory impairment in rats. Heliyon. 2025 Jan 3;11(1):e41626. doi: 10.1016/j.heliyon.2025.e41626.

Abstract

The anxieties and concerns about health hazards caused by microwave has been growing recently. Previous studies have reported microwave induced structural and functional injuries to brain. However, the biological effects caused by compound microwave were largely unexplored. In this study, we investigated the biological effects on rat's hippocampus after sequentially exposure to 2.8 GHz and 9.3 GHz at **average power density of 10 mW/cm^2**. Morris water maze (MWM) navigation tests showed that microwave exposure significantly extended the average escape latency (AEL) at 1d and 3d after exposure, suggesting reduced learning and memory ability. Importantly, compound microwave produced strongest responses at 3 d after exposure. Moreover, microwave also could induce abnormal electroencephalogram (EEG), such as increasing the power of θ wave and δ wave, suggesting brain damage or dysfunction. Histopathological analysis suggested that microwave resulted in obvious structural injuries in hippocampus at 7 d after exposure, and most serious injuries were observed in compound microwave exposed rats. Moreover, disorder of mitochondria and reduced Nissl bodies in hippocampus might contribute to the decrease of cognitive function. However, both the cognitive function and hippocampal structure restored to normal at 28 d after exposure, which might be attributed to self-recovery mechanisms. Gene ontology (GO) and Protein-protein interaction (PPI) analyses of differential expressed genes (DEGs) in peripheral blood suggested that Htt and Bdnf might be potential indicators to predict microwave. In conclusion, compound microwave of 2.8 GHz and 9.3 GHz could elicit reversible structural injuries to hippocampus, which could decrease the cognitive function of rats.

Conclusion

The composite microwave of 2.8 GHz and 9.3 GHz can cause damage to rat hippocampal tissue, inhibit brain activity, and reduce the metabolism of hippocampal neurons through inducing mitochondrial disorder and decreasing Nissl bodies, and ultimately impaired the learning and memory abilities of rats. Notably, compound microwave can reduce the learning and memory ability of rats more than single frequency microwave. Through a combination of GO and PPI analyses, we have also identified Htt, Htra, Psen1, Tp73, and Bdnf as potential predictive indicators of microwave-induced neuronal damage in the nervous system.

Open access paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC11760294/

--

Measurement of non-ionizing radiation (NIR) from Wi-Fi access points and microwave ovens in residential environments

Silva, J. da L. A., Rodrigues, M. E. C., Pinheiro, F. S. R., Silva, G. S. da, Mendonça, H. B., Silva, R. Q. de F. H., … Sousa, V. A. de, Jr. (2025). Measurement of non-ionizing radiation (NIR) from Wi-Fi access points and

microwave ovens in residential environments. *Journal of Electromagnetic Waves and Applications*, 1–22. doi: 10.1080/09205071.2024.2449120.

Abstract

It is essential to monitor the radiation levels to which the general population may be exposed and compare them to limits established by national and international standards. This is especially true in residential environments, where there is a consolidated use of microwave ovens and Wi-Fi access points, in addition to the short distances devices can be from the user, as well as prolonged periods of user exposure. The objective of this work is to measure the exposure levels for daily Wi-Fi and microwave oven use scenarios in residential environments, to compare measured levels with the standards and to verify possible relation with the characteristics of the equipment under analysis. We use the Narda NBM-520 to support our measurements, the same equipment the National Telecommunication Agency (ANATEL) uses for Brazil's Non-ionizing Radiation (NIR) inspection. This work presents measurements in 40 Wi-Fi access points in which the highest exposure level was 4.66V/m (7.64 % of the limit), verified for the 2.4 GHz network (the so-called Situation 1). This work also presents measurements in 50 microwave ovens. It was observed that 72% of them had the highest level of exposure for the position in front of the door.

Conclusion

This work evaluates the level of exposure to NIR in homes, mainly from technologies commonly found in residences: Wi-Fi and microwave ovens. For this, measurements were performed on 40 Wi-Fi access points and 50 microwave ovens following measurement methodologies from suitable governmental agencies.

For Wi-Fi case study, the influence of electric field levels was investigated in four measurement situations, by equipment brand, and by time of use. The highest level of exposure was verified for the 2.4 GHz loaded network (Situation 1), at a value of 4.66V/m (7.64% of limit). As for the influence of the brand, the variation of the equipment model between the samples impacts the dispersion of the data; so Brand 4, which has the highest dispersion, has four different models and only two samples of identical models, while Brands 1 and 2 had the lowest dispersion and contain most of the samples belonging to the same model. Thus, no clear relation was observed regarding the year of acquisition of Wi-FI APs and the NIR leakage.

The microwave oven case study targets the relation between power density level and position, brand, time of use, and conservation. We observe that 72% of the samples had the highest level of exposure in front of the door, followed by 18% on the left side (considering the observer facing the microwave oven as a reference). We speculate that the ventilation outlet located at the left side is responsible for some NIR leakage, since 2 of 3 samples that presented the worst point at the top of the oven, the ventilation outlet is located in this local

Our results do not show the influence of the brand on the power density levels. However, regarding the time of use, there is a tendency for microwave ovens to reach higher levels of power density with age, since microwave ovens with year of acquisition <2016 (oldest) present the highest levels. Likewise, the state of conservation of the equipment restrained the levels of power density so that all microwave

ovens that reached an average power density above 0.6W/m2 showed signs of use. In addition, only two of the nine ovens that obtained peak power above 1W/m2 did not show signs of use.

https://www.tandfonline.com/doi/full/10.1080/09205071.2024.2449120

--

Evaluation of Population Exposure to Radiofrequency Fields in Microenvironments in the Cities of Lima, Cusco and Pucallpa in Perú, Using a Personal Exposimeter

Cruz V, Quispe M. Evaluation of Population Exposure to Radiofrequency Fields in Microenvironments in the Cities of Lima, Cusco and Pucallpa in Perú, Using a Personal Exposimeter. 2024 IEEE XXXI International Conference on Electronics, Electrical Engineering and Computing (INTERCON), Lima, Peru, 2024, pp. 1-8, doi: 10.1109/INTERCON63140.2024.10833492.

Abstract

The objective of this study was to make an evaluation of radiofrequency fields in microenvironments of Cuzco, Lima and Pucallpa using personal exposimeters. To carried it out a literature review was first made, then the urban environ-ments were defined, and 95 microenvironments were selected in the three cities. Subsequently, the exposimeter including the proprietary software for data processing was tested. In total, 1900 measurements of the main telecommunications services were carried out. The electric field was measured for each environment, with a maximum of 2.531 V/m, a minimum of 0.063 V/m and an average of 0.521 V/m for all measurements combined. Based on the ICNIRP 1998 limits, exposure quotients were also obtained for all the measurements combined and for the measurements in each city considering outdoor and indoor measurements, different types of environments and by main frequency bands. For all measurements combined, the maximum exposure quotient was 0.747% and the average was 0.033%. All measurements made were well below international limits, the largest contributors to total exposure were mobile phone base stations, the second largest were broadcasting services, and mobile phone handsets exposure was well below that of mobile phone base stations.

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=10833492&isnumber=10833229

--

Wireless radiation and health: making the case for proteomics research of individual sensitivity

Leszczynski D. Wireless radiation and health: making the case for proteomics research of individual sensitivity. Front. Public Health. Volume 12 doi: 10.3389/fpubh.2024.1543818.

No abstract

Excerpts

"According to the WHO definition of health, just a belief in having EHS and experiencing non-specific symptoms, physiological and/or psychological, is experiencing the health effects of wireless technology. Hence, it is correct to claim that wireless radiation causes health effects."

"Puzzlingly, the frequent observation that the self-declared EHS person can't feel the wireless radiation and can't recognize when the wireless transmitter emits radiation and when it is not transmitting, is considered ultimate proof that the form of individual sensitivity to wireless radiation called EHS is not caused by wireless radiation exposures. This is questionable as no person, sensitive or not, could feel the ionizing radiation or other non-ionizing radiation like ultraviolet in their environment."

"...logically and per analogiam with other environmental factors, individual sensitivity to wireless radiation, which includes EHS, exists as indicated below, and should be studied using biochemical methods."

"Search for sensitive individuals, most commonly using provocation studies where experimentally controlled exposures are followed by inquiries about acutely occurring symptoms and feelings, has failed to detect any sensitivity to wireless radiation. The reason might be that provocation exposures combined with psychological inquiries might be not sensitive enough to detect individual sensitivity to a single agent present in a mix of other environmental agents..."

"There is a need for human volunteer studies where the already proposed, and other potentially useful biomarkers, would be examined in groups of sensitive and non-sensitive persons, ethically exposed to wireless radiation."

"The way forward in EHS research is to discover biomarkers of EHS, molecules that are affected by wireless radiation exposure, by research using high-throughput screening techniques of proteomics, transcriptomics, and metabolomics [...]. For the start, proteomics might be the most promising of these methods."

"The reasons why proteomics is not used to study the physiological effects of wireless radiation exposures in humans are difficult to understand and comprehend. Despite the advantages of research using proteomics methodology, over the last 20 years, only a few proteomics studies have examined proteome changes in response to wireless radiation exposures."

"In conclusion, it is logical to conclude that the individual sensitivity to wireless radiation emitted by wireless communication devices and networks exists and impacts the health of sensitive persons. Clearly, the to-date unsuccessfully used methods of provocation studies were either too crude or too much affected by the perceptions and preexisting opinions of study volunteers."

Open access paper: https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2024.1543818/full

--

Amelioration and Immuno-modulation by Ashwagandha on Wi-fi Induced Oxidative Stress in Regulating Reproduction Via Estrogen Receptor Alpha in Male Japanese Quail

Gupta V, Srivastava R. Amelioration and Immuno-modulation by Ashwagandha on Wi-fi Induced Oxidative Stress in Regulating Reproduction Via Estrogen Receptor Alpha in Male Japanese Quail. Reprod Sci. 2025 Jan 13. doi: 10.1007/s43032-024-01774-7.

Abstract

As global change threatens avian biodiversity, understanding species responses to environmental perturbations due to radiation emitted by enormous increase in the application of wireless communication is very urgent. The study investigates the effect of MW radiation on redox balance, stress level, male fertility and the efficacy of Withania somnifera (WS) root extract (100 mg/kg body weight) orally administered in 8 weeks old mature male Japanese quail exposed to 2.4 GHz MW radiation for 2 h/day for 30 days with power density = 0.1264 mw/cm^2 and SAR = 0.9978 W/Kg. Wi-fi exposure induces a decrease in testicular weight, volume, density and gonado-somatic index (GSI) while Ashwagandha increases them. Oxidative stress parameters increased and activity of SOD, catalase, GSH was reduced in testes of exposed quail while Ashwagandha treatment reinstates the redox balance. Exposure to Wi-fi alters quail reproduction by increase in corticosterone and decreased testosterone with reduced expression of estrogen receptor alpha (ERα) in testis. Wi-fi exposure increases IL1β and reduces IL10 in testis. IL-1β inhibits testicular cell function and promotes apoptosis by increasing NF-κB and decreasing sperm count in exposed quails. Ashwagandha increases expression of ERα, sperm count and immunity in quail testis. Further, decrease in IL1β, NF-κB and increase in IL-10 after administration of Ashwagandha in Wi-fi exposed quail prevents inflammatory damages and enhances gonadal function. Thus, exposure to Wi-fi increases oxidative stress, activates apoptosis, modulates immunity in testis while Ashwagandha reverses them via enhanced ERα expression, increase in sperm count thereby enhancing fertility in male Japanese quail.

https://pubmed.ncbi.nlm.nih.gov/39806168/

--

The Development of a Reverberation Chamber for the Assessment of Biological Effects of Electromagnetic Energy Absorption in Mice

Iskra S, McIntosh RL, McKenzie RJ, Frankland JV, Deng C, Sylvester E, Wood AW, Croft RJ. The Development of a Reverberation Chamber for the Assessment of Biological Effects of Electromagnetic Energy Absorption in Mice. Bioelectromagnetics. 2025 Jan;46(1):e22539. doi: 10.1002/bem.22539.

Abstract

In this paper, we present the design, RF-EMF performance, and a comprehensive uncertainty analysis of the reverberation chamber (RC) exposure systems that have been developed for the use of researchers at the University of Wollongong Bioelectromagnetics Laboratory, Australia, for the purpose of investigating the biological effects of RF-EMF in rodents. Initial studies, at 1950 MHz, have focused on investigating thermophysiological effects of RF exposure, and replication studies related to RF-EMF exposure and progression of Alzheimer's disease (AD) in mice predisposed to AD. The RC exposure system was chosen as it allows relatively unconstrained movement of animals during exposures which can have the beneficial effect of minimizing stress-related, non-RF-induced biological and behavioral changes in the animals. The performance of the RCs was evaluated in terms of the uniformity of the Whole-Body Average-Specific Absorption Rate (WBA-SAR) in mice for a given RF input power level. The expanded uncertainty in WBA-SAR estimates was found to be 3.89 dB. Validation of WBA-SAR estimates based on a selected number of temperature measurements in phantom mice found that the maximum ratio of the temperature-derived WBA-SAR to the computed WBA-SAR was 1.1 dB, suggesting that actual WBA-SAR is likely to be well within the expanded uncertainties.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.22539

--

Continuous exposure to 60 Hz extremely low frequency magnetic field at 10-14 mT promotes various human cell proliferation by activating extracellular-signal-regulated kinase

Goh J, Suh D, Um DY, Chae SA, Park GS, Song K. Continuous exposure to 60 Hz extremely low frequency magnetic field at 10-14 mT promotes various human cell proliferation by activating extracellular-signal-regulated kinase. Biochem Biophys Res Commun. 2025 Jan 28;751:151414. doi: 10.1016/j.bbrc.2025.151414.

Abstract

We previously showed that 60 Hz extremely low-frequency magnetic fields (ELF-MF) at 6 mT promote various humancell proliferation. This study investigated the effects of 60 Hz ELF-MF at 10-16 mT on various mammalian cells, including human cervical carcinoma, rat neuroblastoma, liver cancer stem cells, immortalized normal hepatic cells, and normal fibroblasts. Using a revised ELF-MF-generating device that increases magnetic flux density stably without thermal effects, we exposed cells to 10 and 16 mT ELF-MF for 72 h. All cell types exhibited an approximately 20 % or greater increase in proliferation compared to the sham exposure group at 14 mT, with no further increase observed at 16 mT. In cells with activated proliferation at 14 mT, we observed activation of the MEK-ERK pathway and NF-κB, but not Akt, and a slight increase in S phase population. Intracellular and mitochondrial ROS levels remained unchanged, and the proliferation-activating effects persisted when oxidative phosphorylation was interrupted. No changes in intracellular calcium levels were observed, and the proliferation-activating effects were maintained in the presence of a calcium chelator. These findings suggest that ROS and intracellular calcium do not mediate ELF-MF's proliferation-activating physiological effect. In conclusion, exposure to 60 Hz ELF-MF at 10-14 mT promotes cell proliferation by activating ERK1/2 without affecting intracellular ROS and calcium levels.

https://pubmed.ncbi.nlm.nih.gov/39903967/

--

An amplification mechanism for weak ELF magnetic fields quantum-bio effects in cancer cells

Zandieh A, Shariatpanahi SP, Ravassipour AA, Azadipour J, Nezamtaheri MS, Habibi-Kelishomi Z, Ghanizadeh M, Same-Majandeh A, Majidzadeh-A K, Taheri A, Ansari AM, Javidi MA, Pirnia MM, Goliaei B. An amplification mechanism for weak ELF magnetic fields quantum-bio effects in cancer cells. Sci Rep. 2025 Jan 23;15(1):2964. doi: 10.1038/s41598-025-87235-w.

Abstract

Observing quantum mechanical characteristics in biological processes is a surprising and important discovery. One example, which is gaining more experimental evidence and practical applications, is the effect of weak magnetic fields with extremely low frequencies on cells, especially cancerous ones. In this study, we use a mathematical model of ROS dynamics in cancer cells to show how ROS oscillatory patterns can act as a resonator to amplify the small effects of the magnetic fields on the radical pair dynamics in mitochondrial Complex III. We suggest such a resonator can act in two modes for distinct

states in cancer cells: (1) cells at the edge of mitochondrial oscillation and (2) cells with local oscillatory patches. When exposed to magnetic fields, the first group exhibits high-amplitude oscillations, while the second group synchronizes to reach a whole-cell oscillation. Both types of amplification are frequency-dependent in the range of hertz and sub-hertz. We use UV radiation as a positive control to observe the two states of cells in DU and HELA cell lines. Application of magnetic fields shows frequency-dependent results on both the ROS and mitochondrial potential which agree with the model for both type of cells. We also observe the oscillatory behavior in the time-lapse fluorescence microscopy for 0.02 and 0.04 Hz magnetic fields. Finally, we investigate the dependence of the results on the field strength and propose a quantum spin-forbidden mechanism for the effect of magnetic fields on superoxide production in Q_O site of mitochondrial Complex III.

Excerpts

In a nutshell, our proposed scheme acts in two distinct levels: [I] At the level of superoxide production in mitochondrial ETC where the applied magnetic field can alter production rate of superoxide by closing the energy gap between singlet and triplet state of a radical pair in Complex III. Here the applied magnetic field effectively acts as a static field. And [II] at the level of mitochondrial network, where oscillation of the applied magnetic field (and thus oscillation in the production rate of superoxide) can resonate with the intrinsic frequency of RIRR in cancer cells. At this level, the minor effect of underlying RPM is amplified to affect the physiology of the cell in a frequency-dependent manner....

Conclusion

Our study proposed a quantum biological mechanism for the effect of varying small magnetic fields on cancer cells. The frequency dependent effect of magnetic field is here postulated to be the result of interaction of alternating field with the intrinsically oscillatory system of coupled mitochondria via a novel radical pair phenomenon.

Here the site of action of the field is suggested to be the semiquinone/FeS radical pair in Complex III of ETC. A magnetic field of the order of few hundreds of millitesla is predicted to populate the triplet state of the radical pair whose electron transfer onto oxygen molecule is restricted by a spin forbidden reaction in half of the cases. This leads to a mild decrease in superoxide production in the presence of such a magnetic field. An oscillatory field can amplify such an effect via a resonance phenomenon in a network of coupled mitochondria in many cell lines. This substantial change in ROS balance can potentially cause physiological effect in the cells, including apoptosis. Our simulations show the mentioned effect can be manifested in two resonance modes: Either triggering the oscillation in network or synchronizing the out of phase oscillatory mitochondria. Our experimental results confirm the suggested model which can explain the previous observations for the first time. The hypothesis was validated by the observation of the synchronization when cells were exposed to sudden UV irradiation.

Moreover, we observed a frequency-dependent variation of the ROS concentration and mitochondrial membrane potential under the effect of oscillatory magnetic field in frequencies and intensities suggested by our model.

Further in vitro investigation with higher resolution of microscopy and more extensive profile of magnetic field frequencies and intensities can shed light on the detailed components of the involved mechanism.

Open access paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC11757740/

--

Effect of terahertz radiation on cells and cellular structures (Review)

Rytik AP, Tuchin VV. Effect of terahertz radiation on cells and cellular structures. Front Optoelectron. 2025 Jan 27;18(1):2. doi: 10.1007/s12200-024-00146-y.

Abstract

The paper presents the results of modern research on the effects of electromagnetic terahertz radiation in the frequency range 0.5-100 THz at different levels of power density and exposure time on the viability of normal and cancer cells. As an accompanying tool for monitoring the effect of radiation on biological cells and tissues, spectroscopic research methods in the terahertz frequency range are described, and attention is focused on the possibility of using the spectra of interstitial water as a marker of pathological processes. The problem of the safety of terahertz radiation for the human body from the point of view of its effect on the structures and systems of biological cells is also considered.

Conclusion

The presented data show that THz radiation has a variety of effects on cells, which are manifested in the disruption of the properties of cell membranes, the formation of pores, the activation of ion channels, and changes in their proliferation and viability [110]. Possible mechanisms that determine the reaction of cells to THz radiation may be the following:

- a change in the conformation of membrane proteins, which triggers an intracellular cascade of regulators of the genetic and enzymatic systems and the permeability of cell membranes for various substances;
- a change in the conformation of membrane proteins that perceive external regulatory signals;
- change in the conformation of membrane proteins that are pumps or channels for the transport of various substances into and out of the cell;
- redistribution of the electric charge on the cell membrane;
- excitation of resonant oscillations of macromolecules that make up the cell membrane and the cytoskeleton as a whole.

Thus, fundamentally, THz radiation does not cause the breaking or restoration of chemical covalent bonds, since the quantum energy is insufficient for this, 1 THz − 4.1 meV. However, this radiation, in its frequencies, falls into the region of vibrational–rotational movements of biological molecules and water and can excite energy levels of vibrational–rotational transitions of proteins and water, and thereby change the spatial conformation of proteins, which can affect various interactions between proteins, protein and water molecules.

It is generally accepted that there are several mechanisms that determine the effect of the response of living cells to THz radiation, in particular [1]:

- redistribution of electrical charge on the cell membrane, changing the ratio of concentrations of bound and free water;
- excitation of resonant vibrations of macromolecules that make up the cell membrane and the cytoskeleton as a whole;
- change in the conformation of membrane proteins.

Water molecules themselves can be considered as a universal marker in the THz frequency range, which is sensitive to various vital processes occurring in living tissues and cells. Compared to what is traditionally described in dielectric spectroscopy, in the THz frequency range water as a marker allows one to obtain new information about biological systems. Moving from the gigahertz (GHz) to the THz range, we are gradually approaching various vibrational–rotational processes that are determined by the interaction of water molecules with surrounding molecular systems [23]. Biomacromolecules, being excited, absorb part or all of the energy of electromagnetic waves, depending on the frequency of the incident radiation [26]. Since the generalized terahertz range (0.1–100 THz) partially overlaps with the vibration spectrum of biomolecules, terahertz waves can greatly enhance vibrations of biomolecule bonds such as twisting, stretching, and bending through resonant excitation [111]. However, early studies of the biological effects induced by optical stimulation focused on the infrared region, which promotes the strong absorption of incoming energy by water and its conversion into heat [112, 113]. While heat alters transmembrane capacitance or ion channel activity and hence induces biological responses, it inevitably also causes a transient increase in local temperature. On the other hand, terahertz wave modulation is seen as a promising approach for interfering with biophysical processes without being damaged by electromagnetic radiation. The study of non-thermal biological effects of infrared radiation has attracted close attention from both opticians and biologists. In addition, THz waves with low photon energy are unlikely to cause ionizing effects, thus will not damage genome integrity as other radiation intervention approaches might [112].

It can be concluded that currently there is no full consensus in the scientific community as to whether THz radiation has a damaging effect on biological objects at various levels of organization [83, 114]. Therefore, an increase in studies using THz radiation to monitor the activity of uncontrolled dividing cells is expected in the near future. The development of new high-resolution THz diagnostic methods in combination with AI technologies will take cancer diagnosis and therapy to a new level. It is obvious that more and more new data will appear soon for THz diagnostics and therapy of tumor oncological processes. In addition, theranostics technologies, where THz radiation from the same source is used first for diagnosis and then at increased energy parameters for therapy within a single protocol, have not yet received proper development, but are undoubtedly promising.

Open access paper: https://link.springer.com/article/10.1007/s12200-024-00146-y

--

Metabolomic and gene networks approaches reveal the role of mitochondrial membrane proteins in response of human melanoma cells to THz radiation

Butikova EA, Basov NV, Rogachev AD, Gaisler EV, Ivanisenko VA, Demenkov PS, Makarova AA, Ivanisenko TV, Razumov IA, Kolomeyets DA, Cheresiz SV, Solovieva OI, Larionov KP, Sotnikova YS, Patrushev YV,

Kolchanov NA, Pokrovsky AG, Vinokurov NA, Kanygin VV, Popik VM, Shevchenko OA. Metabolomic and gene networks approaches reveal the role of mitochondrial membrane proteins in response of human melanoma cells to THz radiation. Biochim Biophys Acta Mol Cell Biol Lipids. 2025 Jan 20:159595. doi: 10.1016/j.bbalip.2025.159595.

This paper presents a study on the biological effects of terahertz irradiation with a frequency of 2.3 THz and a power of 0.056 W/cm^2 on SK-MEL-28 cells, using targeted metabolomic screening by liquid chromatography and tandem mass spectrometry (LC-MS/MS) and gene network analysis.

Highlights

• THz radiation exposure induced significant changes in the levels of 40 metabolites, especially those related to purine and pyrimidine pathways. Lipids, such as ceramides and phosphatidylcholines, were also affected by this exposure.
• Gene network reconstruction identified mitochondrial membrane proteins, including components of the respiratory chain complex, ATP synthase, and lipid rafts, as key regulators of the enzymes involved in biosynthesis and degradation of the significantly altered metabolites.
• THz radiation induces reversible disruption of lipid raft macromolecular structure, altering mitochondrial molecule transport while maintaining protein integrity.
• Importantly, the metabolic effects observed were specific to THz radiation and were not due to heating, as evidenced by the distinct responses in control cells exposed to infrared radiation.

Abstract

Terahertz (THz) radiation has gained attention due to technological advancements, but its biological effects remain unclear. We investigated the impact of 2.3 THz radiation on SK-MEL-28 cells using metabolomic and gene network analysis. Forty metabolites, primarily related to purine, pyrimidine synthesis and breakdown pathways, were significantly altered post-irradiation. Lipids, such as ceramides and phosphatidylcholines, were also affected. Gene network reconstruction and analysis identified key regulators of the enzymes involved in biosynthesis and degradation of significantly altered metabolites. Mitochondrial membrane components, such as the respiratory chain complex, the proton-transporting ATP synthase complex, and components of lipid rafts, reacted to THz radiation. We propose that THz radiation induces reversible disruption of the lipid raft macromolecular structure, thereby altering mitochondrial molecule transport while maintaining protein integrity, which explains the high cell survival rate. Our findings enhance the understanding of THz biological effects and emphasize the role of membrane components in the cellular response to THz radiation.

https://www.sciencedirect.com/science/article/abs/pii/S1388198125000034

--

Hidden route of protein damage through oxygen-confined photooxidation

Kim S, Kim E, Park M, Kim SH, Kim BG, Na S, Sadongo VW, Wijesinghe WCB, Eom YG, Yoon G, Jeong H, Hwang E, Lee C, Myung K, Kim CU, Choi JM, Min SK, Kwon TH, Min D. Hidden route of protein damage through oxygen-confined photooxidation. Nat Commun. 2024 Dec 30;15(1):10873. doi: 10.1038/s41467-024-55168-z.

Abstract

Oxidative modifications can disrupt protein folds and functions, and are strongly associated with human aging and diseases. Conventional oxidation pathways typically involve the free diffusion of reactive oxygen species (ROS), which primarily attack the protein surface. Yet, it remains unclear whether and how internal protein folds capable of trapping oxygen (O_2) contribute to oxidative damage. Here, we report a hidden pathway of protein damage, which we refer to as O_2-confined photooxidation. In this process, O_2 is captured in protein cavities and subsequently converted into multiple ROS, primarily mediated by tryptophan residues under blue light irradiation. The generated ROS then attack the protein interior through constrained diffusion, causing protein damage. The effects of this photooxidative reaction appear to be extensive, impacting a wide range of cellular proteins, as supported by whole-cell proteomic analysis. This photooxidative mechanism may represent a latent oxidation pathway in human tissues directly exposed to visible light, such as skin and eyes.

Open access paper: https://www.nature.com/articles/s41467-024-55168-z

--

Structural Analysis of Plasma-Induced Oxidation and Electric Field Effect on the Heat Shock Protein (Hsp60) Structure

Attri P, Okumura T, Koga K, Shiratani M. Structural Analysis of Plasma-Induced Oxidation and Electric Field Effect on the Heat Shock Protein (Hsp60) Structure: A Computational Viewpoint. Chem Biodivers. 2025 Jan 5:e202401243. doi: 10.1002/cbdv.202401243.

Abstract

In recent years, there has been an increase in the study of the mechanisms behind plasma oncology. For this, many wet lab experiments and computational studies were conducted. Computational studies give an advantage in examining protein structures that are costly to extract in enough amounts to analyze the biophysical properties following plasma treatment. Therefore, in this work, we studied the effect of plasma oxidation and electric field on the human mitochondrial heat shock protein (mHsp60). Hsp60, alias chaperonin, is one of the most conserved proteins expressed across all species. Hence, we performed molecular dynamic simulations to calculate the root-mean-square deviation, root-mean-square fluctuation, and solvent-accessible surface area of mHsp60 with and without oxidation. In addition to the oxidation state, we also applied an electric field (0.003 and 2.0 V/nm) to check the changes in the mHsp60 protein. Through simulations, we observed that the electric field strongly affects the structure of mHsp60 protein compared with the oxidation. The combination of oxidation and electric field effect increases the destabilization of the mHsp60 structure compared with their respective control states.

https://pubmed.ncbi.nlm.nih.gov/39756028/

Conclusion

We can conclude that the mHsp60 structure is modified by possible plasma-assisted oxidation, particularly involving Trp, Try, and Met amino acids, resulting in noticeable structural changes. RMSD

values reflect that mHsp60 structure flexibility slightly increases in OXID-2, whereas the structure becomes rigid in OXID-1. This suggests that the oxidation of Met amino acids plays a significant role in enhancing mHsp60 flexibility. On the other hand, applying an electric field (EF1 and EF2) to the control mHsp60 increases the RMSD value more than for oxidized mHsp60 (OXID-1 and OXID-2), indicating that the electric field has a more substantial effect on the mHsp60 structure than oxida- tion alone. Additionally, the RMSD values of OXID-1 and OXID-2 increased after electric field treatment (0.003 V/nm from the x, y, and z axes), demonstrating that the small electric field generated by DBD plasma can significantly influence protein conforma- tional changes in both native and oxidized states. HSPs, including Hsp60, play a vital role in maintaining protein homeostasis, which is crucial for cell integrity, survival, and metabolism. However, when chaperone-assisted protein quality control is compromised due to oxidation or exposure to electric effect, it may trigger the onset and progression of numerous diseases.

--

World Health Organization's EMF Project's Systemic Reviews on the Association Between RF Exposure and Health Effects Encounter Challenges

In a newly-published paper, one of the world's most renowned scientists who has studied the effects of radio frequency (RF) radiation, Dr. James C. Lin, Professor Emeritus at the University of Illinois, Chicago and a former ICNIRP Commissioner, attacks the World Health Organization's systematic reviews of the research on RF radiation that dismiss the substantial evidence for adverse biological and health effects.

He concludes:

"The criticisms and challenges encountered by the published WHO-EMF systematic reviews are brutal, including calls for retraction. Rigorous examinations of the reviews reveal major concerns. In addition to the scientific quality, they appear to have a strong conviction of nothing but heat to worry about with RF radiation. The unsubtle message that cellular mobile phones do not pose a cancer risk is clear. The reviews exhibit a lack of serious concerns for conflicts of interest and display unequivocal support for the recently promulgated ICNIRP RF exposure guidelines for human safety.

From its inception, WHO-EMF had close ties with ICNIRP, a private organization, frequently referred to as the WHO-EMF project's scientific secretariat [18]. What may not be as apparent for the WHO-EMF systematic reviews is the lack of diversity of views. A large number of ICNIRP commissioners and committee members are listed as authors for the WHO-EMF systematic reviews; some also served as lead authors. These concerns advance issues of reviewer independence and potential for conflicts of interest."

Lin JC. World Health Organization's EMF Project's Systemic Reviews on the Association Between RF Exposure and Health Effects Encounter Challenges [Health Matters]. *IEEE Microwave Magazine*, 26(1): 13-15, Jan. 2025, doi: 10.1109/MMM.2024.3476748.

No abstract

Excerpts

1) Regarding "The WHO-EMF systematic review on the association between RF exposure and adverse health effects pertaining to reproductive health (pregnancy and birth outcome.... While the WHO-EMF

systematic review presents itself as thorough, scientific, and relevant to human health, numerous issues were identified, suggesting the WHO-EMF review was severely flawed. The found flaws skewed the results in support of the review's conclusion that there is no conclusive evidence for effects other than RF-induced tissue heating. It showed that the underlying data, when relevant studies are cited correctly, support the opposite conclusion: "There are clear indications of detrimental nonthermal effects" from RF exposure. The authors identified a multitude of flaws in the methodology. To those scientists, the methodology and low quality of the systematic review were highly concerning "as it threatens to undermine the trustworthiness and professionalism of the WHO-EMF project in the area of human health hazards from man-made RF radiation."

2) Regarding "The WHO-EMF systematic review of human observational studies on the occurrence of migraine, headaches, tinnitus, sleep disturbances, and nonspecific symptoms in the general and working population .... An ensuing critical appraisal by three accomplished senior researchers documented major problems with the WHO-EMF-commissioned review and called for its retraction [8]. The meta-analysis for the handful of very heterogeneous primary studies identified for each of the analyzed exposure and outcome combinations appeared fundamentally inappropriate. The number is very small, and the methodological quality of the relevant primary studies is low. In contrast, this peer-reviewed publication concluded that the body of evidence reviewed is inadequate to either support or refute the safety of current exposure limits.

3) "Some skepticism has been expressed regarding a third WHO-EMF systematic review on RF-induced oxidative stress [9]. The study identified 11,599 studies on oxidative stress in the frequency range 800–2,450 MHz and then eliminated 11,543 of them as not meeting the criteria for inclusion. Of the remaining 56 papers, there were 45 animal studies and 11 in vitro cellular studies....For many years, Henry Lai, a leading researcher in RF oxidative responses and professor emeritus at the University of Washington, Seattle, has maintained a bibliography of RF-oxidative stress papers. As of mid-August, his list includes 367 studies, published between 1997 and 2024. By his count, 89% showed significant effects. Lai's assessment of the WHO-EMF review is that it left out a large portion of RF-oxidative effect studies and appears to have only considered oxidative molecular reactions among the possible oxidative effects [10]. As reported, others have opined that "this systematic review methodically excluded most of the relevant research.""

4) Regarding the WHO systematic review to assess the evidence provided by human epidemiological studies for the cancer risk from radiofrequency (RF): " This WHO-EMF review was picked up and reported on by many Western media outlets. Actually, there are truly few data that are new in this review. For sure, the assessment of scientific evidence in this subject has been controversial and less than uniform. The question is, "Is this review really the definitive word on the long-standing issue of whether cell phone radiations pose a cancer risk?" My answer is, far from it!"

"Microwave News [12] published a meticulously researched investigative report in the historical context of the latest WHO-EMF cancer review. Five years ago, the lead author [13] with some members of the same team made similar efforts to terminate the RF–cancer debate with basically the same no-risk message. However, "it was not well received" by the scientific community, since the analysis excluded some people older than 59 years of age, the largest segment of the brain cancer population."

"The WHO-IARC, NIH-NTP, and Ramazzini outcomes, under normal circumstances, would likely have provided the justification for raising WHO-IARC's current possible cancer risk designation to the probable cancer-causing classification, if not higher.

https://ieeexplore.ieee.org/document/10795296

--

A mechanistic understanding of human magnetoreception validates the phenomenon of electromagnetic hypersensitivity (EHS)

Henshaw DL, Philips A. (2024). A mechanistic understanding of human magnetoreception validates the phenomenon of electromagnetic hypersensitivity (EHS). *International Journal of Radiation Biology*, 1–19. doi: 10.1080/09553002.2024.2435329.

Abstract

Background Human electromagnetic hypersensitivity (EHS) or electrosensitivity (ES) symptoms in response to anthropogenic electromagnetic fields (EMFs) at levels below current international safety standards are generally considered to be nocebo effects by conventional medical science. In the wider field of magnetoreception in biology, our understanding of mechanisms and processes of magnetic field (MF) interactions is more advanced.

Methods We consulted a range of publication databases to identify the key advances in understanding of magnetoreception across the wide animal kingdom of life.

Results We examined primary MF/EMF sensing and subsequent coupling to the nervous system and the brain. Magnetite particles in our brains and other tissues can transduce MFs/EMFs, including at microwave frequencies. The radical pair mechanism (RPM) is accepted as the main basis of the magnetic compass in birds and other species, acting via cryptochrome protein molecules in the eye. In some cases, extraordinary sensitivity is observed, several thousand times below that of the geomagnetic field. Bird compass disorientation by radio frequency (RF) EMFs is known.

Conclusions Interdisciplinary research has established that all forms of life can respond to MFs. Research shows that human cryptochromes exhibit magnetosensitivity. Most existing provocation studies have failed to confirm EHS as an environmental illness. We attribute this to a fundamental lack of understanding of the mechanisms and processes involved, which have resulted in the design of inappropriate and inadequate tests. We conclude that future research into EHS needs a quantum mechanistic approach on the basis of existing biological knowledge of the magnetosensitivity of living organisms.

Conclusions and recommendations

1. At the scientific level, researchers working in the field of *magnetoreception in biology* should be made aware of EHS as a human public health concern and funded to address the issue as part of their scientific research.

2. All interested parties, especially EHS sufferers and medical professionals, should be made aware of the considerable growth in understanding in recent decades of the mechanisms by which all forms of life sense MFs/EMFs, even at extremely low levels. EHS research to date has been significantly hindered by a fundamental lack of knowledge among many medical scientists and EHS researchers regarding the current scientific understanding of quantum biology mechanisms and processes. This has resulted in the design and analysis of inappropriate provocation tests.

3. Almost all existing epidemiological and provocation studies have failed adequately to determine and measure the necessary dependent and independent variables. In particular:

    1. to characterize in proper technical detail the EMF/RF exposures (including electric and MF levels; average and peak power-density levels; frequencies involved; and modulation characteristics).

    2. to triage participants effectively to remove 'electrophobic' and other volunteers self-reporting apparent EHS-related problems.

    3. in provocation studies, to fail to recognize the nonlinear nature of EHS responses and the extremely low levels of exposure (<100 nT) that have effects and, instead, use relatively high exposures fairly close to the ICNIRP and IEEE guidance levels.

    4. in provocation studies, to provide a participant-comfortable extremely low EMF/RF test location, screened from anthropogenic sources and allow adequate time (days rather than hours) for adverse effects to washout between exposures.

4. EHS studies should move away from current, nonforensic epidemiological approaches and human subjective provocation studies (Leszczynski 2022; Röösli et al. 2024). Instead, objective measurements of biological parameters, such as heart rate variability, brain wave activity (e.g. fMRI and wide-bandwidth EEG), and the immune response to oxidative stress should be investigated (Caswell et al. 2016; Gurfinkel et al. 2018; Pishchalnikova et al. 2019; Wang et al. 2019; Thoradit et al. 2024). We caution that these approaches require sophisticated design and analysis and advanced design personal exposure meters.

5. We recommend that the WHO properly reevaluates its understanding of EHS to align it with the substantial body of available scientific literature showing mechanistic evidence of interactions of all forms of life, including humans, with low levels of electric and magnetic fields.

Open access paper: https://www.tandfonline.com/doi/full/10.1080/09553002.2024.2435329

--

Gaps in Knowledge Relevant to the "ICNIRP Guidelines for Limiting Exposure to Time-Varying Electric, Magnetic and Electromagnetic Fields (100 kHz TO 300 GHz)"

International Commission on Non-Ionizing Radiation Protection (ICNIRP). Gaps in Knowledge Relevant to the "ICNIRP Guidelines for Limiting Exposure to Time-Varying Electric, Magnetic and Electromagnetic Fields (100 kHz TO 300 GHz)". Health Phys. 2025 Feb 1;128(2):190-202. doi: 10.1097/HP.0000000000001944.

Abstract

In the last 30 y, observational as well as experimental studies have addressed possible health effects of exposure to radiofrequency electromagnetic fields (EMF) and investigated potential interaction mechanisms. The main goal of ICNIRP is to protect people and the environment from detrimental exposure to all forms of non-ionizing radiation (NIR), providing advice and guidance by developing and disseminating exposure guidelines based on the available scientific research on specific parts of the electromagnetic spectrum. During the development of International Commission on Non-Ionizing Radiation Protection's (ICNIRP's) 2020 radiofrequency EMF guidelines some gaps in the available data were identified. To encourage further research into knowledge gaps in research that would, if addressed, assist ICNIRP in further developing guidelines and setting revised recommendations on limiting exposure, data gaps that were identified during the development of the 2020 radiofrequency EMF guidelines, in conjunction with subsequent consideration of the literature, are described in this Statement. Note that this process and resultant recommendations were not intended to duplicate more traditional research agendas, whose focus is on extending knowledge in this area more generally but was tightly focused on identifying the highest data gap priorities for guidelines development more specifically. The result of this distinction is that the present data gap recommendations do not include some gaps in the literature that in principle could be relevant to radiofrequency EMF health, but which were excluded because either the link between exposure and endpoint, or the link between endpoint and health, was not supported sufficiently by the literature. The evaluation of these research areas identified the following data gaps: (1) Issues concerning relations between radiofrequency EMF exposure and heat-induced pain; (2) Clarification of the relation between whole-body exposure and core temperature rise from 100 kHz to 300 GHz, as a function of exposure duration and combined EMF exposures; (3) Adverse effect thresholds and thermal dosimetry for a range of ocular structures; (4) Pain thresholds for contact currents under a range of exposure scenarios, including associated dosimetry; and (5) A range of additional dosimetry studies to both support future research, and also to improve the application of radiofrequency EMF exposure restrictions in future guidelines.

Open access paper:https://journals.lww.com/health-physics/fulltext/2025/02000/gaps_in_knowledge_relevant_to_the__icnirp.8.aspx

--

Sensory Processing Sensitivity, and Not Gender, Drives Electromagnetic Hypersensitivity and Nature Connection

Watten RG, Volden F, Visnes H. Sensory Processing Sensitivity, and Not Gender, Drives Electromagnetic Hypersensitivity and Nature Connection. Ecopsychology. Nov 28, 2024. doi: 10.1089/eco.2024.0027

Abstract

Electromagnetic hypersensitivity (EHS) is a biomedical condition associated with exposure to man-made radio-frequency electromagnetic fields. Common sources are devices like mobile phones, wireless networks, base stations, computers, TVs, and more. Identifying individuals at risk of developing EHS is important. Those with heightened sensory processing sensitivity (SPS; the ability to perceive, process, and react to environmental stimuli, approximately 25% to 30% of the population) are of special interest. SPS could also be associated with connectedness to nature (CNS) and EHS. In the current gender-matched cross-sectional study (*n* = 450; 225 men and 225 women) we assessed gender differences in environmental sensitivity assessed as SPS, EHS, and CNS. Women had higher general EHS prevalence

than men (13.3% vs. 5.3%), higher mean values on perceived sensitivity for 5 out of 9 categories of electromagnetic equipment (computers, electrical appliances, fluorescent lighting, mobile phones, and television) and they had higher scores on the three EHS Scales EHS General, EHS Scale, and EHS Index, in addition to higher SPS and CNS scores. The gender differences vanished when adjusting for SPS in the multivariate analyses of variance (MANOVA) analyses, suggesting that the differences in these scales were associated with SPS. The statistical equation modeling (SEM) results showed a significant positive direct effect of SPS on CNS, and a significant indirect effect mediated by EHS. The present study provides the first empirical evidence that highly sensitive individuals are capable of perceiving electromagnetic radiation. The results indicate that perceived EHS and CNS are driven by SPS and not by gender.

https://www.liebertpub.com/doi/10.1089/eco.2024.0027

--

Effect of Exposure to Mobile Phones on Electrical Cardiac Measurements: Multivariate Analysis & Variable Selection Algorithm to Detect Relationship With Mean Changes

Alharbi N, Alassiri M. The Effect of Exposure to Mobile Phones on Electrical Cardiac Measurements: A Multivariate Analysis and a Variable Selection Algorithm to Detect the Relationship With Mean Changes. Int J Cell Biol. 2024 Oct 3;2024:7093771. doi: 10.1155/2024/7093771.

Abstract

Background: The exponential growth in mobile phone usage has raised concerns about electromagnetic field (EMF) exposure and its health risks. Blood pressure and BMI, which impair heart function due to decreased adrenoreceptor responsiveness, parasympathetic tone withdrawal, and increased sympathetic activity, may further exacerbate these risks. However, the effects of radiofrequency electromagnetic (RF-EM) exposure from mobile phones on electrocardiograms (ECGs) and heart rate variability (HRV) in individuals remain unclear.

Purpose: Building upon our previous findings on HRV changes due to mobile phone proximity, this study is aimed at significantly enhancing the analytical approach used to assess the effects of mobile phones on cardiac parameters. This study exploits data from a previous study but with a different purpose. The aim of this study is twofold: (a) to examine whether exposure to mobile phones changes the five variables (P-R, QRS, QT, ST, and HR) in a multivariate manner and (b) to examine whether the blood pressure and/or the body mass index (BMI), which acts as a proxy for obesity, have an effect on the change of these five variables. For both aspects of the study, four cycles are performed.

Method: We conducted multivariate analysis on previously collected electrical cardiac measurement data from 20 healthy male subjects exposed to mobile phone EMF, with the mobile phones placed at four different body locations. The one-sample Hotelling T^2 test on the mean vector of differences was utilised instead of multiple paired t-tests. This multivariate method comprehensively analyzes data features and accounts for variable correlations, unlike multiple univariate analyses. Given our small sample size, we employed the MMPC variable selection algorithm to identify predictor variables significantly related to mean changes.

Results: Significant alterations in ECG intervals and heart rate were noted in the subjects before and

after the first EMF exposure cycle, independent of their BMI. Notably, heart rate, P-R, and QRS intervals fell postexposure while QT and ST intervals increased. These changes were influenced by variations in systolic blood pressure, with BMI showing no significant effect.

Conclusion: The observed modifications in cardiac electrical measurements due to mobile phone EMF exposure are attributed to the effects of EMF itself, with no impact from BMI on the extent of these changes.

Note: iPhone 5 Plus was used.

Conclusion

Given the evidence linking EMFs from mobile phones to adverse effects on heart health, it is crucial for regular users of such devices, especially those at risk of CVDs, to take steps towards reducing their overall exposure. This can be achieved through measures such as limiting talk time or using hands-free headsets during calls. Additionally, ongoing research in this area is essential to deepen our understanding of how these potentially harmful frequencies impact our bodies over both short- and long-term periods.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1155/2024/7093771

--

The Role of Excess Charge Mitigation in Electromagnetic Hygiene: An Integrative review

Jamieson IA, Bell JNB, Holdstock P. The Role of Excess Charge Mitigation in Electromagnetic Hygiene: An Integrative review. Biomedical Journal. 2024, doi: 10.1016/j.bj.2024.100801.

Abstract

The electromagnetic characteristics of many environments have changed significantly in recent decades. This is in large part due to the increased presence of equipment that emits electromagnetic radiation and materials that may often readily gain excess charge. The presence of excess charge can often increase risk of infection from pathogens, and likelihood of individuals experiencing compromised performance, respiratory problems and other adverse health issues from increased uptake of particulate matter. It is proposed that adopting improved electromagnetic hygiene measures, including optimized humidity levels, to reduce the presence of inappropriate levels of electric charge can help reduce the likelihood of ill health, infection and poor performance arising from contaminant inhalation and deposition, plus reduce the likelihood of medical devices and other electronic devices getting damaged and/or having their data compromised. It is suggested that such measures should be more widely adopted within clinical practice guidelines and water, sanitation and hygiene programs.

Highlights

Electric fields can increase localized deposition of pathogens.

High charge of either polarity increases contaminant deposition.

40-60% relative humidity reduces likelihood of infection.

Proper specification of materials reduces infection risks.

Electromagnetic hygiene can reduce risk of infection.

Open access paper: https://www.sciencedirect.com/science/article/pii/S2319417024001045

--

5G RF-EMF Effects on the Human Sleep Electroencephalogram: A Randomized Controlled Study in Healthy Volunteers (pre-print, not peer-reviewed)

My note: Most studies to date that claim to evaluate the effects of 5G exposure did not employ a 5G signal generator. This study, however, used an exposure system (sXh5G) developed by the IT'IS Foundation that provides a controlled and well-characterized 5G EMF exposure at two different carrier frequencies.

Sousouri G, Eicher C, D'Angelo RM, Billecocq M, Fussinger T, Studler M, Capstick M, Kuster N, Achermann P, Huber R, Landolt H-P. 5G Radio-Frequency-Electromagnetic-Field Effects on the Human Sleep Electroencephalogram: A Randomized Controlled Study in CACNA1C Genotyped Healthy Volunteers. MedRxiv. Dec 26, 2024. doi: 10.1101/2024.12.16.24319082.

Abstract

Background: The introduction of 5G technology as the latest standard in mobile telecommunications has raised concerns about its potential health effects. Prior studies of earlier generations of radiofrequency electromagnetic fields (RF-EMF) demonstrated narrowband spectral increases in the electroencephalographic (EEG) spindle frequency range (11-16 Hz) in non-rapid-eye-movement (NREM) sleep. However, the impact of 5G RF-EMF on sleep remains unexplored. Additionally, RF-EMF can activate L-type voltage-gated calcium channels (LTCC), which have been linked to sleep quality and EEG oscillatory activity.

Objective: This study investigates whether the allelic variant rs7304986 in the CACNA1C gene, encoding the α1C subunit of LTCC, modulates 5G RF-EMF effects on EEG spindle activity during NREM sleep.

Methods: Thirty-four healthy, matched participants, genotyped for rs7304986 (15 T/C and 19 T/T carriers), underwent a double-blind, sham-controlled study with standardized left-hemisphere exposure to two 5G RF-EMF signals (3.6 GHz and 700 MHz) for 30 min before sleep. Sleep spindle activity was analyzed using high-density EEG and the Fitting Oscillations & One Over f (FOOOF) algorithm.

Results: T/C carriers reported longer sleep latency compared to T/T carriers. A significant interaction between RF-EMF exposure and rs7304986 genotype was observed, with 3.6 GHz exposure in T/C carriers inducing a faster spindle center frequency in the central, parietal, and occipital cortex compared to sham.

Conclusion: These findings suggest 3.6 GHz 5G RF-EMF modulates spindle center frequency during NREM sleep in a CACNA1C genotype-dependent manner, implicating LTCC in the physiological response to RF-EMF and underscoring the need for further research into 5G effects on brain health.

Excerpts

All participants completed three experimental nights with different, standardized exposure conditions according to a randomized, double-blind, cross-over design: 1) 30-min, pre-sleep exposure to an active

5G EMF at a carrier frequency of 700 MHz, 20 MHz bandwidth, and 12.5 Hz applied power control, 2) 30-min, pre-sleep exposure to an active 5G EMF at a carrier frequency of 3.6 GHz, 100 MHz bandwidth, and 12.5 Hz applied power control, and 3) a 30-min sham exposure without an active field....

All exposure conditions were administered with the same exposure system (sXh5G), provided by the IT'IS Foundation for Research on Information Technologies in Society (IT'IS Foundation, Zurich, Switzerland), which ensures controlled and well-characterized 5G EMF exposure. Following detailed simulated dosimetry (Figure 1), the signal intensity was calibrated to ensure the specific absorption rate (SAR) for the head (averaged over 10 g of tissue) did not exceed 2 W/kg. The active field remained within the SAR limit for the general population established by the International Commission on Non-Ionizing Radiation Protection (ICNIRP) and posed no known health risks. The two active fields administered are 5G uplink signals generated in the 5G frequency range. The lower frequency signal has a carrier frequency of 700 MHz, 20 MHz bandwidth, Frequency Division Duplexing/Orthogonal Frequency-Division Multiplexing (FDD/OFDM) with 24 resource blocks, 16 time slots, 60 kHz sub-carrier spacing and Quadrature Phase Shift Keying (QPSK) modulation, with an output power of 4.28 W. The higher frequency signal has a carrier frequency of 3.6 GHz, 100 MHz bandwidth, Time Division Duplexing (TDD)/QPSK OFDM with 135 resource blocks, 16 time slots, 60 kHz sub-carrier spacing and QPSK modulation, with an output power of 1.63 W. In the signals used, only uplink communication is implemented, and all frames are identical with 16 time slots. Both signals have identical power control applied that introduces low frequency amplitude modulation at 12.5 Hz on top of the modulation due to the occupied time slots which have a dominant power modulation frequency of 200 Hz resulting in a 14.2 dB peak to average power ratio (PAPR) (Supp. Figure 1, 2 & 3). The exposure levels in grey and white matter, thalamus and all tissues in the brain averaged over 0.125 g which is a cube of side length ~5 mm are reported in Table 1 for both 700 MHz and 3.6 GHz....

... we found a significant interaction between exposure and the genetic variant in the center frequency of sleep spindles. Specifically, we demonstrated a topographically widespread acceleration of spindle center frequency in the T/C carriers after exposure to the 3.6 GHz RF-EMF in comparison to sham....

By leveraging the strengths of spectral parameterization, we demonstrated a widespread shift in the center frequency of sleep spindles towards faster oscillatory activity in T/C allele carriers after exposure to a 5G RF-EMF with a carrier frequency of 3.6 GHz. This effect was evident in central, parietal, and occipital cortical areas coinciding with areas that predominantly express faster spindles. A number of previous studies also reported enhanced EEG spectral power in the upper spindle range after exposure to RF-EMF of earlier generation (Huber et al., 2000, 2002; Schmid, Loughran, et al., 2012; Schmid, Murbach, et al., 2012)....

The discrepancy between the deeper penetration of the 700 MHz signal revealed by the simulated SAR distribution in the brain and the more pronounced effects on the EEG sleep spindles observed following exposure to the 3.6 GHz signal remains unclear. Notably, the pulse modulation, which has been identified as critical for the biological effects of RF-EMF (Huber et al., 2002), was identical at 12.5 Hz in both fields and the psSAR10gr was consistently set at 2 W/kg. The findings underscore the necessity for a comprehensive investigation into the complex characteristics of the new 5G signals. Furthermore, they may suggest that the dielectric and conductive properties of the tissues associated with the minor allele may not be adequately represented by the current simulation parameters. Alternatively, the observed effects may indicate a distinct mode of action that is unrelated to SAR distribution....

The differential effects observed between the 700 MHz and 3.6 GHz exposures highlight the importance of considering signal characteristics and tissue properties in understanding RF-EMF interactions. Overall, our results provide new insights into the genetic and biophysical factors underlying RF-EMF effects on sleep, emphasizing the need for more targeted studies to elucidate these mechanisms.

Open access pre-print: https://www.medrxiv.org/content/10.1101/2024.12.16.24319082v1

--

RF-EMF Exposure near 5G NR Small Cells

Aerts S, Deprez K, Verloock L, Olsen RG, Martens L, Tran P, Joseph W. RF-EMF Exposure near 5G NR Small Cells. *Sensors*. 2023; 23(6):3145. doi: 10.3390/s23063145.

Abstract

Of particular interest within fifth generation (5G) cellular networks are the typical levels of radiofrequency (RF) electromagnetic fields (EMFs) emitted by 'small cells', low-power base stations, which are installed such that both workers and members of the general public can come in close proximity with them. In this study, RF-EMF measurements were performed near two 5G New Radio (NR) base stations, one with an Advanced Antenna System (AAS) capable of beamforming and the other a traditional microcell. At various positions near the base stations, with distances ranging between 0.5 m and 100 m, both the worst-case and time-averaged field levels under maximized downlink traffic load were assessed. Moreover, from these measurements, estimates were made of the typical exposures for various cases involving users and non-users. Comparison to the maximum permissible exposure limits issued by the International Commission on Non-Ionizing Radiation Protection (ICNIRP) resulted in maximum exposure ratios of 0.15 (occupational, at 0.5 m) and 0.68 (general public, at 1.3 m). The exposure of non-users was potentially much lower, depending on the activity of other users serviced by the base station and its beamforming capabilities: 5 to 30 times lower in the case of an AAS base station compared to barely lower to 30 times lower for a traditional antenna.

Excerpt

Scaled to small-cell powers, the measured exposure levels in this study were below the MPE limits for both occupational (at distances between 0.5 m and 1 m from the base station) and general public exposure (>1 m) issued by the International Commission on Non-Ionizing Radiation Protection (ICNIRP) [17]: the maximum exposure ratios were 0.15 (occupational) and 0.68 (general public). These theoretical worst-case exposures were higher than the actual maxima measured in situ by maximizing the downlink traffic load, and both types of worst-case exposures were much higher (3–12 times) than the exposures of a typical user, which in turn were much higher (4–9 times) than the exposures without any users. Finally, the exposure of a non-user within a mature 5G NR network depends on the distribution of users, their usage, and the AAS capabilities of the base station radio.

Although adverse health effects at non-thermal exposure levels cannot be ruled out [20], the ICNIRP reference levels are still relevant to calculate exposure ratios against. The measurement values obtained in this study and reported in this paper can be directly compared to other reference or limit levels (e.g., from legislation or scientific literature), depending on the scope of the study.

For a 5G NR AAS base station, the actual exposure of a given user will generally be less than the theoretical maximum exposure E_{max} for several reasons. First, other users (the number may vary) may be in beams other than the one the given user is using. Hence, the RF energy directed toward these users will not add (much) to the exposure of the given user. Second, the usage by the given user will generally be less than the maximum assumed for which E_{max} was defined. Third, there may be dynamic power control to reduce base station power to the minimum needed for communication. This was not taken into account in this study. Finally, the base station beam may be narrower or wider than that used to measure E_{max}. Hence, the problem of determining actual RF exposure becomes a statistical one that depends on several different variables. However, the exposure will almost always be less than E_{max}.

Similarly, for the non-user, the difference lies in whether they are in a beam or not. Generally, the RF exposure from the base station for the non-user will be smaller than that for a user (here by at least a factor of 5 for an AAS), unless there are many users around and the MaMIMO capabilities of the base station are limited (Table 4) [19].

Open access paper: https://www.mdpi.com/1424-8220/23/6/3145

--

Improving Monitoring of Indoor RF-EMF Exposure Using IoT-Embedded Sensors and Kriging Techniques

Jabeur R, Alaerjan A. Improving Monitoring of Indoor RF-EMF Exposure Using IoT-Embedded Sensors and Kriging Techniques. *Sensors*. 2024; 24(23):7849. https://doi.org/10.3390/s24237849

Abstract

Distributed wireless sensor networks (WSNs) are widely used to enhance the quality and safety of various applications. These networks consist of numerous sensor nodes, often deployed in challenging terrains where maintenance is difficult. Efficient monitoring approaches are essential to maximize the functionality and lifespan of each sensor node, thereby improving the overall performance of the WSN. In this study, we propose a method to efficiently monitor radiofrequency electromagnetic fields (RF-EMF) exposure using WSNs. Our approach leverages sensor nodes to provide real-time measurements, ensuring accurate and timely data collection. With the increasing prevalence of wireless communication systems, assessing RF-EMF exposure has become crucial due to public health concerns. Since individuals spend over 70% of their time indoors, it is vital to evaluate indoor RF-EMF exposure. However, this task is complicated by the complex indoor environments, furniture arrangements, temporal variability of exposure, numerous obstructions with unknown dielectric properties, and uncontrolled factors such as people's movements and the random positioning of furniture and doors. To address these challenges, we employ a sensor network to monitor RF-EMF exposure limits using embedded sensors. By integrating Internet of Things-embedded sensors with advanced modeling techniques, such as kriging, we characterize and model indoor RF-EMF downlink (DL) exposure effectively. Measurements taken in several buildings within a few hundred meters of base stations equipped with multiple cellular antennas (2G, 3G, 4G, and 5G) demonstrate that the kriging technique using the spherical model provides superior RF-EMF prediction compared with the exponential model. Using the spherical model, we constructed a high-resolution coverage map for the entire corridor, showcasing the effectiveness of our approach.

Conclusions

This study proposes the use of WSN to monitor the indoor RF-EMF exposure induced by cellular networks. To this end, we first proposed a measurement system based on the Narda NBM-550 and Nucleo-F401RE microcontroller board. The aim is to characterize and model indoor RF-EMF DL exposure using the collected measurements and kriging technique. First, several indoor measurements are conducted in an area covered by various frequency bands, including those used for 5G. By comparing the spherical and exponential models, we demonstrated that the spherical model provides a superior fit for predicting RF-EMF exposure levels. The high-resolution coverage map constructed using the spherical model revealed that the maximum average RF-EMF DL exposure levels within the corridor are well below the limits established by the ICNIRP. These findings underscore the effectiveness of the kriging technique in accurately modeling and predicting RF-EMF exposure in complex indoor environments.

For future work, several avenues can be explored to enhance the understanding and assessment of indoor RF-EMF exposure. Firstly, expanding the measurement campaign to include a wider variety of indoor environments, such as residential buildings, offices, and public transport, would provide a more comprehensive dataset. Additionally, incorporating temporal variations by conducting long-term measurements could offer insights into the fluctuations of RF-EMF exposure over time. Furthermore, integrating advanced machine learning algorithms with the kriging technique could improve the accuracy and efficiency of exposure predictions. Another important research axis involves the analysis of measurement uncertainty, which is planned for future investigation. For large-scale deployments, the Narda NBM-550 can be replaced with frequency-selective equipment such as the ExpoM-RF4, MVG EME Spy Evolution, or Narda SRM-3006. This substitution ensures that only downlink bands are considered and enables the reconstruction of RF-EMF exposure maps for each frequency band, facilitating a more detailed assessment by frequency band. Finally, investigating the impact of emerging wireless technologies, such as beyond 5G, on indoor RF-EMF exposure will be crucial as these technologies become more widespread. These future directions will contribute to a more thorough understanding of indoor RF-EMF exposure and help address public concerns regarding wireless communication systems.

Open access paper: https://www.mdpi.com/1424-8220/24/23/7849

--

AI-based optimization of EM radiation estimates from GSM base stations using traffic data

Lal R, Singh RK, Nishad DK, *et al.* AI-based optimization of EM radiation estimates from GSM base stations using traffic data. *Discov Appl Sci* 6, 655 (2024). doi: 10.1007/s42452-024-06395-y

Abstract

The fast expansion of mobile networks has sparked worries regarding base station EM radiation's health impacts. Traffic load is commonly ignored when evaluating EM radiation levels using maximum power output. This study proposes utilizing AI and ML on real network traffic data to optimize GSM base station EM radiation estimations. We obtained EM radiation measurements and traffic data from selecting GSM base stations by location and configuration. To predict EM radiation levels, traffic patterns were used to train linear regression, random forests, and neural networks. Base stations were clustered by radiation profile using unsupervised learning. Considering regulatory restrictions and

measurement feasibility, an optimization methodology was created to minimize EM radiation estimate inaccuracy. The results show better prediction accuracy than power-based estimations and high generalisability across base station types. Site-specific factors influenced daily EM radiation patterns after clustering. EM radiation levels can be monitored using traffic data and the optimized AI/ML model. This research helps telecom operators and regulators analyze EM radiation more accurately and efficiently. Future projects should include 5G and small cell network extensions and intelligent city platform integration. The suggested method develops data-driven, AI-powered Public Safety and mobile network trust solutions.

Open access paper: https://link.springer.com/article/10.1007/s42452-024-06395-y

--

Advancements in electromagnetic microwave absorbers: Ferrites and carbonaceous materials

Mohapatra PP, Singh HK, Dobbidi P. Advancements in electromagnetic microwave absorbers: Ferrites and carbonaceous materials. Adv Colloid Interface Sci. 2024 Dec 14;337:103381. doi: 10.1016/j.cis.2024.103381.

Abstract

Heightened levels of electromagnetic (EM) radiation emitted by electronic devices, communication equipment, and information processing technologies have become a significant concern recently. So, substantial efforts have been devoted for developing novel materials having high EM absorption properties. This critical review article provides an overview of the advancements in understanding and developing such materials. It delves into the interaction between EM radiation and absorbing materials, focusing on phenomena like multiple reflections, scattering, and polarization. Additionally, the study discusses various types of losses that impact microwave absorber performance, like magnetic loss, and dielectric loss. Each of these losses has distinct implications for microwave absorbers' effectiveness. Furthermore, the review offers detailed insights into different microwave-absorbing materials, such as metal composites, magnetic materials, conducting polymers, and carbonaceous materials (composites with carbon fiber, porous carbon, carbon nanotube, graphene oxide, etc.). Overall, it highlights the progress achieved in microwave-absorbing materials and emphasizes optimizing various loss mechanisms for enhanced performance.

Conclusions

This study provides a comprehensive overview of recent advancements in magnetic and carbon-based dielectric composites, showcasing their potential as promising materials for microwave absorption. The examples underscore that a single dielectric or magnetic system alone cannot consistently achieve optimal microwave absorption performance, necessitating the formulation of composite mixtures that incorporate dielectric and magnetic fillers. These composites' exceptional microwave absorption capabilities can be attributed to several key factors: optimizing intrinsic properties: The amalgamation of a carbon-based material with dielectric components yields distinctive complementary responses in their intrinsic properties, resulting in optimized impedance matching. Inducing Interfacial Polarization: Ample heterogeneous interfaces between different components effectively induce strong interfacial polarization, enhancing the overall microwave absorption performance. Facilitating Conductive Networks: Integrating multiple components facilitates the formation of a conductive network, hopping

electrons and fostering the migration and fortifying conductivity loss. Precision in Microstructure Design: Meticulously designed microstructures offer additional propagation paths for incident electromagnetic waves, stimulating multiple reflections and scatterings that efficiently consume electromagnetic energy.

Despite remarkable progress in carbon-based dielectric systems, persistent challenges warrant a strategic approach: In-Depth Exploration of EM Loss Characteristics: A meticulous exploration of the electromagnetic loss characteristics of each component is crucial. It emphasizes the need for a rational combination of elements rather than arbitrarily preparing multicomponent composites. Defect Engineering: While defects like grain boundaries, atom vacancies, and heteroatoms positively affect polarization and conductivity losses, the intricate relationship between defect sites and microwave absorption performance necessitates further exploration. Defect engineering is pivotal in guiding the fabrication of high-performance magnetic and carbon-based dielectric composites. Expanding the Effective Frequency Range: The effective frequency range of most composites is confined to 8.0–18.0 GHz, limiting their applicability in the electronics industry, where many devices operate at frequencies lower than 8.0 GHz. A rational construction approach for multicomponent composites with well-designed microstructures holds the potential to overcome this limitation and enhance low-frequency attenuation capabilities. Simplified Preparation Methods: The preparation methods, especially for multicomponent composites, are often complicated, posing challenges for large-scale production. Streamlining these methods is crucial for overcoming production difficulties.

In conclusion, the future outlook for high-performance materials against electromagnetic pollution lies in developing novel carbon-based dielectric systems, magnetic fillers with well-balanced compositions, and intricate microstructures. Addressing the outlined challenges will contribute to realizing these materials' potential to mitigate electromagnetic interference.

https://pubmed.ncbi.nlm.nih.gov/39700971/

--

Effect of radiation emitted from mobile phone on innate immunity in mice

Pei, Y., Gao, H., Zhang, M., Zhou, F., Zhu, Y., Wang, X., & Sun, J. (2024). Effect of radiation emitted from mobile phone on innate immunity in mice. *Radiation Effects and Defects in Solids*, *179*(11–12), 1585–1596. doi: 10.1080/10420150.2024.2352845

Abstract

This present study aims to explore the potential impact of cell phone radiation on innate immunity in mice. Ninety-six male BALB/C mice aged 2–3 weeks were randomly distributed into 4 groups as blank control, control, TD-SCDMA and LTE-Advanced respectively, with 32 mice in each group. Mice were designed to be exposed to cell phone radiation for 4–8 weeks. Eight mice in each group were taken out for measurement given exposure periods were 4, 6 and 8 weeks respectively. Cell biological technique was conducted to assess the chemotaxis of neutrophils, and a morphological method was performed for the detection of phagocytosis of neutrophil and macrophage, while microbiological means was carried out to test the relative activity of lysozyme in serum of mice. As a result, the chemotaxis ratio of neutrophils was with little statistical difference among the four groups given a shorter exposure period. However, the ratios in TD-SCDMA and LTE-Advanced groups were decreased significantly on the condition that the exposure period was more than 6 weeks. No statistical difference was observed

among the four groups during the entire exposure period in terms of the chemotaxis index. Phagocytosis of the innate cells as neutrophil and macrophage showed little change in the two control groups during the whole experimental stages, while the percentage in the two treated groups decreased statistically, and this kind of reduction was prone to feature time dependence. The activities of lysozyme in TD-SCDMA and LTE-Advanced groups declined significantly, further to that, the impact was climbing paralleled with the prolonged duration. It could be deduced that radiation from cell phones could weaken innate immunity in experimental mice; moreover, this adverse effect was seemingly more severe as the radiation exposure continued.

https://www.tandfonline.com/doi/full/10.1080/10420150.2024.2352845

--

Numerical dosimetry of specific absorption rate of insects exposed to far-field radiofrequency electromagnetic fields

Jeladze V, Nozadze T, Partsvania B, Thielens A, Shoshiashvili L, Gogoladze T (2025). Numerical dosimetry of specific absorption rate of insects exposed to far-field radiofrequency electromagnetic fields. *International Journal of Radiation Biology*, 1–14. doi: 10.1080/09553002.2024.2442693.

Abstract

Purpose This paper reports a study of electromagnetic field (EMF) exposure of several adult insects: a ladybug, a honey bee worker, a wasp, and a mantis at frequencies ranging from 2.5 to 100 GHz. The purpose was to estimate the specific absorption rate (SAR) in insect tissues, including the brain, in order to predict the possible biological effects caused by EMF energy absorption.

Method Numerical dosimetry was executed using the finite-difference time-domain (FDTD) method. Insects were modeled as 3-tissue heterogeneous dielectric objects, including the cuticle, the inner tissue, and the brain tissue. The EMF source was modeled as sinusoidal plane waves at a single frequency (far-field exposure).

Results The whole-body averaged, tissue averaged, and 1 milligram SAR values were determined in insects for all considered frequencies for 10 different incident plane waves. SAR values were normalized to the incident power density of 1 mW/cm2. Maximal EMF absorption in the inner and brain tissues was observed at 6, 12, and 25 GHz for the considered insects, except the brain tissue of a ladybug (max at 60 GHz).

Conclusion The paper presented the first estimation of the SAR for multiple insects over a wide range of RF frequencies using 3-tissue heterogenous insect 3D models created for this specific research. The selection of tissues' dielectric properties was validated. The obtained results showed that EMF energy absorption in insects highly depends on frequency, polarization, and insect morphology.

Conclusion

The paper presented a study of RF-EMF dosimetry of honeybee worker, wasp, mantis, and ladybug from 2.5 to 100 GHz, including frequencies that will be utilized in future 5 G technologies.

Discrete, 3-tissue heterogenous insect 3D models were created and used for FDTD modeling.

The whole-body averaged SAR values and tissue-averaged SAR values were estimated in insects' tissues for 9 considered frequencies and 10 polarizations of incident plane wave. For the first time, 1 mg SAR values were determined in insect tissues.

The obtained results showed SAR values in honeybee, wasp, ladybug, and mantis body tissues, which depend on the direction of the incident plane wave and polarization, frequency, and the insects' body sizes and peculiarities.

The highest values of the peak 1 mg SAR for the honeybee and wasp − 39.2 W/kg and 169.2 W/kg for an incident field strength of 1 mW/cm^2, were observed when E-field polarization was directed along the insect's length (pol. E3, E5, E9).

The obtained results showed maximal tissue-specific SAR values in the brain at 25 GHz for the honeybee (3.6 W/kg), 12 GHz for the wasp (5.4 W/kg), 25 GHz for the mantis (5.2 W/kg), and 60 GHz for the ladybug (10 W/kg), all for an incident power density of 1 mW/cm^2. Maximal EMF absorption in the inner tissue was observed at 12 GHz, 4.3 W/kg, 5.9 W/kg, 4.8 W/kg for the honey bee, wasp, and ladybug, respectively, and for the mantis 3.3 W/kg at 6 GHz for an incident power density of 1 mW/cm^2. The absorption in insects' cuticles increased proportionally with frequency. For example, for the ladybug, the tissue-specific SAR in the cuticle was 0.1 W/kg at 2.5 Ghz and 11.9 W/kg at 100 GHz for the same incident power density of 1 mW/cm^2.

Future studies will consider introducing insect models that will be obtained using micro-CT scanning, examining the effects of high-frequency electromagnetic fields on other insects of different forms and sizes, conducting thermal simulations along with EM simulations, and evaluating temperature rise in insect tissues.

Based on the present results, we expect this research to have an impact on (environmental) policymaking and standardization and regulation regarding RF-EMF emissions. We expect to contribute to the harmonization of 5G EMF safety and compliance doses and to the development of future recommendations about safe frequencies and doses of 5G-EMF on the organisms studied in the present research.

https://www.tandfonline.com/doi/full/10.1080/09553002.2024.2442693

--

Emerging cancer therapies: targeting physiological networks and cellular bioelectrical differences with non-thermal systemic electromagnetic fields in the human body – a comprehensive review

Costa FP, Wiedenmann B, Schöll E, Tuszynski J. Emerging cancer therapies: targeting physiological networks and cellular bioelectrical differences with non-thermal systemic electromagnetic fields in the human body – a comprehensive review. Frontiers in Network Physiology. Vol. 4, 2024. doi: 10.3389/fnetp.2024.1483401.

Abstract

A steadily increasing number of publications support the concept of physiological networks, and how

cellular bioelectrical properties drive cell proliferation and cell synchronization. All cells, especially cancer cells, are known to possess characteristic electrical properties critical for physiological behavior, with major differences between normal and cancer cell counterparts. This opportunity can be explored as a novel treatment modality in Oncology. Cancer cells exhibit autonomous oscillations, deviating from normal rhythms. In this context, a shift from a static view of cellular processes is required for a better understanding of the dynamic connections between cellular metabolism, gene expression, cell signaling and membrane polarization as states in constant flux in realistic human models. In oncology, radiofrequency electromagnetic fields have produced sustained responses and improved quality of life in cancer patients with minimal side effects. This review aims to show how non-thermal systemic radiofrequency electromagnetic fields leads to promising therapeutic responses at cellular and tissue levels in humans, supporting this newly emerging cancer treatment modality with early favorable clinical experience specifically in advanced cancer.

Open access paper:

https://www.frontiersin.org/journals/network-physiology/articles/10.3389/fnetp.2024.1483401

--

Effect of 6 GHz radiofrequency electromagnetic field on the development of fetal bones

Karamazı Y, Emre M, Uçar S, Aksoy G, Emre T, Tokuş M. (2024). Effect of 6 GHz radiofrequency electromagnetic field on the development of fetal bones. *Electromagnetic Biology and Medicine*, 1–9. doi: 10.1080/15368378.2024.2438608.

Abstract

This study examined the impact of 6 GHz (0.054 W/kg SAR) Radiofrequency-Electromagnetic Field (RF-EMF) on prenatal bone development. In this study, 20 female and 20 male Wistar Albino rats divided into four groups. The Control group received no treatment, while in Group-I, only male rats were exposed to RF-EMF, female rats had no exposure. Group-II, both male and female rats received RF-EMF treatment. While in Group-III, only female rats were exposed to RF-EMF, male rats had no exposure. The exposure lasted 4 hours per day for 6 weeks. The rats were then allowed to mate within the group. After pregnancy, pregnant rats (Group-II and III) were exposed 4 hours per day for 18 days. On the 18th day of gestation, fetuses were removed and their weight and various lengths were measured. The skeletal system development of fetuses was examined with double skeletal staining method and assessed ossification in the extremities. In the study, fetal weights, head-tail length, occipital-frontal and parietal-parietal lengths significantly increased in all exposure groups when compared to the control group ($p < 0.001$). Although occipital-frontal length was smallest in Group-I, Group-II and Group-III were more higher than the control group ($p < 0.001$). The bones of the anterior and posterior extremities showed significant increases in length, ossification zone length, and ossification percentage in all experimental groups compared to the control group ($p < 0.001$). Our study showed that rats exposed to 6 GHz (0.054 W/kg) RF-EMF during the prenatal period had significant increases in bone development.

Plain-Language Summary

Radiofrequency Electromagnetic Field (RF-EMF) sources are one of the most widely used technology systems in daily life. This study examined the impact of 6 GHz RF-EMF on prenatal bone development. In

this study, 20 female and 20 male Wistar Albino rats divided into four groups. The control group rats received no treatment, while in Group-I, only male rats, Group-II, both male and female rats, while in Group-III, only female rats were exposed to RF-EMF. The exposure lasted 4 hours/day for 6 weeks. The rats were then allowed to mate within the group. After pregnancy, pregnant rats (Group-II and III) were exposed 4 hours/day for 18 days. On the 18th day of gestation, fetuses were removed 10 fetuses each group were randomly selected in each group and their weight and various lengths were measured. The skeletal system development of fetuses was examined with double skeletal staining method. The ImageJ program was used assess ossification in the extremities.

https://www.tandfonline.com/doi/full/10.1080/15368378.2024.2438608

--

Effects of 4G Long-Term Evolution Electromagnetic Fields on Thyroid Hormone Dysfunction and Behavioral Changes in Adolescent Male Mice

Kim H-Y, Son Y, Jeong YJ, Lee S-H, Kim N, Ahn YH, Jeon SB, Choi H-D, Lee H-J. Effects of 4G Long-Term Evolution Electromagnetic Fields on Thyroid Hormone Dysfunction and Behavioral Changes in Adolescent Male Mice. *International Journal of Molecular Sciences*. 2024; 25(20):10875. doi: 10.3390/ijms252010875.

Abstract

Radiofrequency electromagnetic fields (RF-EMFs) can penetrate tissues and potentially influence endocrine and brain development. Despite increased mobile phone use among children and adolescents, the long-term effects of RF-EMF exposure on brain and endocrine development remain unclear. This study investigated the effects of long-term evolution band (LTE) EMF exposure on thyroid hormone levels, crucial for metabolism, growth, and development. Four-week-old male mice (C57BL/6) were exposed to LTE EMF (whole-body average specific absorption rate [SAR] 4 W/kg) or a positive control (lead; Pb, 300 ppm in drinking water) for 4 weeks. Subsequently, the mice underwent behavioral tests including open field, marble burying, and nest building. Blood pituitary and thyroid hormone levels, and thyroid hormone-regulating genes within the hypothalamus–pituitary–thyroid (HPT) axis were analyzed. LTE exposure increased T3 levels, while Pb exposure elevated T3 and T4 and decreased ACTH levels. The LTE EMF group showed no gene expression alterations in the thyroid and pituitary glands, but hypothalamic Dio2 and Dio3 expressions were significantly reduced compared to that in the sham-exposed group. Pb exposure altered the hypothalamic mRNA levels of Oatp1c1 and Trh, pituitary mRNA of Trhr, and Tpo and Tg expression in the thyroid. In conclusion, LTE EMF exposure altered hypothalamic Dio2 and Dio3 expression, potentially impacting the HPT axis function. Further research is needed to explore RF-EMF's impacts on the endocrine system.

Open access paper: https://www.mdpi.com/1422-0067/25/20/10875

--

Evaluation of the Thyroids of Offsprings Exposed to 2450 MHz Radiofrequency Radiation During Pregnancy: A Sixth Month Data

Ozyilmaz C, Oktay MF, Dasdag S, Ulukaya E, Genel ME, Tansuker HD, Emre F, Yeğin K. Evaluation of the

Thyroids of Offsprings Exposed to 2450 MHz Radiofrequency Radiation During Pregnancy: A Sixth Month Data. Journal of International Dental & Medical Research, 2024, 17(2): 925-930.

Abstract

This study aimed to determine whether the exposure to radiofrequency emitted by wireless internet providers (2450 MHz) throughout the day during rats' pregnancy causes a problem in the thyroid tissues of their offspring.

The pregnant rats in the experimental group were exposed to radiofrequency radiation (RFR) (24 hours/day) at a 2450 MHz frequency in pulse wave mode with 1 W output strength by a generator simulating Wi-Fi waves. The offspring in the control and experimental groups were randomized selected (n:8). At the end of the sixth month, the thyroid tissues were removed and evaluated histopathologically and biochemically. Mann–Whitney U-tests and T-tests were used for statistical analysis. The threshold for statistical significance was $p < 0.05$.

There was a significant difference in mononuclear cell infiltration ($p=0.03$) and vascular increase in congestion ($p<0.001$). There was no difference in the TUNEL-positive cell percentage ($p=0.62$) and H2A.X antibody levels ($p=0.68$) between the rats in the control and experimental groups. In this study, 2450 MHz RFR exposure during the prenatal period did not cause a statistically significant difference in terms of H2A.X levels and TUNEL-positive cell percentages in the thyroid tissue of rats.

Open access paper: http://www.jidmr.com/journal/wp-content/uploads/2024/06/71-M24_3112_Suleyman_DASDAG_Turkey-Exp-2.pdf

--

Effects of extremely low frequency magnetic fields on animal cancer and DNA damage: a systematic review and meta-analysis

Brabant C, Honvo G, Demonceau C, Tirelli E, Léonard F, Bruyère O. Effects of extremely low frequency magnetic fields on animal cancer and DNA damage: a systematic review and meta-analysis. Prog Biophys Mol Biol. 2024 Dec 31:S0079-6107(24)00116-0. doi: 10.1016/j.pbiomolbio.2024.12.005.

Abstract

The objective of this systematic review and meta-analysis is to assess the carcinogenic effects of extremely low frequency magnetic fields (ELF-MF) by analyzing animal and comet assay studies. We have performed a global meta-analysis on all the animal studies on the relation between ELF-MF and cancer incidence and separate meta-analyses on the incidence of cancer, leukemia, lymphoma, breast cancer, brain cancer and DNA damage assessed with the comet assay. Of the 5145 references identified, 71 studies have been included in our systematic review and 22 studies in our meta-analyses. Our global meta-analysis indicated that ELF-MF exposure had no significant impact on the incidence of cancers in rodents (19 studies, OR = 1.10; 95% CI 0.91-1.32). However, our separate meta-analyses showed that ELF-MF increased the odds of developing leukemia in mice (4 studies, OR = 4.45; 95% CI 1.90-10.38) but not in rats. Our systematic review also suggests that ELF-MF can damage DNA in certain cell types like brain cells. Nevertheless, a meta-analysis on three comet assay studies indicated that ELF-MF did not increase DNA damage in neuroblastoma cells (SMD = -0.08; 95% CI -0.18-0.01). Overall, our results

suggest that exposure to ELF-MF does not represent a major hazard for mammals and the carcinogenic effects of these magnetic fields could be limited to leukemia.

Highlights

ELF-MF do not affect the odds of lymphoma, brain cancer and breast cancer in rodents

ELF-MF could increase the odds of developing leukemia in mice but not in rats

ELF-MF have no influence on survival and body weight in rodents

ELF-MF do not increase DNA damage in neuroblastoma cells

ELF-MF could damage DNA of certain cell types like brain cells in rodents

https://pubmed.ncbi.nlm.nih.gov/39746455/

--

Effects of extremely low-frequency (50 Hz) electromagnetic fields on vital organs of adult Wistar rats and viability of mouse fibroblast cells

Tekam CKS, Majumdar S, Kumari P, Prajapati SK, Sahi AK, Singh R, Krishnamurthy S, Mahto SK. Effects of extremely low-frequency (50 Hz) electromagnetic fields on vital organs of adult Wistar rats and viability of mouse fibroblast cells. Radiat Prot Dosimetry. 2024 Dec 4:ncae220. doi: 10.1093/rpd/ncae220.

Abstract

In recent years, scientific communities have been concerned about the potential health effects of periodic electromagnetic field exposure (≤1 h/d). The objective of our study is to determine the impact of extremely low-frequency pulsed electromagnetic fields (ELF-PEMF) (1-3 mT, 50 Hz) on mouse fibroblast (red fluorescent protein (RFP)-L929) cells and adult Wistar rats to gain a comprehensive understanding of biological effects. We observed that RFP-L929 exhibits no significant changes in cell proliferation and morphology but mild elevation in aspartate aminotransferases, alanine aminotransferases, total bilirubin, serum creatinine, and creatine kinase-myocardial band levels in ELF-PEMF exposed groups under in vitro and in vivo conditions. However, the histological examination showed no significant alterations in tissue structure and morphologies. Our result suggests that 50-Hz ELF-PEMF exposure (1-3 mT, 50 Hz) with duration (<1 h/d) can trigger mild changes in biochemical parameters, but it is insufficient to induce any pathological alterations.

Conclusions

The present study demonstrates the effects of 50-Hz ELF-PEMF (1–3 mT) using *in vitro* (RFP-L929 mouse fibroblast cells) and *in vivo* (adult male Wistar rats) models. The results reveal that exposure duration of 20 min (each) with a 4-h gap is non-destructive for RFP-L929 cells and causes mild alterations in biochemical parameters but not in organ coefficient, tissue structure, and morphology of adult Wistar rats. The results demonstrated that 50-Hz ELF-PEMF exposure did not cause significant cellular fragmentation and changes in the morphology of mouse fibroblast cells. We observed mild alterations in the biochemical parameters of rats among MF exposed and control groups. Conversely, histological analysis of the selected liver, kidney, and heart sections following ELF-PEMF exposure revealed no significant changes in tissue structure and morphology. Our efforts provide conceptual and experimental

support to establish a link between 50-Hz ELF-PEMF exposure and biological systems/tissues in residential and occupational environments.

https://pubmed.ncbi.nlm.nih.gov/39656829/

--

Extremely Low-Frequency Electromagnetic Field (ELF-EMF) Increases Mitochondrial Electron Transport Chain Activities and Ameliorates Depressive Behaviors in Mice

Teranishi M, Ito M, Huang Z, Nishiyama Y, Masuda A, Mino H, Tachibana M, Inada T, Ohno K. Extremely Low-Frequency Electromagnetic Field (ELF-EMF) Increases Mitochondrial Electron Transport Chain Activities and Ameliorates Depressive Behaviors in Mice. *International Journal of Molecular Sciences*. 2024; 25(20):11315. https://doi.org/10.3390/ijms252011315

Abstract

Compromised mitochondrial electron transport chain (ETC) activities are associated with depression in humans and rodents. However, the effects of the enhancement of mitochondrial ETC activities on depression remain elusive. We recently reported that an extremely low-frequency electromagnetic field (ELF-EMF) of as low as 10 μT induced hormetic activation of mitochondrial ETC complexes in human/mouse cultured cells and mouse livers. Chronic social defeat stress (CSDS) for 10 consecutive days caused behavioral defects mimicking depression in mice, and using an ELF-EMF for two to six weeks ameliorated them. CSDS variably decreased the mitochondrial ETC proteins in the prefrontal cortex (PFC) in 10 days, which were increased by an ELF-EMF in six weeks. CSDS had no effect on the mitochondrial oxygen consumption rate in the PFC in 10 days, but using an ELF-EMF for six weeks enhanced it. CSDS inactivated SOD2 by enhancing its acetylation and increased lipid peroxidation in the PFC. In contrast, the ELF-EMF activated the Sirt3-FoxO3a-SOD2 pathway and suppressed lipid peroxidation. Furthermore, CSDS increased markers for mitophagy, which was suppressed by the ELF-EMF in six weeks. The ELF-EMF exerted beneficial hormetic effects on mitochondrial energy production, mitochondrial antioxidation, and mitochondrial dynamics in a mouse model of depression. We envisage that an ELF-EMF is a promising therapeutic option for depression.

Open access paper: https://www.mdpi.com/1422-0067/25/20/11315

--

Effects of light, electromagnetic fields and water on biological rhythms

Martel J, Rouleau N, Murugan NJ, Chin WC, Ojcius DM, Young JD. Effects of light, electromagnetic fields and water on biological rhythms. Biomed J. 2024 Dec 11:100824. doi: 10.1016/j.bj.2024.100824.

Abstract

The circadian rhythm controls a wide range of functions in the human body and is required for optimal health. Disruption of the circadian rhythm can produce inflammation and initiate or aggravate chronic diseases. The modern lifestyle involves long indoor hours under artificial lighting conditions as well as eating, working, and sleeping at irregular times, which can disrupt the circadian rhythm and lead to poor health outcomes. Seasonal solar variations, the sunspot cycle and anthropogenic electromagnetic fields can also influence biological rhythms. The possible mechanisms underlying these effects are discussed,

which include resonance, radical-pair formation in retina cryptochromes, ion cyclotron resonance, and interference, ultimately leading to variations in melatonin and cortisol. Intracellular water, which represents a coherent, ordered phase that is sensitive to infrared light and electromagnetic fields, may also respond to solar variations and man-made electromagnetic fields. We describe here various factors and underlying mechanisms that affect the regulation of biological rhythms, with the aim of providing practical measures to improve human health.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S2319417024001276?via%3Dihub

--

The origins of light-independent magnetoreception in humans

Shibata T, Hattori N, Nishijo H, Kuroda S, Takakusaki K. The origins of light-independent magnetoreception in humans. Front Hum Neurosci. 2024 Nov 29;18:1482872. doi: 10.3389/fnhum.2024.1482872.

Abstract

The Earth's abundance of iron has played a crucial role in both generating its geomagnetic field and contributing to the development of early life. In ancient oceans, iron ions, particularly around deep-sea hydrothermal vents, might have catalyzed the formation of macromolecules, leading to the emergence of life and the Last Universal Common Ancestor. Iron continued to influence catalysis, metabolism, and molecular evolution, resulting in the creation of magnetosome gene clusters in magnetotactic bacteria, which enabled these unicellular organisms to detect geomagnetic field. Although humans lack a clearly identified organ for geomagnetic sensing, many life forms have adapted to geomagnetic field-even in deep-sea environments-through mechanisms beyond the conventional five senses. Research indicates that zebrafish hindbrains are sensitive to magnetic fields, the semicircular canals of pigeons respond to weak potential changes through electromagnetic induction, and human brainwaves respond to magnetic fields in darkness. This suggests that the trigeminal brainstem nucleus and vestibular nuclei, which integrate multimodal magnetic information, might play a role in geomagnetic processing. From iron-based metabolic systems to magnetic sensing in neurons, the evolution of life reflects ongoing adaptation to geomagnetic field. However, since magnetite-activated, torque-based ion channels within cell membranes have not yet been identified, specialized sensory structures like the semicircular canals might still be necessary for detecting geomagnetic orientation. This mini-review explores the evolution of life from Earth's formation to light-independent human magnetoreception, examining both the magnetite hypothesis and the electromagnetic induction hypothesis as potential mechanisms for human geomagnetic detection.

Conclusion

Eukaryotic cells, and vertebrates have developed magnetoreception systems to adapt to the geomagnetic field. Numerous studies on magnetoreception in birds, particularly concerning the upper beak and inner ear, suggest that in humans, the trigeminal nerve, vestibular nerve, and hindbrain might be involved in light-independent magnetoreception pathways. However, the specific sensory organ in humans that detects the geomagnetic field has not yet been identified. Unlike traditional senses, geomagnetic information is transmitted without attenuation through the scalp, bones, and

cerebrospinal fluid, similar to gravity. It also generates eddy currents and lorentz forces accompanying the relative movement of the geomagnetic field (Arago and Flourens, 1856). Considering these unique properties, vertebrates might have evolved to detect the geomagnetic field in a light-independent manner, not only through direct detection using torque-based magnetic particles but also through indirect detection of electric potentials using electromagnetic induction. Further research into this evolutionary adaptation could help unravel the mystery of geomagnetic field detection in humans.

Open access paper: https://www.frontiersin.org/journals/human-neuroscience/articles/10.3389/fnhum.2024.1482872/full

--

Effects of anthropogenic electromagnetic fields used for subsurface oil and gas exploration (controlled-source electromagnetics, CSEM) on the early development of Atlantic haddock

Guillebon C, Perrichon P, Browman HI, Cresci A, Sivle LD, Skiftesvik AB, Zhang G, Durif CMF. Effects of anthropogenic electromagnetic fields used for subsurface oil and gas exploration (controlled-source electromagnetics, CSEM) on the early development of Atlantic haddock (Melanogrammus aeglefinus). Mar Pollut Bull. 2024 Dec 12;211:117425. doi: 10.1016/j.marpolbul.2024.117425.

Abstract

Controlled source electromagnetics (CSEM) uses electromagnetic fields (EMF) to detect oil reservoirs. Atlantic haddock, Melanogrammus aeglefinus, is a commercially important demersal fish species that can potentially be impacted by such surveys due to potential overlap with egg distribution. In this study, haddock eggs were exposed to EMF, replicating CSEM survey conditions in a laboratory. Three different EMF intensities were used to replicate different distances between the EMF source and the organism. Exposures lasted for 15 min. A worst-case scenario, i.e. 1 h exposure at the highest EMF level was also carried out. None of the treatments caused malformations, mortality or affected hatching of eggs. However, EMF exposure induced tachycardia in newly hatched larvae and reduced the size of their yolk sac reserve. The effect was significant at the lowest EMF intensity (corresponding to 1000 m between the EMF source and the exposed subject) and increased with exposure time and intensity

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0025326X24014024?via%3Dihub

--

The effect of mobile phone electromagnetic fields on the human resting state wake EEG and event-related potential: A systematic review and meta-analysis

Prins AC, Baas K, van der Meer JN, Jacobs M, Nederveen AJ. The effect of mobile phone electromagnetic fields on the human resting state wake EEG and event-related potential: A systematic review and meta-analysis. Bioelectromagnetics. 2024 Nov 22. doi: 10.1002/bem.22531.

Abstract

The rapid growth of mobile phone usage and its use of radiofrequency electromagnetic fields (RF-EMF) have raised concerns about potential health risks. Researchers have conducted studies to examine the effects of RF-EMF on the brain using electroencephalography (EEG). We conducted a systematic quality

assessment and meta-analysis of published research in this field to establish high-quality studies as references for future protocols. The electronic search yielded 244 records from which a total of 51 studies were included in the review after excluding studies based on study design, and data or report availability. Of these 51 studies, 31 (61%) focused on resting state wake EEG and 20 (39%) on event-related potentials (ERP). None of the 51 studies were free from risk of bias. From the 51 included studies, we were able to use seven studies to create three different groups for meta-analysis for resting state wake EEG and five studies to create 10 different groups for meta-analysis for ERP. Per group the number of studies varies from 1 to 5. Our procedure is the first systematic quality assessment in this field and revealed three important findings. First, there is evidence of an effect on the EEG of a 2G protocol using an eyes-open condition. Second, we did not find evidence for EEG effects during task performance. This suggests that the impact of EMF during task performance is less pronounced compared to the resting state condition. Third, this meta-analysis shows that the field is unable to create an evidence base for most comparisons due to heterogeneity. We therefore advise that all future studies are double-blind in nature, adhere to the methodological standard of randomized experiments, and publish their protocols first.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.22531

--

Electromagnetic fields exposure on fetal and childhood abnormalities: Systematic review and meta-analysis

Kashani ZA, Pakzad R, Fakari FR, Haghparast MS, Abdi F, Kiani Z, Talebi A, Haghgoo SM. Electromagnetic fields exposure on fetal and childhood abnormalities: Systematic review and meta-analysis. Open Med (Wars). 2023 May 12;18(1):20230697. doi: 10.1515/med-2023-0697

Abstract

Today, in the modern world, people are often exposed to electromagnetic waves, which can have undesirable effects on cell components that lead to differentiation and abnormalities in cell proliferation, deoxyribonucleic acid (DNA) damage, chromosomal abnormalities, cancers, and birth defects. This study aimed to investigate the effect of electromagnetic waves on fetal and childhood abnormalities. PubMed, Scopus, Web of Science, ProQuest, Cochrane Library, and Google Scholar were searched on 1 January 2023. The Cochran's *Q*-test and I^2 statistics were applied to assess heterogeneity, a random-effects model was used to estimate the pooled odds ratio (OR), standardized mean difference (SMD), and mean difference for different outcomes, and a meta-regression method was utilized to investigate the factors affecting heterogeneity between studies. A total of 14 studies were included in the analysis, and the outcomes investigated were: change in gene expression, oxidant parameters, antioxidant parameters, and DNA damage parameters in the umbilical cord blood of the fetus and fetal developmental disorders, cancers, and childhood development disorders. Totally, the events of fetal and childhood abnormalities were more common in parents who have been exposed to EMFs compared to those who have not (SMD and 95% confidence interval [CI], 0.25 [0.15-0.35]; I^2, 91%). Moreover, fetal developmental disorders (OR, 1.34; CI, 1.17-1.52; I^2, 0%); cancer (OR, 1.14; CI, 1.05-1.23; I^2, 60.1%); childhood development disorders (OR, 2.10; CI, 1.00-3.21; I^2, 0%); changes in gene expression (mean difference [MD], 1.02; CI, 0.67-1.37; I^2, 93%); oxidant parameters (MD, 0.94; CI, 0.70-1.18; I^2, 61.3%); and DNA damage parameters (MD, 1.01; CI, 0.17-1.86; I^2, 91.6%) in parents who have been exposed to

EMFs were more than those in parents who have not. According to meta-regression, publication year has a significant effect on heterogeneity (coefficient: 0.033; 0.009-0.057). Maternal exposure to electromagnetic fields, especially in the first trimester of pregnancy, due to the high level of stem cells and their high sensitivity to this radiation, the biochemical parameters of the umbilical cord blood examined was shown increased oxidative stress reactions, changes in protein gene expression, DNA damage, and increased embryonic abnormalities. In addition, parental exposure to ionizing and non-ionizing radiation can lead to the enhancement of different cell-based cancers and developmental disorders such as speech problems in childhood.

Conclusions

Studies are associated with mothers' exposure to EMFs during pregnancy and non-ionizing radiation (RF and ELF-EMFs) with fetal complications such as significant enhancement of oxidant factors, decrease of antioxidant factors, and increase in DNA damage parameters, as well as changes in expression proteins in cord blood genes. On the other hand, close maternal exposure in prenatal and postnatal (residence or occupational exposure) with EMFs of high voltages power lines more than 1 mG or 50 Hz with congenital anomalies (CNS defect, spina bifida) and fetal developmental disorders (such as reduced embryonic bud length) and neurodevelopmental disorders in childhood (e.g., speech problems in children) are associated. Moreover, parents' exposure to ionizing and non-ionizing radiation (X-ray, ELF-EMF, and RF examinations) before and after birth is also associated with enhancement of the risk of cancers (such as rhabdomyosarcoma, ALL, brain tumors, and neuroectoderm) in childhood and adolescence. However, due to the limitations of studies, such as inaccurate measurement of exposure to ELF-EMF (e.g., interviews based on participants' reminders) or inaccurate measure of the actual rate of exposure to EMF or case–control model of most studies, the effects of EMF on fetal and childhood abnormalities should be interpreted with caution. Considering the widespread exposure to non-ionizing radiation, a little enhancement of exposure to EMF could lead to unacceptable health consequences for future generations. Although the number of epidemiological studies examining the undesirable effects of EMF exposure in humans is limited, the findings of this study should prompt further research on this significant environmental danger for pregnant women.

Open access paper: https://www.degruyter.com/document/doi/10.1515/med-2023-0697/html?lang=en

--

Systematic Review of Exposure Studies to Radiofrequency Electromagnetic Fields: Spot Measurements and Mixed Methodologies

Ramirez-Vazquez R, Escobar I, Arribas E, Vandenbosch GAE. Systematic Review of Exposure Studies to Radiofrequency Electromagnetic Fields: Spot Measurements and Mixed Methodologies. *Applied Sciences*. 2024; 14(23):11161. https://doi.org/10.3390/app142311161

Abstract

This work presents a review and evaluation of studies measuring exposure to Radiofrequency Electromagnetic Fields (RF-EMF). The review meets the basic quality criteria and the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines after the eligibility criteria of the PECO (Population, Exposure, Comparator, and Outcome) methodology and the instrument Critical Appraisal Skills Programme Español (CASPe). A total of 86 papers published between 1 January 1998 and

31 December 2023 are included: 61 studies with spot measurements and 25 studies with mixed methodologies (spot measurements, personal measurements with volunteers or with a trained researcher and prediction models) are highlighted. Forty-three percent of the studies use Spectrum Analyzers in the spot measurements, mainly the Narda SRM–3006, followed by the Narda SRM-3000, highlighting the introduction and use of Sensors for this kind of study. The minimum mean value was measured in Palestine at 0.0600 $\mu W/m^2$, and the maximum mean value was measured in Norway at 200,000 $\mu W/m^2$. The RF-EMF exposure levels measured in the different microenvironments are minimal and far from the maximum levels established by the ICNIRP guidelines.

Conclusions

This study reviewed 86 scientific works on personal exposure to RF-EMF that conducted measurements through spot measurements and/or using different methods (mixed method) combining spot measurements, personal measurements with volunteers or with trained researchers, spatiotemporal exposure measurements and in some cases using model predictions.

In addition to the exposure measurement studies, we want to refer to a study recently published by McKenzie [143] in which electromagnetic fields produced by “smart” devices used daily in a modern “smart” home are measured. It is interesting to see the activity of these EMFs in different periods of time while they are used. This study has not been included in this review because although specific measures have been taken, exposure to these fields has not been measured.

As far as we know, today, despite the different methodologies used, the results are comparable with international reference levels and comparable between microenvironments and countries; although it is true that some values are higher than others, they are still lower than these reference levels. The minimum average was measured in Palestine at 0.0600 μW/m2 in 2017, and the maximum average was measured in Norway at 200,000 μW/m2 in 2022, equivalent to 0.200 W/m2, far below the permitted international reference levels. During the development of this work, we have seen that the statistical data confirm that the exposure levels to RF-EMF are much lower than the maximum levels allowed in the ICNIRP guidelines, both for the general public scenario (10 W/m2) and for the occupational scenario (50 W/m2). However, these values can change over time, considering the new 5G technologies introduced everywhere right at this moment, so it is recommended to stay vigilant and continue expanding research in this field.

As previously indicated, with this work, we want to help researchers scrutinize this field with a review that classifies, orders, and synthesizes, thus making a reference for future research and comparisons. We believe that in the future, some research areas may be prioritized in the field of radiofrequency electromagnetic exposure fields: exploring new measurement devices, such as the use of sensors that have already begun to be used to monitor RF-EMF [111], as well as carrying out longitudinal studies to evaluate possible long-term effects.

We want to highlight that in this society in which we currently live, in which we are always immersed in a sea of electromagnetic waves, the importance of clear and effective communication of the possible risks associated with RF-EMF exposure, as well as the need to educate the public on how to minimize exposure in everyday environments.

Open access paper: https://www.mdpi.com/2076-3417/14/23/11161

--

Causal associations between mobile phone usage and glaucoma risk: A Mendelian randomization study

Song R, Wang Y, Kong Y, Fan X, Yuan C, Zha X. Causal associations between mobile phone usage and glaucoma risk: A Mendelian randomization study. Medicine (Baltimore). 2024 Nov 29;103(48):e40666. doi: 10.1097/MD.0000000000040666.

Abstract

Previous research has indicated a possible link between mobile phone usage and the incidence of glaucoma. This study employs a 2-sample Mendelian randomization (MR) analysis to examine the causal relationship between mobile phone use and glaucoma risk. We used single nucleotide polymorphisms (SNPs) from publicly accessible genome-wide association study (GWAS) datasets as instrumental variables (IVs). The primary analytical method was the inverse variance weighted (IVW) approach, with MR-Egger and weighted median analyses serving as complementary methods. Sensitivity was evaluated using Cochran's Q test and MR-Egger regression. The results demonstrate a causal effect of mobile phone usage on an increased risk of glaucoma (OR IVW = 1.358, 95% CI: 1.052-1.752, P = .019; OR MR-Egger = 1.882, 95% CI: 0.53-6.682, P = .337; OR Weighted median = 1.387, 95% CI: 1.012-1.900, P = .042; OR MR-PRESSO = 1.358, 95% CI: 1.052-1.752, P = .026). Sensitivity analyses confirmed the robustness and reliability of these findings. The study identifies mobile phone usage as a potentially modifiable risk factor for glaucoma, providing new avenues for exploring the specific mechanisms underlying these ocular disorders.

Open access paper: https://journals.lww.com/md-journal/fulltext/2024/11290/causal_associations_between_mobile_phone_usage_and.12.aspx

--

Association between mobile phone use and incidence of dementia: A prospective cohort study using the UK Biobank

Zhao H, Wen Q, Zhuo L, Wang S, Zhan S. Association between mobile phone use and incidence of dementia: A prospective cohort study using the UK Biobank. Gerontology. 2023 Jul 26. doi: 10.1159/000531847.

Abstract

Introduction: The rapid growth in mobile phone use has led to public concern about its potential effects on the risk of dementia. This study aimed to investigate the association between mobile phone use in daily life and the risk of dementia incidence in community-dwelling adults based on the data from the UK Biobank.

Methods: Participants in the UK Biobank aged 60 years or older with no diagnosis of dementia at the time of recruitment were included in this prospective cohort study. A Cox regression model adjusted for sociodemographic characteristics, general health factors, mental health, lifestyle factors, comorbidities,

and medication use was used to estimate the hazard ratio (HR) and confidence interval (CI) of the association between mobile phone use and dementia risk.

Results: The final analyses included 213,181 participants. During a median follow-up period of 12.4 years, 6,344 cases of incident dementia occurred. Mobile phone use displayed a modest association with lower risk of dementia incidence, with HRs of 0.85 (95% CI: 0.79-0.91), 0.85 (95% CI: 0.80-0.91), 0.78 (95% C: 0.71-0.86), 0.86 (95% CI: 0.77-0.96), and 0.83 (95% CI: 0.70-0.98) for participants who reported phone call usage of fewer than 5 minutes, 5-29 minutes, 30-59 minutes, 1-3 hours, and more than 3 hours per week, respectively, compared with non-users. In addition, the proportions of the association medicated by family/friend visits and other leisure/social activities were 2.62% (95% CI: -0.64-6.51) and 2.22% (95% CI: 1.12-4.12), respectively.

Conclusions: Daily mobile phone use is significantly associated with a reduced risk of incident dementia in community-dwelling adults in the UK Biobank population. This association seems to be mediated by improved social and mental activities.

https://pubmed.ncbi.nlm.nih.gov/37494916/

--

Protective Properties of Botanical Extracts against 5G Radiation-induced Damage to Human Skin, as Demonstrated in Preliminary Data from a Keratinocyte Cell Culture Model

Havas F, Cohen M, Krispin S, Attia-Vigneau J. Protective Properties of Botanical Extracts against 5G Radiation-induced Damage to Human Skin, as Demonstrated in Preliminary Data from a Keratinocyte Cell Culture Model. *Front. Biosci. (Landmark Ed)* 2024, 29(1). doi: 10.31083/j.fbl2901031.

Abstract

Background: Next-generation 5G communication technology involves increasing use of 3–100 GHz wireless bands in population centers. Though still non-ionizing, this implies higher radiation energy vs. existing bands. The range is also shorter, needing more numerous emitters, closer to the user—resulting in higher electromagnetic energy exposure. With no universal consensus regarding exposure risks, there is some concern among the public and the scientific community, following indications that 5G radiation can impact immune function, trigger inflammatory responses, and influence expression of genes affecting protein folding, oxidative stress, tissue/extracellular matrix (ECM) matrix turnover, and more. This work aims at identifying botanical extracts for protection of human skin from these impacts, based on a preliminary cell culture-based model.

Methods: We irradiated human epidermal keratinocytes at 6 GHz, evaluating effects on Interleukin1-α (IL1-α), a key inflammatory cytokine; TIMP metallopeptidase inhibitor 1 (TIMP1), shown to inhibit collagenase; Angiopoietin-like protein 4 (ANGPLT4), which plays a role in wound healing and epidermal differentiation; and S100 calcium-binding protein A9 (S100A9), involved in immune recruitment during injury, by enzyme-linked immunosorbent assay (ELISA) and immunostaining. We next used this model to identify substances able to mitigate the effects of 5G irradiation, through the evaluation of the influence of treatment by one of several botanical extracts on the observed effects of 5G irradiation.

Results: After a remarkably short 1-h exposure, clear effects on keratinocyte function were observed: increased inflammatory cytokine IL1-α; reduced collagenase inhibitor TIMP1; increased wound healing/differentiation facilitator ANGPLT4; and increased SA100A9, involved in immune recruitment during injury. On this basis, we then showed the protective effects of selected botanical extracts, capable of reducing the increase in IL1-α induced by 5G exposure, possibly in part due to anti-inflammatory and antioxidant properties of compounds present in these extracts.

Conclusions: Our results show a clear influence of 5G irradiation on the keratinocytes, possibly indicating injury and damage responses. What's more, we showed how these preliminary data can be used to identify botanical extracts capable of offering some protection against these effects for users of 5G technology, e.g., when employed as active ingredients in protective cosmetic applications.

Open access paper: https://doi.org/10.31083/j.fbl2901031

--

Preliminary Study on the Impact of 900MHz Radiation on Human Sperm: An In Vitro Molecular Approach

Keskin I, Karabulut S, Kaplan AA, Alagöz M, Akdeniz M, Tüfekci KK, Davis DL, Kaplan S. Preliminary Study on the Impact of 900MHz Radiation on Human Sperm: An In Vitro Molecular Approach. Reprod Toxicol. 2024 Nov 4:108744. doi: 10.1016/j.reprotox.2024.108744.

Abstract

The use of technologies that produce and emit electromagnetic fields (EMF) is growing exponentially worldwide. The biological effects of EMF-emitting equipment, such as mobile phones and other wireless devices, have been studied in the last decade using in vitro and in vivo methods. Infertility is a growing health problem, and nearly half of cases are because of male-factor. This study investigated the direct in vitro effects of 900 MHz radiation exposure on sperm parameters, genetic status, apoptotic markers, and the PI3K/AKT signaling pathway in healthy normozoospermic men. Semen samples were divided into four groups, two control (30 min and 1 h) and two EMF exposure (30 min and 1 h). Sperm parameters (motility, progressive motility, acrosomal index, morphology), genetic status (DNA fragmentation and chromatin integrity), apoptotic markers (cytokine-c and caspase-3 expression) and the PI3K/AKT signaling pathway (phosphoinoitide 3-kinase-PI3K- and phosphorylated AKT- p-AKT-) were analysed. Sperm motility were significantly reduced in 30 min EMF exposure while a significant increase in the expression of p-AKT were observed in 1 h EMF exposure group. An increased vacuolisation, acrosomal defect, extension of subacrosomal space, uncondensed chromatin structure, apoptotic signs and disrupted axoneme were observed in both EMF groups which were not observed in the control group. Other sperm parameters (morphology and acrosomal index), genetic status, apoptotic markers and the PI3K expression rates had no significant change.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0890623824002119

--

Sensory Processing Sensitivity, and Not Gender, Drives Electromagnetic Hypersensitivity and Nature Connection

Watten RG, Volden F, Visnes H. Sensory Processing Sensitivity, and Not Gender, Drives Electromagnetic Hypersensitivity and Nature Connection. Ecopsychology. Nov 28, 2024. https://doi.org/10.1089/eco.2024.0027

Abstract

Electromagnetic hypersensitivity (EHS) is a biomedical condition associated with exposure to man-made radio-frequency electromagnetic fields. Common sources are devices like mobile phones, wireless networks, base stations, computers, TVs, and more. Identifying individuals at risk of developing EHS is important. Those with heightened sensory processing sensitivity (SPS; the ability to perceive, process, and react to environmental stimuli, approximately 25% to 30% of the population) are of special interest. SPS could also be associated with connectedness to nature (CNS) and EHS. In the current gender-matched cross-sectional study (*n* = 450; 225 men and 225 women) we assessed gender differences in environmental sensitivity assessed as SPS, EHS, and CNS. Women had higher general EHS prevalence than men (13.3% vs. 5.3%), higher mean values on perceived sensitivity for 5 out of 9 categories of electromagnetic equipment (computers, electrical appliances, fluorescent lighting, mobile phones, and television) and they had higher scores on the three EHS Scales EHS General, EHS Scale, and EHS Index, in addition to higher SPS and CNS scores. The gender differences vanished when adjusting for SPS in the multivariate analyses of variance (MANOVA) analyses, suggesting that the differences in these scales were associated with SPS. The statistical equation modeling (SEM) results showed a significant positive direct effect of SPS on CNS, and a significant indirect effect mediated by EHS. The present study provides the first empirical evidence that highly sensitive individuals are capable of perceiving electromagnetic radiation. The results indicate that perceived EHS and CNS are driven by SPS and not by gender.

https://www.liebertpub.com/doi/10.1089/eco.2024.0027

--

Exploring RF-EMF levels in Swiss microenvironments: An evaluation of environmental and auto-induced downlink and uplink exposure in the era of 5G

Veludo AF, Stroobandt B, Van Bladel H, Sandoval-Diez N, Guxens M, Joseph W, Röösli M. Exploring RF-EMF levels in Swiss microenvironments: An evaluation of environmental and auto-induced downlink and uplink exposure in the era of 5G. Environmental Research, 2024, doi: 10.1016/j.envres.2024.120550.

Highlights

A new protocol was created to measure environmental and auto-induced RF-EMF levels.

Environmental RF-EMF was mainly attributed to downlink frequency bands

Inducing downlink and uplink traffic increased RF-EMF exposure levels notably

Auto-induced downlink exposure was mainly attributed to the 5G band at 3.5 GHz

The main contributor to auto-induced uplink exposure was the band at 2.1 GHz

Abstract

The advancement of cellular networks requires updating measurement protocols to better study radiofrequency electromagnetic field (RF-EMF) exposure emitted from devices and base stations. This paper aims to present a novel activity-based microenvironmental survey protocol to measure environmental, auto-induced downlink (DL), and uplink (UL) RF-EMF exposure in the era of 5G. We present results when applying the protocol in Switzerland. Five study areas with different degrees of urbanization were selected, in which microenvironments were defined to assess RF-EMF exposure in the population. Three scenarios of data transmission were performed using a user equipment in flight mode (non-user), inducing DL traffic (max DL), or UL traffic (max UL). The exposimeter ExpoM-RF 4, continuously measuring 35 frequency bands ranging from broadcasting to Wi-Fi sources, was carried in a backpack and placed 30cm apart from the user equipment. The highest median RF-EMF levels during the non-user scenario were measured in an urban business area (1.02 mW/m2). Here, DL and broadcasting bands contributed the most to total RF-EMF levels. Compared to the non-user scenario, exposure levels increased substantially during max DL due to the 5G band at 3.5 GHz with 50% of the median levels between 3.20-12.13 mW/m2, mostly in urban areas. Note that the time-division nature of this band prevents distinguishing between exposure contribution from DL beamforming or UL signals emitted at this frequency. The highest levels were measured during max UL, especially in rural microenvironments, with 50% of the median levels between 12.08-37.50 mW/m2. Mobile UL 2.1 GHz band was the primary contributor to exposure during this scenario. The protocol was successfully applied in Switzerland and used in nine additional countries. Inducing DL and UL traffic resulted in a substantial increase in exposure, whereas environmental exposure levels remained similar to previous studies. This data is important for epidemiological research and risk communication/management.

Conclusion

A novel activity-based microenvironmental survey protocol was developed and successfully carried out to disentangle environmental from auto-induced downlink and uplink exposure in the era of 5G. The measurements conducted in Switzerland demonstrate that higher RF-EMF exposure levels were measured when inducing maximum downlink and uplink traffic using a user equipment, with the 5G band at 3.5 GHz and the UL band at 2.1 GHz the main contributors to exposure, respectively. This data is important for epidemiological research, risk communication and risk management, but also for future dosimetry and modelling studies. Future research understanding auto-induced DL and UL exposure from more realistic case scenarios remains necessary for a better characterization of the exposure levels. Future research will consist of the application of the proposed protocol in various countries and the comparison of the exposure values.

Open access paper: https://www.sciencedirect.com/science/article/pii/S001393512402454X

--

Electromagnetic intensity investigation of emitted non-ionizing radiation from base transcriptive stations in the urban region of southern Iran

Abbasi F, Badeenezhad A, Abouee E, Shademanpour Z, Janghorban F, Janatshoar H, Naserpour M, Mohammadpour A. Electromagnetic intensity investigation of emitted non-ionizing radiation from base

transcriptive stations in the urban region of southern Iran. Int J Environ Health Res. 2024 Dec 2:1-12. doi: 10.1080/09603123.2024.2435484. Epub ahead of print. PMID: 39620927.

Abstract

Their monitoring in urban regions is essential for policymakers and the population. In this study, the electromagnetic intensity (EMI) was measured around 30 stations of base transcriptive stations (BTS) at both distances lower than 20m and higher than 200m using real-time equipment SMP2-dual in summer and winter. Results have shown that EMI in summer (range: 2-6500 mW/m2) was more than in winter (1.5-5000), and the intensity of about 93% of samples exceeded WHO standards. EMI has consistently decreased with the increasing distance from BTS. There was a negative correlation between the temperature and EMI in summer and humidity and EMI for both distances. The mapping of EMI depicted the highest value of EMI across the central region from south to eastern north at a distance lower than 20 m in winter. The clustering of the EMI in this region was influenced by the geographical location of BTS.

https://pubmed.ncbi.nlm.nih.gov/39620927/

--

Effect of microwave radiation on adult neurogenesis and behavior of prenatally exposed rats

Popovičová A, Račeková E, Martončíková M, Fabianová K, Raček A, Žideková M. Effect of microwave radiation on adult neurogenesis and behavior of prenatally exposed rats. IBRO Neuroscience Reports, 17:234-244, 2024. doi: 10.1016/j.ibneur.2024.08.007.

Abstract

Postnatal neurogenesis appears to be highly sensitive to environmental factors, including microwave electromagnetic radiation (MWR). Here, we investigated the impact of MWR during intrauterine development on juvenile and adult neurogenesis in the rostral migratory stream (RMS) and the dentate gyrus of the hippocampus in the rat brain, as well as its effect on animal behavior. Female rats were exposed to MWR at a frequency of 2.45 GHz for 2 hours daily throughout pregnancy. The offspring of irradiated mothers survived to either juvenile age or adulthood. The brains of the rats were subjected to morphological analysis, assessing cell proliferation and death in both neurogenic regions. In the RMS, the differentiation of nitrergic neurons was also investigated. The effect of MWR on behavior was evaluated in rats surviving to adulthood. Prenatal MWR exposure caused significant changes in the number of proliferating and dying cells, depending on the age of the animals and the observed neurogenic region. In addition, MWR attenuated the maturation of nitrergic neurons in the RMS in both juvenile and adult rats. Morphological alterations in neurogenesis were accompanied by changes in animals' behavior. Affected neurogenesis and changes in animal behavior suggest a high sensitivity of the developing brain to MWR.

Open access paper: https://www.sciencedirect.com/science/article/pii/S2667242124000782

--

The histological investigation of the effects of electromagnetic radiation on rat ovaries

Kartal B, Alimoğulları E, Akkurt G, Alimogulları M, Çaylı S. The histological investigation of the effects of electromagnetic radiation on rat ovaries. J Mol Histol. 2024 Dec 4;56(1):29. doi: 10.1007/s10735-024-10319-w.

Abstract

People are now exposed to higher levels of electromagnetic radiation (EMR) due to the widespread use of mobile phones in recent years. The possible effects of this exposure on human health are related to EMR. It has been suggested that exposure to EMR has serious effects on reproduction. The study aimed to investigate the impact of exposure to EMR (4.5 GB; 2600 MHz) emitted by mobile phones on rat ovaries. 18 adult female Wistar albino rats were used in the study, and the animals were divided into three groups (n = 6): control, stand-by, and dialing. For 8 weeks, the experimental groups were subjected to 4.5 GB EMR at 2600 MHz while on standby and making 10-min calls every hour. The rats in the control group received no exposure. Hematoxylin-eosin (H&E) staining of ovarian tissues was performed for histomorphological examinations. Additionally, immunoexpression of autophagy-related protein Beclin-1, apoptosis marker Caspase-3, ovarian reserve marker FSH, and oxidative stress marker iNOS were investigated in the rat ovaries. Microscopic examinations showed follicular degeneration in the ovaries of the rats in the stand-by and dialing groups. The immunoexpression of Beclin-1, Caspase-3, FSH, and iNOS was detected in granulosa cells and the corpus luteum in ovarian tissues obtained from the two EMR-exposed groups. There was a significant increase in the immunoexpression of Beclin-1 and Caspase-3 in the dialing group compared to the other two groups. Additionally, the iNOS and FSH expressions were increased in both EMR exposure groups compared to the control. Our results suggest that EMR exposure harms the ovaries, and autophagy and apoptosis are involved in this process.

https://pubmed.ncbi.nlm.nih.gov/39630206/

--

Exposure to a 0.9-GHz electromagnetic field on postnatal days 21-45 may trigger the renin-angiotensin system in male rat: a histological and biochemical study

Keleş Aİ, Kaya H, Keleş G, Erol HS, Mercantepe T, Odaci E. Exposure to a 0.9-GHz electromagnetic field on postnatal days 21-45 may trigger the renin-angiotensin system in male rat: a histological and biochemical study. J Mol Histol. 2024 Dec 4;56(1):22. doi: 10.1007/s10735-024-10317-y.

Abstract

The aim of this study was to examine the relationship between the renin-angiotensin system (RAS) and histological and biochemical changes occurring in the kidney tissue of male rats exposed to a 0.9 GHz electromagnetic field (EMF). Twelve male rats aged 21 days were randomly assigned to control (C-Gr) and EMF (EMF-Gr) groups. No procedure was performed on C-Gr, while the EMF-Gr rats were exposed to a 0.9 GHz EMF on postnatal days 21-45 (one hour a day for 25 days). Tissues were removed at the end of the experiment and evaluated using biochemical, and histopathological methods. Increased kidney tissue volume and weight and total body weight were determined in the group exposed to EMF. Lipid peroxidation, glutathione, catalase, and superoxide dismutase also increased in the kidney tissue of the EMF-Gr rats. Histopathological evaluation revealed cortical/medullary bleeding/obstruction and widespread fibrosis, dilatation, vacuolization, and degeneration in distal and proximal tubules, decreased and atypical parietal cells, and degeneration in epithelial cells. Additionally, dilated and

degenerated glomeruli in the Malpighian body, Bowman's membrane degeneration and degeneration in the vascular pole, podocyte, pedicel and mesangial cells were also observed. As a result of exposure to EMF, oxidative stress, tissue volume and weight increased, and histopathological changes caused the formation of a pathway that triggers RAS in kidney tissues. In conclusion, long-term exposure to 0.9 GHz EMF can activate the renin-angiotensin system in the rat kidney, and we think that such activation may be associated with structural, histopathological, and biochemical changes occurring in renal tissue.

https://pubmed.ncbi.nlm.nih.gov/39627484/

--

Oxidative Stress and Energy Metabolism in Male Reproductive Damage from Single and Combined High-Power Microwave Exposure at 1.5 and 4.3Ghz

Li Y, Yao B, Men J, Pang Y, Gao J, Bai Y, Wang H, Zhang J, Zhao L, Xu X, Dong J, Li C, Peng R. Oxidative Stress and Energy Metabolism in Male Reproductive Damage from Single and Combined High-Power Microwave Exposure at 1.5 and 4.3Ghz. Reprod Toxicol. 2024 Nov 29:108759. doi: 10.1016/j.reprotox.2024.108759.

Abstract

The effect of multi-frequency electromagnetic environments on male reproduction has attracted the medical community's interest. Studies have investigated the effects and mechanisms of single-frequency microwave exposure on male reproduction, but comparative research on high-power microwave (HPM) composite and single exposure remains scarce. This study aimed to examine the effects and mechanisms of combined 1.5GHz and 4.3GHz microwave exposure on male reproduction. Male Wistar rats were exposed to 1.5GHz (L-band) and 4.3GHz (C-band) electromagnetic radiation for 15minutes. The four groups were: sham, 10mW/cm² L-band, 10mW/cm² C-band, and 5mW/cm² L-band and 5mW/cm² C-band compound. Assessments were made on the pathological structures of testes, sperm viability, serum sex hormones, oxidative stress, and energy metabolism levels after radiation. Exposure to 1.5GHz and 4.3GHz microwaves individually resulted in testicular tissue damage and reduced sperm quality. There was little difference between the damage caused by HPM composite and single exposure. The exposed groups showed histological and ultrastructural changes, with reduced spermatozoa viability, motility parameters, and serum testosterone, luteinizing hormone, follicle-stimulating hormone, and serum inhibin-B on days 1 and 7 after exposure. These tended to recover partially by day 14. Adenosine triphosphate content and lactate dehydrogenase and succinate dehydrogenase activities in the exposed testicular tissue decreased, corresponding to decreased superoxide dismutase activity and increased malondialdehyde content. Both single and combined exposure to L- and C-band HPM affect the male reproductive system. Exposure to single and compound HPM shows no significant difference in risks, with oxidative stress and energy metabolism disturbances playing key roles.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0890623824002260?via%3Dihub

--

Impact of exposure to extremely low-frequency magnetic fields on blood pressure, heart rate variation and disturbance to quality of sleep on industrial workers in Korea

Amila Madhushanka Weerasinghe SDNAM, Liyanage S, Rajitha Kawshalya MAD, Hong SC. Impact of exposure to extremely low-frequency magnetic fields on blood pressure, heart rate variation and disturbance to quality of sleep on industrial workers in Korea. Int J Occup Saf Ergon. 2024 Nov 23:1-8. doi: 10.1080/10803548.2024.2413816.

Abstract

Objectives: This study investigates the potential health risks associated with exposure to extremely low-frequency magnetic fields (ELF-MFs), focusing on the impact on sleep quality. The hypothesis is that increasing ELF-MF exposure will correlate with elevated blood pressure (BP), altered heart rate (HR) dynamics and poorer sleep quality among occupational workers in Korea.

Methods: The study involved 85 participants. Assessment of ELF-MF exposure was carried out continuously for 24 h in the two groups. Measurements of BP parameters, HR and sleep quality were conducted accordingly.

Results: The study found significant ELF-MF exposure levels in both working and sleeping times for industrial factory workers (0.19 ± 0.38 and 0.17 ± 0.23 μT, $p < 0.05$). Long-term exposure among industrial workers may contribute to a significant difference in systolic blood pressure (SBP) (125.61 ± 9.60 mm Hg, $p < 0.05$). Additionally, a significant difference in deep sleep analysis was observed between the high-exposure and low-exposure groups (67.13 ± 31.15 min, $p < 0.05$).

Conclusion: The hypothesis was confirmed, indicating possible effects of high ELF-MF exposure on SBP and deep sleep. However, further research on long-term exposure and its association with sleep quality disturbances is needed for validation

Open access paper: https://www.tandfonline.com/doi/full/10.1080/10803548.2024.2413816

--

Biological issue of electromagnetic fields and waves

Koyama S. Biological issue of electromagnetic fields and waves. Chapter 12 in Shinohara N. Wireless Power Transfer Technologies: Theory and Technologies (2nd edition). Aug 6, 2024.

Abstract

In the past few decades, the use of electromagnetic fields (EMF) has become prevalent throughout the world, and we no longer intend to stop using this technology. The widespread use of electromagnetic sources in daily life has induced an inevitable exposure to EMF. Recently, the rapid introduction of EMF has witnessed an upsurge in the use of EMF for telecommunication. Mobile phone use has become ubiquitous. Wireless communication devices emit nonionizing electromagnetic radiofrequency (RF) fields in the range of 300 MHz to 300 GHz, raising public concern regarding the increasing use of mobile phones and their potential health-related risks. The intermediate frequency (IF) EMF (300 Hz to 10 MHz) is also now widely used for wireless power transmission (WPT) or domestic kitchen appliances such as induction heating (IH) cooking. Unfortunately, although plenty of research has been conducted on RF, little information exists on the potential health effects associated with the exposure to IF magnetic

fields. With growing concerns regarding the potential health hazards, it has become necessary to investigate the risks of IF magnetic fields in more detail.

12.5 Conclusions on IF and RF studies

A handful of cellular studies exist on IF exposure. The usage of IF is now spreading all over the world, and more devices are using this frequency. Although the research on IF is urgent, so far it seems that the low-power intensity of IF might not induce any adverse effects on the cellular response. However, no general conclusions can be drawn on this frequency range.

There are still contradictory results about the effects of exposure to RF on the cellular response. However, RF energy does not directly cleave intercellular DNA since most genotoxicity studies have indicated negative results. Some positive data at high SARs seem to relate mainly thermal effect by exposure to RF. Several interesting results of HSP and apoptosis should be validated shortly. Further studies on RF effects using improved biotechnological methods are warranted for precise conclusions.

https://doi.org/10.1049/PBPO256E_ch12

--

Fifty-hertz magnetic fields induce DNA damage through activating mPTP associated mitochondrial permeability transition in senescent human fetal lung fibroblasts

Sun C, Wang S, Zhang J, Zhou X, Zhu T, Mao G. Fifty-hertz magnetic fields induce DNA damage through activating mPTP associated mitochondrial permeability transition in senescent human fetal lung fibroblasts. Biophysical Chemistry. Volume 318, 2025, doi: 10.1016/j.bpc.2024.107367.

Abstract

With the rapid development and using of electromagnetic technology, artificial electromagnetic fields (EMFs) have become an emerging environmental factor in our daily life. Extremely-low-frequency (ELF) magnetic fields (MFs), generally generated by power lines and various electric equipment, is one of the most common EMFs in the environment which were concerned for the potential impact on human health. Base on limited evidence, ELF-MFs have been classified as possible carcinogen to human by International Agency for Research on Cancer (IARC), but the mechanisms have not been fully elucidated. Senescent cells are a group of special cells, characterized by cell cycle arrest, senescence-associated secretory phenotype (SASP), accumulation of macromolecular damage, and metabolic disturbance, play important role in fetal development, tissue aging, and even carcinogenesis. Thus, EMFs may promote carcinogenesis by affecting senescent cells, however, there are few studies. In this study, we found that exposure to 50 Hz MFs at 1.0 mT for 24 h could induce significant DNA damage in senescent but not non-senescent human fetal lung fibroblast suggested that senescent cells are more sensitive to 50 Hz MFs on DNA damage, and further results revealed that reactive oxygen species (ROS) generation mediated by mitochondrial permeability transition pore (mPTP) activation play critical role in this process. Our results indicated that cellular senescence can lead to cell sensitivity to the DNA damage effect of 50 Hz MFs, however, whether this play important role in mediating the carcinogenesis of EMFs await further study.

https://www.sciencedirect.com/science/article/pii/S0301462224001960

--

Effects of electric fields on the release and content of extracellular vesicles

Wang Y, Worrell GA, Wang HL. Effects of electric fields on the release and content of extracellular vesicles. J Extracell Biol. 2024 Nov 29;3(11):e70018. doi: 10.1002/jex2.70018.

Abstract

Extracellular vesicles (EVs) are small membrane-bound structures that originate from various cell types and carry molecular cargos to influence the behaviour of recipient cells. The use of EVs as biomarkers for diagnosis and as delivery vehicles for treatment in a wide range of human disease is a rapidly growing field in research and clinical practice. We hypothesized that electric fields (EFs) could influence the release and content of EVs. To examine this hypothesis, we developed a specialized bioreactor enabling cells to thrive in a three-dimensional setting, replicating in-vivo conditions amidst programmable EF environments. We established a three-step EV purification protocol to achieve high-density production of EVs. We also performed mass spectrometry-based proteomics analysis on EV-carrying proteins and used high-resolution nanoparticle flowcytometry for single-vesicle analysis. Findings from this report suggest that electrical stimulation, employing physiologically relevant amplitudes typical in therapeutic deep brain stimulation, influences the release of EVs and their cargo content in a frequency-dependent fashion. This conclusion could carry significant implications for both fundamental biological understanding and medical advancements. First, it raises an intriguing question about how the endogenous electrical activity of neuronal and other cellular assemblies influence the production and composition of EVs. Second, it reveals a novel underlying mechanism of how therapeutic electrical stimulations can modulate EVs and treat human brain disorders. Third, it provides a novel approach to utilize electrical stimulation for generating desired EV cargos in a programmable setting.

Open access paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC11605478/

--

Extremely low frequency magnetic fields (ELF-MF) in Switzerland: From exposure monitoring to daily exposure scenarios

Loizeau N, Haas D, Zahner M, Stephan C, Schindler J, Gugler M, Fröhlich J, Ziegler T, Röösli M. Extremely low frequency magnetic fields (ELF-MF) in Switzerland: From exposure monitoring to daily exposure scenarios. Environment International. 2024. doi: 10.1016/j.envint.2024.109181.

Abstract

Exposure to extremely low frequency magnetic fields (ELF-MF) is ubiquitous in our daily environment. This study aims to provide a comprehensive overview of the ambient ELF-MF exposure in Switzerland and presents a novel environmental exposure matrix for exposure assessment and risk communication. Magnetic flux density levels (μT) were measured using a portable exposimeter carried in a backpack for the main ELF sources: railway power (16.7 Hz), domestic power (50 Hz), and tram ripple current (300 Hz). We collected ELF-MF levels between 2022 and 2024 in various environments representative of

the Swiss population: 300 outdoor areas (e.g. city centres, residential areas), 245 public spaces (e.g. train stations, schools), 348 transport journeys (e.g. train, cars), and in 59 homes (e.g. bedrooms, living rooms). Over all environments, the highest ELF-MF exposure levels were measured in train stations (median: 0.48 µT), trains (median: 0.40 µT), and in living rooms near (<200 m) highest voltage lines of 220 kV and 380 kV (median: 0.37 µT). ELF-MF median levels measured two years apart showed high Pearson correlation coefficients in the same 150 outdoor areas (r = 0.88) and 86 public spaces (r = 0.87), without any significant changes. All measurements are well below the Swiss ambient regulatory limit based on the ICNIRP 1998 guidelines (median: 0.2 %). Finally, we derived an environmental exposure matrix and modelled 27 daily time-weighted average ELF-MF exposure scenarios by combining typical time spent at home, work and transport environments. People who do not live near highest voltage lines or work in highly exposed environments are typically exposed to less than 0.3 µT on average, while those who do are likely to exceed this level. This novel environmental exposure matrix is a useful tool for public communication and agent-based exposure modelling for future epidemiological research.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024007670

--

Biophysical mechanism of animal magnetoreception, orientation and navigation

Panagopoulos DJ, Karabarbounis A, Chrousos GP. Biophysical mechanism of animal magnetoreception, orientation and navigation. Sci Rep. 2024 Dec 3;14(1):30053. doi: 10.1038/s41598-024-77883-9.

Abstract

We describe a biophysical mechanism for animal magnetoreception, orientation and navigation in the geomagnetic field (GMF), based on the ion forced oscillation (IFO) mechanism in animal cell membrane voltage-gated ion channels (VGICs) (IFO-VGIC mechanism). We review previously suggested hypotheses. We describe the structure and function of VGICs and argue that they are the most sensitive electromagnetic sensors in all animals. We consider the magnetic force exerted by the GMF on a mobile ion within a VGIC of an animal with periodic velocity variation. We apply this force in the IFO equation resulting in solution connecting the GMF intensity with the velocity variation rate. We show that animals with periodic velocity variations, receive oscillating forces on their mobile ions within VGICs, which are forced to oscillate exerting forces on the voltage sensors of the channels, similar or greater to the forces from membrane voltage changes that normally induce gating. Thus, the GMF in combination with the varying animal velocity can gate VGICs and alter cell homeostasis in a degree depending, for a given velocity and velocity variation rate, on GMF intensity (unique in each latitude) and the angle between velocity and GMF axis, which determine animal position and orientation.

Open access paper: https://www.nature.com/articles/s41598-024-77883-9

--

Relationship between radiofrequency-electromagnetic radiation from cellular phones and brain tumor: meta-analyses using various proxies for RF-EMR exposure-outcome assessment

My Note: The journal *Environmental Health* published a systematic review and meta-analytic study on cell phone use and brain tumor risk by Moon et al. (2024). The study found significantly elevated risks

for three types of brain tumors when examining tumors on the side of the head where cell phones were held and for heavy, long-term cell phone use.

Six other systematic reviews and meta-analyses of case-control studies published in peer-reviewed journals since 2016 have also found significant associations between heavy, long-term cellphone use and brain tumor risk (Wang & Guo, 2016; Bortkiewicz et al., 2017; Carlberg & Hardell, 2017; Prasad et al., 2017, Yang et al., 2017; Choi et al., 2020).

These seven peer-reviewed meta-analytic studies contradict the conclusion of the recent WHO systematic review conducted by Karipidis et al. (2024) that there is no evidence cell phone use causes brain cancer. To learn about serious problems with the WHO systematic reviews on the health effects of radiofrequency radiation see:

Biased WHO-commissioned review claims no cancer link to cellphone use

WHO Radiofrequency EMF Health Risk Assessment Monograph (EHC series)

References

Wang & Guo (2016). Meta-analysis of association between mobile phone use and glioma risk. *J Cancer Research Therapy* http://bit.ly/2o1dVcn

Bortkiewicz et al (2017). Mobile phone use and risk of intracranial tumors and salivary gland tumors - A meta-analysis. *Int J Occ Med Envir Health*. http://bit.ly/2nVJC5d

Carlberg & Hardell (2017). Evaluation of mobile phone and cordless phone use and glioma risk using the Bradford Hill viewpoints from 1965 on association or causation. *Biomed Res Int*. http://bit.ly/2WwBX1K

Prasad et al (2017). Mobile phone use and risk of brain tumours: a systematic review of association between study quality, source of funding, and research outcomes. *Neurol Sci*. http://bit.ly/2Xxp83P

Yang et al (2017). Mobile phone use and glioma risk: A systematic review and meta-analysis. *PLOS One*. https://bit.ly/3U0kafd

Choi, Moskowitz, et al (2020). Cellular phone use and risk of tumors: Systematic review and meta-analysis. *Int J Envir Res Public Health*. https://doi.org/10.3390/ijerph17218079.

Karipidis et al. (2020). The effect of exposure to radiofrequency fields on cancer risk in the general and working population: A systematic review of human observational studies - Part I: Most researched outcomes. *Environ Int.* https://doi.org/10.1016/j.envint.2024.108983

--

Moon J, Kwon J, Mun Y. Relationship between radiofrequency-electromagnetic radiation from cellular phones and brain tumor: meta-analyses using various proxies for RF-EMR exposure-outcome assessment. *Environ Health* **23**, 82 (2024). https://doi.org/10.1186/s12940-024-01117-8.

Abstract

Introduction

The authors conducted meta-analyses regarding the association between cellular and mobile phone use and brain tumor development by applying various radiofrequency-electromagnetic radiation (RF-EMR) exposure subcategories. With changing patterns of mobile phone use and rapidly developing Wireless

Personal Area Network (WPAN) technology (such as Bluetooth), this study will provide insight into the importance of more precise exposure subcategories for RF-EMR.

Methods

The medical librarian searched MEDLINE (PubMed), EMBASE, and the Cochrane Library until 16 December 2020.

Results

In these meta-analyses, 19 case-control studies and five cohort studies were included. Ipsilateral users reported a pooled odds ratio (OR) of 1.40 (95% CI 1.21–1.62) compared to non-regular users. Users with years of use over 10 years reported a pooled OR of 1.27 (95% CI 1.08–1.48). When stratified by each type of brain tumor, only meningioma (OR 1.20 (95% CI 1.04–1.39)), glioma (OR 1.45 (95% CI 1.16–1.82)), and malignant brain tumors (OR 1.93 (95% CI 1.55–2.39)) showed an increased OR with statistical significance for ipsilateral users. For users with years of use over 10 years, only glioma (OR 1.32 (95% CI 1.01–1.71)) showed an increased OR with statistical significance. When 11 studies with an OR with cumulative hours of use over 896 h were synthesized, the pooled OR was 1.59 (95% CI 1.25–2.02). When stratified by each type of brain tumor, glioma, meningioma, and acoustic neuroma reported the pooled OR of 1.66 (95% CI 1.13–2.44), 1.29 (95% CI 1.08–1.54), and 1.84 (95% CI 0.78–4.37), respectively. For each individual study that considered cumulative hours of use, the highest OR for glioma, meningioma, and acoustic neuroma was 2.89 (1.41–5.93) (both side use, > 896 h), 2.57 (1.02–6.44) (both side use, > 896 h), and 3.53 (1.59–7.82) (ipsilateral use, > 1640 h), respectively. For five cohort studies, the pooled risk ratios (RRs) for all CNS tumors, glioma, meningioma, and acoustic neuroma, were statistically equivocal, respectively. However, the point estimates for acoustic neuroma showed a rather increased pooled RR for ever-use (1.26) and over 10 years of use (1.61) compared to never-use, respectively.

Excerpts

“In consideration of these rapidly changing mobile phone technologies, the currently used proxies for RF-EMR exposure assessment are crude and insufficient to clarify the relationship between RF-EMR exposure from cell phones and brain tumor incidence.... The usual exposure measures, such as the years of mobile phone use, the cumulative duration of calls, and the number of calls per week, are rough indicators of mobile phone use.... the authors conducted a series of meta-analyses and subgroup analyses using various exposure measuring categories, from crude to more precise ones. In consideration of crude exposure classifications used in previous meta-analyses, this study will give insight into the importance of more precise exposure subcategories in investigating this topic."

"... the authors analyzed the risk of bias regarding selection and recall bias for the amount of cell phone use and misclassification and recall bias for ipsilateral/contralateral use. A major reason was that typical risk of bias rating tools such as the National Toxicology Program Office of Health Assessment and Translation Risk of Bias rating tool (NTP OHAT RoB rating tool, Supplementary material B) were not appropriate for assessing individual studies regarding this topic." [Note: The WHO systematic reviews relied on this rating tool.]

"With the conversion from 2G cellular phones through 3G and 4G mobile phones to current 5G mobile phones, transmission of large data became possible. With the introduction of 3G technology, all aspects

of our society and daily lives have changed drastically. Currently, we are using mobile phones nearly continuously and putting mobile phones near our bodies even when we are not using them. For example, if people use their mobile phone for morning-alarming purposes, they might put their mobile phone near the bed, sometimes even beside their head, all night. These changed patterns of mobile phone use could increase exposure to RF-EMR from cellular and mobile phones. Therefore, precise exposure assessment for RF-EMR from mobile phones would become more complex in future studies."

"The results of cumulative meta-analyses according to precision indicated that the pooled OR was biased downwards with the addition of studies with lower precision. This indicates that the results of studies with a lower precision should be interpreted cautiously."

"Because brain tumors require a latency period to develop [29], an accurate assessment of brain tumor risk associated with RF-EMR exposure requires a long observation span. However, each included study did not consider a sufficient latency period in their study design. This could have led to a possible underestimation of brain tumor risk. Future studies with long observation spans might resolve this problem."

"In this meta-analysis, as the applied exposure subcategories became more concrete, the pooled ORs showed more increased values with statistical significance. Even though the meta-analysis of cohort studies showed statistically equivocal pooled effect estimates, (i) as the number of included studies increases and (ii) as the applied exposure subcategory becomes more concrete, the pooled RRs could show a different aspect in future studies. Furthermore, changing patterns of mobile phone use and increasing use of earphones or headphones with WPAN technology should be sufficiently considered in future studies. Relatively short observation spans for brain tumor incidence and age of starting exposure and brain tumor diagnosis should also be considered in future studies. Previous studies that adjusted for selection and recall bias for the amount of cellphone use and misclassification and recall bias for ipsilateral/contralateral use showed possible underestimations of previous risk estimates. Future studies should try to adjust for these biases in their study design."

Open access paper: https://ehjournal.biomedcentral.com/articles/10.1186/s12940-024-01117-8

Supplementary material: https://static-content.springer.com/esm/art%3A10.1186%2Fs12940-024-01117-8/MediaObjects/12940_2024_1117_MOESM1_ESM.docx

--

Electromagnetic fields of cellular communication as risk factors able to produce negative effects on the central nervous system of children and adolescents (review)

Khorseva NI, Grigoriev PE. Electromagnetic fields of cellular communication as risk factors able to produce negative effects on the central nervous system of children and adolescents (review). Part 1. Modeling. Parameters of electroencephalography and sensorimotor reactions. Health Risk Analysis, 2024, no. 2, pp. 162–169. doi: 10.21668/health.risk/2024.2.15.eng.

Abstract

It is quite relevant to investigate possible outcomes of exposure to radio frequency electromagnetic fields (RF EMF) since contemporary children and adolescents have become active users of the most

advanced technologies. They are especially susceptible to electromagnetic factors; therefore, it is necessary to have a proper insight into outcomes of such exposures for the body.

The central nervous system (CNS) is one of the main targets under exposure to RF EMF. In most cases, users hold mobile phones close to their heads thereby directly exposing their brains to RF EMF.

As the analysis of literature data has shown, there are few studies in this area; however, proposed options for assessing the impact of RF EMF on children and adolescents are very diverse.

This part of the review focuses on various types of modeling. These are not only phantom, voxel models or the finite difference method but also new approaches such as distribution matrices, Monte Carlo simulations and an integrated radio frequency model based on the results of magnetic resonance imaging of the brain and other methods.

The review provides the results obtained by investigating encephalography under exposure to RF EMF created by mobile communication devices. They are rather contradictory; however, changes in the bioelectrical activity of the brain are detected in most cases, in particular, a decrease in the alpha rhyme.

Since the characteristics of sensorimotor reactions quite clearly reflect the power relations in the cerebral cortex, we analyzed changes in the parameters of simple auditory-motor and visual-motor reactions in children and adolescents who were mobile communication users. In addition, the review covers the results of changes in working capacity, fatigue, the duration of an individual minute and the reproduction of a given rhythm.

Open access paper: health-risk-analysis-2024-2-15.pdf

Assessing the risk of negative effects produced by electromagnetic fields of cellular communication on the central nervous system of children and adolescents (review). Part 2. Indicators of cognitive processes

Khorseva NI, Grigoriev PE. Assessing the risk of negative effects produced by electromagnetic fields of cellular communication on the central nervous system of children and adolescents (review). Part 2. Indicators of cognitive processes. Health Risk Analysis, 2024, no. 3, pp. 146–154. doi: 10.21668/health.risk/2024.3.15.eng

Abstract

This paper continues the authors' review that dwells on modeling radiofrequency electromagnetic fields (RF EMF) and results obtained by measuring electroencephalography indicators, sensorimotor reactions, fatigue, work capacity, duration of an individual minute and the reproduction of a given rhythm in children and adolescents.

Health risk assessment is always based on data obtained by either laboratory tests or epidemiological studies. This paper analyses publications that describe effects of RF EMF exposure, including Wi-Fi, on cognitive processes in children and adolescents as well as methodical approaches to investigating this exposure. However, there are few such studies; in particular, effects produced by Wi-Fi exposure on cognitive indicators of adolescents aged 14–17 years, were found only in two publications.

Literature analysis has established that research findings do not always give an unambiguous estimation of RF EMF effects. The review covers the reasons for ambiguous interpretation of research results: a variable range of test-systems used for investigating indicators of cognitive processes; simultaneous analysis of single exposures including descriptions of ‘effect of improvement’ in indicators; changes in cognitive indicators registered for a group of children and adolescents in a wide age range.

Nevertheless, most results give evidence of negative changes in attention and memory of children and adolescents. Given that, longitudinal studies are becoming especially relevant since they estimate changes in various indicators in dynamics, including those induced by changes in mobile phone use. The review highlights the relevance of comprehensive investigations with their focus on health outcomes of RF EMF exposure intrinsic to 5G technologies considering their global implementation.

Open access paper: https://journal.fcrisk.ru/eng/2024/3/15

Russian version: https://journal.fcrisk.ru/sites/journal.fcrisk.ru/files/upload/article/787/health-risk-analysis-2024-3-15.pdf

--

Autonomous nervous system responses to environmental-level exposure to 5G's first deployed band (3.5 GHz) in healthy human volunteers

Jamal L, Michelant L, Delanaud S, Hugueville L, Mazet P, Lévêque P, Baz T, Bach V, Selmaoui B. Autonomous nervous system responses to environmental-level exposure to 5G's first deployed band (3.5 GHz) in healthy human volunteers. Exp Physiol. 2024 Oct 15. doi: 10.1113/EP092083.

Abstract

Following the global progressive deployment of 5G networks, considerable attention has focused on assessing their potential impact on human health. This study aims to investigate autonomous nervous system changes by exploring skin temperature and electrodermal activity (EDA) among 44 healthy young individuals of both sexes during and after exposure to 3.5 GHz antenna-emitted signals, with an electrical field intensity ranging from 1 to 2 V/m. The study employed a randomized, cross-over design with triple-blinding, encompassing both 'real' and 'sham' exposure sessions, separated by a maximum interval of 1 week. Each session comprised baseline, exposure and postexposure phases, resulting in the acquisition of seven runs. Each run initiated with a 150 s segment of EDA recordings stimulated by 10 repeated beeps. Subsequently, the collected data underwent continuous decomposition analysis, generating specific indicators assessed alongside standard metrics such as trough-to-peak measurements, global skin conductance and maximum positive peak deflection. Additionally, non-invasive, real-time skin temperature measurements were conducted to evaluate specific anatomical points (hand, head and neck). The study suggests that exposure to 3.5 GHz signals may potentially affect head and neck temperature, indicating a slight increase in this parameter. Furthermore, there was a minimal modulation of certain electrodermal metrics after the exposure, suggesting a potentially faster physiological response to auditory stimulation. However, while the results are significant, they remain within the normal physiological range and could be a consequence of an uncontrolled variable. Given the preliminary nature of this pilot study, further research is needed to confirm the effects of 5G exposure.

Open access paper: https://physoc.onlinelibrary.wiley.com/doi/10.1113/EP092083

--

Does radiofrequency radiation impact sleep? A double-blind, randomised, placebo-controlled, crossover pilot study

Bijlsma N, Conduit R, Kennedy G, Cohen M. Does radiofrequency radiation impact sleep? A double-blind, randomised, placebo-controlled, crossover pilot study. Front Public Health. 2024 Oct 29;12:1481537. doi: 10.3389/fpubh.2024.1481537.

Abstract

The most common source of Radiofrequency Electromagnetic Field (RF-EMF) exposures during sleep includes digital devices, yet there are no studies investigating the impact of multi-night exposure to electromagnetic fields emitted from a baby monitor on sleep under real-world conditions in healthy adults. Given the rise in the number of people reporting to be sensitive to man-made electromagnetic fields, the ubiquitous use of Wi-Fi enabled digital devices and the lack of real-world data, we investigated the effect of 2.45 GHz radiofrequency exposure during sleep on subjective sleep quality, and objective sleep measures, heart rate variability and actigraphy in healthy adults. This pilot study was a 4-week randomised, double-blind, crossover trial of 12 healthy adults. After a one-week run-in period, participants were randomised to exposure from either an active or inactive (sham) baby monitor for 7 nights and then crossed over to the alternate intervention after a one-week washout period. Subjective and objective assessments of sleep included the Pittsburgh Insomnia Rating Scale (PIRS-20), electroencephalography (EEG), actigraphy and heart rate variability (HRV) derived from electrocardiogram. Sleep quality was reduced significantly ($p < 0.05$) and clinically meaningful during RF-EMF exposure compared to sham-exposure as indicated by the PIRS-20 scores. Furthermore, at higher frequencies (gamma, beta and theta bands), EEG power density significantly increased during the Non-Rapid Eye Movement sleep ($p < 0.05$). No statistically significant differences in HRV or actigraphy were detected. Our findings suggest that exposure to a 2.45 GHz radiofrequency device (baby monitor) may impact sleep in some people under real-world conditions; however, further large-scale real-world investigations with specified dosimetry are required to confirm these findings.

Open access paper:

https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2024.1481537/full

--

A review on the consequences of molecular & genomic alterations following exposure to EMF: Remodeling of neuronal network & cognitive changes

Abtin S, Seyedaghamiri F, Aalidaeijavadi Z, Farrokhi AM, Moshrefi F, Ziveh T, Zibaii MI, Aliakbarian H, Rezaei-Tavirani M, Haghparast A. A review on the consequences of molecular and genomic alterations following exposure to electromagnetic fields: Remodeling of neuronal network and cognitive changes. Brain Research Bulletin, Volume 217, 2024. doi: 10.1016/j.brainresbull.2024.111090.

Abstract

The use of electromagnetic fields (EMFs) is essential in daily life. Since 1970, concerns have grown about potential health hazards from EMF. Exposure to EMF can stimulate nerves and affect the central nervous system, leading to neurological and cognitive changes. However, current research results are often vague and contradictory. These effects include changes in memory and learning through changes in neuronal plasticity in the hippocampus, synapses and hippocampal neuritis, and changes in metabolism and neurotransmitter levels. Prenatal exposure to EMFs has negative effects on memory and learning, as well as changes in hippocampal neuron density and histomorphology of hippocampus. EMF exposure also affects the structure and function of glial cells, affecting gate dynamics, ion conduction, membrane concentration, and protein expression. EMF exposure affects gene expression and may change epigenetic regulation through effects on DNA methylation, histone modification, and microRNA biogenesis, and potentially leading to biological changes. Therefore, exposure to EMFs possibly leads to changes in cellular and molecular mechanisms in central nervous system and alter cognitive function.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0361923024002247

--

Impact of non-ionising radiation of male fertility: a systematic review

Motchidlover L, Sari-Minodier I, Sunyach C, Metzler-Guillemain C, Perrin J. Impact of non-ionising radiation of male fertility: a systematic review. French Journal of Urology. 35(1), 2025, doi: 10.1016/j.fjurol.2024.102800.

Abstract

Exposure to non-ionizing radiation has become inevitable because people cannot escape sources of electromagnetic fields, such as Wi-Fi or cell phones. Among the mechanisms mentioned, the energy emitted by this non-ionizing radiation could cause heating which would have harmful effects on semen quality. The objective of our study was to carry out a systematic review of the literature concerning the impact of exposure to non-ionizing radiation from mobile phones (or other sources) on sperm parameters. We selected 12 studies: the majority of in vivo studies in humans and in vitro studies in animals report a significant impact on sperm count, mobility and vitality. Mobility and vitality seem to be the parameters most regularly impacted by exposure to non-ionizing radiation. Additional studies are necessary to complete this study in order to deepen knowledge with new generations of mobile phones which can raise health concerns.

https://www.sciencedirect.com/science/article/pii/S2950393024002675

--

Exposure to Radiofrequency Electromagnetic Fields Enhances Melanin Synthesis by Activating the P53 Signaling Pathway in Mel-Ab Melanocytes

Kim JH, Kang D-J, Seok JY, Kim M-H, Kim D-S, Jeon S-B, Choi H-D, Moon JI, Kim N, Kim HR. Exposure to Radiofrequency Electromagnetic Fields Enhances Melanin Synthesis by Activating the P53 Signaling Pathway in Mel-Ab Melanocytes. *International Journal of Molecular Sciences*. 2024; 25(22):12457. https://doi.org/10.3390/ijms252212457

Abstract

The skin is the largest body organ that can be physiologically affected by exposure to radiofrequency electromagnetic fields (RF-EMFs). We investigated the effect of RF-EMFs on melanogenesis; Mel-Ab melanocytes were exposed to 1760 MHz radiation with a specific absorption rate of 4.0 W/kg for 4 h/day over 4 days. Exposure to the RF-EMF led to skin pigmentation, with a significant increase in melanin production in Mel-Ab melanocytes. The phosphorylation level of cAMP response element binding protein (CREB) and the expression of microphthalmia-associated transcription factor (MITF), which regulate the expression of tyrosinase, were significantly increased in Mel-Ab after RF-EMF exposure. Interestingly, the expression of tyrosinase was significantly increased, but tyrosinase activity was unchanged in the RF-EMF-exposed Mel-Ab cells. Additionally, the expression of p53 and melanocortin 1 receptor (MC1R), which regulate MITF expression, was significantly increased. These results suggest that the RF-EMF induces melanogenesis by increasing phospho-CREB and MITF activity. Importantly, when Mel-Ab cells were incubated at 38 °C, the melanin production and the levels of tyrosinase significantly decreased, indicating that the increase in melanin synthesis by RF-EMF exposure is not due to a thermal effect. In conclusion, RF-EMF exposure induces melanogenesis in Mel-Ab cells through the increased expression of tyrosinase via the activation of MITF or the phosphorylation of CREB, which are initiated by the activation of p53 and MC1R.

Conclusion: In summary, we have shown that 1760 MHz RF-EMF exposure (SAR of 4.0 W/kg for 4 h/d over 4 d) could induce hyperpigmentation by activating the melanin synthesis pathway in Mel-Ab melanocytes, suggesting that skin pigmentation could be affected by RF-EMF exposure in a way that is distinct from simple thermal effects.

Open access paper: https://www.mdpi.com/1422-0067/25/22/12457

--

Genotoxicity of radiofrequency electromagnetic fields on mammalian cells in vitro: A systematic review with narrative synthesis

My note: This is not one the ten systematic reviews commissioned by the WHO. Nonetheless, this review of *in vitro* studies dismisses most evidence of adverse effects from RF-EMF.

"Genotoxicity, a key characteristic of human carcinogens (Smith et al. 2016), is one of the most investigated outcomes in experimental studies on the effects of RF-EMF (SCHEER 2023). However, none of the ten systematic reviews commissioned by the World Health Organization (WHO) in the framework of an updated RF-EMF hazard and risk assessment, deals with this topic (Verbeek et al. 2021)."

Romeo S, Sannino A, Scarfì MR, Lagorio S, Zeni O. Genotoxicity of radiofrequency electromagnetic fields on mammalian cells in vitro: A systematic review with narrative synthesis. Environment International, 2024, doi: 10.1016/j.envint.2024.109104.

Abstract

Background: Over the last decades, great concern has been raised about possible adverse effects to human health due to exposures to radiofrequency electromagnetic fields (RF-EMF, 100 kHz – 300 GHz) emitted by wireless communication technologies. In 2011 the International Agency for Research on Cancer classified RF-EMF as possibly carcinogenic to humans, highlighting that the evidence was weak

and far from conclusive. Updated systematic reviews of the scientific literature on this topic are lacking, especially for mechanistic studies.

Objectives: To perform a systematic review of the scientific literature on genotoxic effects induced by RF-EMF in in vitro experimental models. The overall aim is to assess the confidence and level of evidence of the induced effects in mammalian cell cultures.

Methods: Full details regarding the eligibility criteria, information sources, and methods developed to assess risk of bias in the included study, are reported in our published protocol (Romeo et al. 2021). The databases NCBI PubMed, Web of Science, and EMF-Portal were used as information sources (last searched on 31st December 2022). In developing the systematic review, we followed the guidelines provided by the National Toxicology Program-Office of Health Assessment and Translation (NTP-OHAT), adapted to the evaluation of in vitro studies. A narrative synthesis of the body of evidence was performed by tabulating data classified according to meaningful groups (endpoints) and sub-groups (exposure parameters). This report, abstract included, conforms to the PRISMA 2020 (Preferred Reporting Items for Systematic reviews and Meta-Analyses) guidelines.

Results: Out of 7750 unique records identified, 159 articles were eligible for inclusion. From the extracted data, we identified 1111 experiments (defined as independent specific combinations of diverse biological and electromagnetic parameters). The large majority (80%) of experiments reviewed did not show statistically significant genotoxic effects of RF-EMF exposures, and most “positive” studies were rated as of moderate to low quality, with negative ratings in the key bias domains. A qualitative evidence appraisal was conducted at the endpoint level, and then integrated across endpoints.

Discussion: To the best of our knowledge, this is the first systematic review of the scientific literature on genotoxic effects in mammalian cell cultures in relation to RF-EMF exposure, which confirms and strengthens conclusions from previous syntheses of this specific topic thanks to the use of transparently reported methods, pre-defined inclusion criteria, and formal assessment of susceptibility to bias. Limitations of the evidence included were the frequent reporting of findings in graphical display only, and the large heterogeneity of experimental data, which precluded a meta-analysis.

Conclusions: In the assessment restricted to studies reporting a significant effect of the exposure on the outcome, we reached an overall assessment of “low” confidence in the level of evidence that RF-EMF induce genotoxic effects in mammalian cells However, 80% of experiments reviewed showed no effect of RF exposure on the large majority of endpoints, especially the irreversible ones, independently of the exposure features, level, and duration (moderate evidence of no effect). Therefore, we conclude that the analysis of the papers included in this review, although only qualitative, suggests that RF exposure does not increase the occurrence of genotoxic effects in vitro.

Framework and funding: This systematic review addresses one of the evidence streams considered in a larger systematic review of the scientific literature on the potential carcinogenicity of RF-EMF, performed by scientists from several Italian public research agencies. The project is supported by the Italian Workers’ Compensation Authority (INAIL) in the framework of the CRA with the Istituto Superiore di Sanità “BRiC 2018/06 – Scientific evidence on the carcinogenicity of electromagnetic fields”.

Excerpt

Conclusions

The main conclusion of our systematic analysis is that the certainty of evidence for genotoxic effects of RF-EMF in mammalian cell cultures is weak. No genotoxic effects of RF exposure were observed in most experiments, especially those concerning irreversible endpoints, independently of the exposure features, level, and duration (moderate evidence of no effect). In the whole, we conclude that the analysis of the papers included in this review, although only qualitative, suggests that RF exposure does not increase the occurrence of genotoxic effects *in vitro*, also considering that the absence of significant effects prevails among the studies classified in tier-1 and tier-2 categories, whereas for the tier-3 studies the trend is reversed, highlighting how methodological quality affects the studies outcome (Simko et al. 2016).

This systematic review may provide support to panels of experts involved in policy making regarding safety of exposure to RF-EMF.

Furthermore, it is possible to state that, although the number of publications on this topic has considerably increased in recent years, this has not translated into an improvement of the experimental quality. Only a small number of the reviewed studies qualified for inclusion in the tier 1 category, while most of them, including those of moderate quality, still present considerable criticalities in key experimental features (use of sham samples, adequate dosimetry, exposure to a uniformly distributed electromagnetic field, appropriateness of biological methods, presence of positive control and outcome analysis blind to the exposure). It is evident that methodological quality is demanded for future research into this field, which will aim at producing sufficiently similar and unbiased data to be aggregated in meaningful categories.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0160412024006901?via%3Dihub

--

The role of digital device use on the risk of migraine: a univariable and multivariable Mendelian randomization study

He Z, Qiu F, Yang J, Zhao M. The role of digital device use on the risk of migraine: a univariable and multivariable Mendelian randomization study. Front Neurol. 2024 Oct 30;15:1462414. doi: 10.3389/fneur.2024.1462414.

Abstract

Background: The pervasive integration of digital devices into daily life has raised concerns about their potential health impacts. This study aimed to explore the causal relationships between digital device use and the risk of migraine using Mendelian randomization (MR).

Methods: Genetic data on digital device use and migraines were sourced from large-scale genome-wide association studies conducted by the UK Biobank, the FinnGen study, and the International Headache Genetics Consortium. Univariable MR (UVMR), meta-analysis, and multivariable MR (MVMR) approaches were conducted to explore and verify the causal effects of digital device use (including mobile phone use, computer use, playing computer games, and watching television) on migraine risk.

Sensitivity analyses were conducted using Cochran's Q, MR-Egger intercept test, MR pleiotropy residual sum and outlier, MR Radial, MR Steiger, and leave-one-out methods.

Results: UVMR analyses revealed that genetically predicted mobile phone use was significantly associated with an increased risk of overall migraine (odds ratio [OR] = 2.39, *p* = 9.78e-5) and migraine without aura (MO) (OR = 2.25, *p* = 0.024). Additionally, there were significant positive associations between genetically predicted television watching and the risk of overall migraine (OR = 1.63, *p* = 2.12e-5) and MO (OR = 2.10, *p* = 4.98e-5). These results were further supported by the meta-analysis and MVMR analysis. Sensitivity analysis indicated no heterogeneity or pleiotropy.

Conclusion: This comprehensive MR study provides preliminary evidence for the causal impact of mobile phone use and television watching on the risk of migraines. Further studies are needed to explore these associations across different populations.

Open access paper:
https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2024.1462414/full

--

The Effect of Exposure to Mobile Phones on Electrical Cardiac Measurements: A Multivariate Analysis and a Variable Selection Algorithm to Detect the Relationship With Mean Changes

Alharbi N, Alassiri M. The Effect of Exposure to Mobile Phones on Electrical Cardiac Measurements: A Multivariate Analysis and a Variable Selection Algorithm to Detect the Relationship With Mean Changes. Int J Cell Biol. 2024 Oct 3;2024:7093771. doi: 10.1155/2024/7093771.

Background: The exponential growth in mobile phone usage has raised concerns about electromagnetic field (EMF) exposure and its health risks. Blood pressure and BMI, which impair heart function due to decreased adrenoreceptor responsiveness, parasympathetic tone withdrawal, and increased sympathetic activity, may further exacerbate these risks. However, the effects of radiofrequency electromagnetic (RF-EM) exposure from mobile phones on electrocardiograms (ECGs) and heart rate variability (HRV) in individuals remain unclear.

Purpose: Building upon our previous findings on HRV changes due to mobile phone proximity, this study is aimed at significantly enhancing the analytical approach used to assess the effects of mobile phones on cardiac parameters. This study exploits data from a previous study but with a different purpose. The aim of this study is twofold: (a) to examine whether exposure to mobile phones changes the five variables (P-R, QRS, QT, ST, and HR) in a multivariate manner and (b) to examine whether the blood pressure and/or the body mass index (BMI), which acts as a proxy for obesity, have an effect on the change of these five variables. For both aspects of the study, four cycles are performed.

Method: We conducted multivariate analysis on previously collected electrical cardiac measurement data from 20 healthy male subjects exposed to mobile phone EMF, with the mobile phones placed at four different body locations. The one-sample Hotelling T 2 test on the mean vector of differences was utilised instead of multiple paired t-tests. This multivariate method comprehensively analyzes data features and accounts for variable correlations, unlike multiple univariate analyses. Given our small sample size, we employed the MMPC variable selection algorithm to identify predictor variables

significantly related to mean changes.

Results: Significant alterations in ECG intervals and heart rate were noted in the subjects before and after the first EMF exposure cycle, independent of their BMI. Notably, heart rate, P-R, and QRS intervals fell postexposure while QT and ST intervals increased. These changes were influenced by variations in systolic blood pressure, with BMI showing no significant effect.

Conclusion: The observed modifications in cardiac electrical measurements due to mobile phone EMF exposure are attributed to the effects of EMF itself, with no impact from BMI on the extent of these changes.

https://regulatoryinfo.apple.com/rfexposure/iphone5,1/en

Open access paper: https://onlinelibrary.wiley.com/doi/10.1155/2024/7093771

--

Use of Mobile Phones and Radiofrequency-Emitting Devices in the COSMOS-France Cohort

Deltour I, Guida F, Ribet C, Zins M, Goldberg M, Schüz J. Use of Mobile Phones and Radiofrequency-Emitting Devices in the COSMOS-France Cohort. *International Journal of Environmental Research and Public Health*. 2024; 21(11):1514. doi: 10.3390/ijerph21111514.

Abstract

COSMOS-France is the French part of the COSMOS project, an international prospective cohort study that investigates whether the use of mobile phones and other wireless technologies is associated with health effects and symptoms (cancers, cardiovascular diseases, neurologic pathologies, tinnitus, headaches, or sleep and mood disturbances). Here, we provide the first descriptive results of COSMOS-France, a cohort nested in the general population-based cohort of adults named Constances.

Methods: A total of 39,284 Constances volunteers were invited to participate in the COSMOS-France study during the pilot (2017) and main recruitment phase (2019). Participants were asked to complete detailed questionnaires on their mobile phone use, health conditions, and personal characteristics. We examined the association between mobile phone use, including usage for calls and Voice over Internet Protocol (VoIP), cordless phone use, and Wi-Fi usage with age, sex, education, smoking status, body mass index (BMI), and handedness.

Results: The participation rate was 48.4%, resulting in 18,502 questionnaires in the analyzed dataset. Mobile phone use was reported by 96.1% (N = 17,782). Users reported typically calling 5–29 min per week (37.1%, N = 6600), making one to four calls per day (52.9%, N = 9408), using one phone (83.9%, N = 14,921) and not sharing it (80.4% N = 14,295), mostly using the phone on the side of the head of their dominant hand (59.1%, N = 10,300), not using loudspeakers or hands-free kits, and not using VoIP (84.9% N = 15,088). Individuals' age and sex modified this picture, sometimes markedly. Education and smoking status were associated with ever use and call duration, but neither BMI nor handedness was. Cordless phone use was reported by 66.0% of the population, and Wi-Fi use was reported by 88.4%.

Conclusion: In this cross-sectional presentation of contemporary mobile phone usage in France, age and sex were important determinants of use patterns.

Open access paper: https://www.mdpi.com/1660-4601/21/11/1514

--

Effects of 1800 MHz and 2100 MHz mobile phone radiation on the blood-brain barrier of New Zealand rabbits

Kizilçay AO, Tütüncü B, Koçarslan M, Gözel MA. Effects of 1800 MHz and 2100 MHz mobile phone radiation on the blood-brain barrier of New Zealand rabbits. Med Biol Eng Comput. 2024 Nov 16. doi: 10.1007/s11517-024-03238-1.

Abstract

In this study, the impact of mobile phone radiation on blood-brain barrier (BBB) permeability was investigated. A total of 21 New Zealand rabbits were used for the experiments, divided into three groups, each consisting of 7 rabbits. One group served as the control, while the other two were exposed to electromagnetic radiation at frequencies of 1800 MHz with a distance of 14.5 cm and 2100 MHz with a distance of 17 cm, maintaining a constant power intensity of 15 dBm, for a duration equivalent to the current average daily conversation time of 38 min. The exposure was conducted under non-thermal conditions, with RF radiation levels approximately ten times lower than normal values. Evans blue (EB) dye was used as a marker to assess BBB permeability. EB binds to plasma proteins, and its presence in brain tissue indicates a disruption in BBB integrity, allowing for a quantitative evaluation of radiation-induced permeability changes. Left and right brain tissue samples were analyzed using trichloroacetic acid (TCA) and phosphate-buffered solution (PBS) solutions to measure EB amounts at 620 nm via spectrophotometry. After the experiments, BBB tissue samples were collected from the right and left brains of all rabbits in the three groups and subjected to a series of medical procedures. Samples from Group 1 were compared with those from Group 2 and Group 3 using statistical methods to determine if there were any significant differences. As a result, it was found that there was no statistically significant difference in the BBB of rabbits exposed to 1800 MHz radiation, whereas there was a statistically significant difference at a 95% confidence level in the BBB of rabbits exposed to 2100 MHz radiation. A decrease in EB values was observed upon the arithmetic examination of the BBB.

https://pubmed.ncbi.nlm.nih.gov/39548043/

Excerpts

When compromised, the BBB is linked to neurological disorders such as Alzheimer's disease, stroke, and multiple sclerosis, leading researchers to develop models for in vitro studies of its mechanisms [9, 10]. However, BBB permeability can be influenced by various factors, including pathological conditions, multiple sclerosis, stroke, and diabetes, as well as exposure to ionizing and non-ionizing radiation [11]. For instance, radiofrequency (RF) radiation has been shown to increase BBB permeability, allowing normally excluded substances to penetrate the brain, potentially impacting neurological health [5]. Furthermore, ultra-wideband electromagnetic pulses have demonstrated the ability to disrupt BBB integrity in animal models [8]....

The two primary frequencies utilized in contemporary mobile telecommunications, 1800 MHz and 2100 MHz, have garnered substantial attention due to their widespread application in cellular networks worldwide [13, 14]. For example, in ref. [15], it has been demonstrated that radiation from mobile phones at 900, 1800, and 2100 MHz frequencies can increase oxidative damage in the frontal lobes of rat brain tissues, an increase in lipid peroxidation, and oxidative DNA damage. Additionally, exposure to 2100 MHz radiofrequency radiation has been shown to induce single-strand breaks in DNA. In ref. [16], it was observed and presented that exposure to 900–1800 MHz radiation caused oxidative stress in the tissues of pregnant mothers and their offspring. In ref. [17], it has been observed that radiofrequency radiation at 2100 MHz can damage the nasal septal mucosa and disrupt mucociliary clearance in rats. While numerous studies have explored the biological effects of radiofrequency radiation, the specific impact of these frequencies on BBB permeability remains an area of ongoing investigation [18,19,20]....

Our research was conducted with three distinct groups, each consisting of seven female rabbits with an average weight ranging from 2 to 5 kg. These groups were categorized as follows: with “N” signifying the number of subjects in each group: Group I, control female group (N = 7); Group II, 1800 MHz GSM group (N = 7); and Group III, 2100 MHz GSM group (N = 7)....

The exposure duration of rabbits to RF radiation was determined to be 38 min, taking into account the current average daily talk time of 19 min for mobile phone users [29]....

It is evident that various studies have investigated the impact of electromagnetic radiation at different frequencies on brain tissue. The analysis of these studies indicates that while lower frequencies may affect brain structure over a longer time frame, higher frequencies like 2.1 GHz have more immediate, statistically significant impacts, suchas compromising BBB integrity. This comparison underscores the importance of further investigating the threshold effects of different frequencies, especially those commonly used in mobile telecommunications.

--

Effects of 4G Long-Term Evolution Electromagnetic Fields on Thyroid Hormone Dysfunction and Behavioral Changes in Adolescent Male Mice

Kim HY, Son Y, Jeong YJ, Lee SH, Kim N, Ahn YH, Jeon SB, Choi HD, Lee HJ. Effects of 4G Long-Term Evolution Electromagnetic Fields on Thyroid Hormone Dysfunction and Behavioral Changes in Adolescent Male Mice. Int J Mol Sci. 2024 Oct 10;25(20):10875. doi: 10.3390/ijms252010875.

Abstract

Radiofrequency electromagnetic fields (RF-EMFs) can penetrate tissues and potentially influence endocrine and brain development. Despite increased mobile phone use among children and adolescents, the long-term effects of RF-EMF exposure on brain and endocrine development remain unclear. This study investigated the effects of long-term evolution band (LTE) EMF exposure on thyroid hormone levels, crucial for metabolism, growth, and development. Four-week-old male mice (C57BL/6) were exposed to LTE EMF (whole-body average specific absorption rate [SAR] 4 W/kg) or a positive control (lead; Pb, 300 ppm in drinking water) for 4 weeks. Subsequently, the mice underwent behavioral tests including open field, marble burying, and nest building. Blood pituitary and thyroid hormone levels, and thyroid hormone-regulating genes within the hypothalamus-pituitary-thyroid (HPT) axis were

analyzed. LTE exposure increased T3 levels, while Pb exposure elevated T3 and T4 and decreased ACTH levels. The LTE EMF group showed no gene expression alterations in the thyroid and pituitary glands, but hypothalamic *Dio2* and *Dio3* expressions were significantly reduced compared to that in the sham-exposed group. Pb exposure altered the hypothalamic mRNA levels of *Oatp1c1* and *Trh*, pituitary mRNA of *Trhr*, and *Tpo* and *Tg* expression in the thyroid. In conclusion, LTE EMF exposure altered hypothalamic *Dio2* and *Dio3* expression, potentially impacting the HPT axis function. Further research is needed to explore RF-EMF's impacts on the endocrine system.

Open access paper: https://www.mdpi.com/1422-0067/25/20/10875

--

Characterising core body temperature response of free-moving C57BL/6 mice to 1.95 GHz whole-body radiofrequency-electromagnetic fields

Sylvester E, Deng C, McIntosh R, Iskra S, Frankland J, McKenzie R, Croft RJ. Characterising core body temperature response of free-moving C57BL/6 mice to 1.95 GHz whole-body radiofrequency-electromagnetic fields. Bioelectromagnetics. 2024 Oct 14. doi: 10.1002/bem.22527.

Highlights

- Exposure of 1.95 GHz electromagnetic fields at 5 W/kg whole-body average specific absorption rate increases core body temperature by 0.4°C.
- The increased thermal energy at 3.75 W/kg is effectively managed by thermoregulation (max increase = 0.24°C).
- An extended habituation period prior to RF-EMF exposure is necessary to account for the large effect of handling on mice core body temperature.

Abstract

The present study investigated the core body temperature (CBT) response of free-moving adult male and female C57BL/6 mice, during and following a 2-h exposure to 1.95 GHz RF-EMF within custom-built reverberation chambers, using temperature capsules implanted within the intraperitoneal cavity and data continuously logged and transmitted via radiotelemetry postexposure. Comparing RF-EMF exposures (WBA-SAR of 1.25, 2.5, 3.75, and 5 W/kg) to the sham-exposed condition, we identified a peak in CBT within the first 16 min of RF-EMF exposure (+0.15, +0.31, +0.24, +0.37°C at 1.25, 2.5, 3.75, and 5 W/kg respectively; statistically significant at WBA-SAR ≥ 2.5 W/kg only), which largely dissipated for the remainder of the exposure period. Immediately before the end of exposure, only the CBT of the 5 W/kg condition was statistically differentiable from sham. Based on our findings, it is apparent that mice are able to effectively compensate for the increased thermal load at RF-EMF strengths up to 5 W/kg. In addition, the elevated CBT at the end of the exposure period in the 5 W/kg condition was statistically significantly reduced compared to the sham condition immediately after RF-EMF exposure ceased. This would indicate that measures of CBT following the end of an RF-EMF exposure period may not reflect the actual change in the CBT of mice caused by RF-EMF exposure in mice.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.22527

--

Effects of radiofrequency electromagnetic radiation emitted by mobile phones on rat parotid gland histology - an experimental study

Matei LI, Neag MA, Mocan LP, Suflețel RT, Cuțaș A, Onofrei MM, Gherman LM, Armencea G, Mihu C, Ilea A, Mihu CM, Bordea IR, Inchingolo F, Dipalma G, Melincovici CS. The effects of radiofrequency electromagnetic radiation emitted by mobile phones on rat parotid gland histology - an experimental study. Eur Rev Med Pharmacol Sci. 2024 Oct;28(20):4405-4419. doi: 10.26355/eurrev_202410_36864.

Abstract

Objective: The advancement of telecommunication technology and devices promptly transformed mobile phones into indispensable objects in our day-to-day lives, but their biological effects remain unclear. Therefore, this study aimed to investigate the potential histopathological changes induced by mobile phone radiation in the parotid gland and the nearby tissues.

Materials and methods: Thirty female Rattus Norvegicus rats were divided into three groups: group 1 (exposed for 30 days), group 2 (exposed for 60 days), and control group (non-exposed). Each subject was exposed to mobile phone radiation in the form of a phone call for two hours every day for their subsequent exposure time. The exposure was always directed towards the same side of the face throughout the whole exposure period. At the end of the exposure period, a comprehensive examination was conducted, including inspection of the orofacial structures, tissue sections of the parotid glands, overlying skin, oral mucosa, and cervical lymph nodes, as well as obtaining smears from the oral cavity. To highlight the presence of micronuclei within the exfoliated squamous cells of the oral epithelium, Feulgen stain was performed.

Results: The results showed a significant activation of the fibroblasts in the parotid gland septa, in both exposed experimental groups, compared to the control group. We also detected significant cervical lymph node reactive changes, hyperkeratosis of the oral epithelium, and activated fibroblasts in the dermis and oral mucosa lamina propria in both experimental groups. Dermal fibrosis and lamina propria fibrosis were significantly increased in the second experimental group, compared to the control group. Moreover, vascular congestion in the parotid gland, dermal, and lamina propria fibrosis were significantly increased in the second study group compared to the first one.

Conclusions: These findings suggest that exposure to mobile phone radiation may lead to pathological changes in the parotid gland and nearby tissues of experimental rats.

Open access paper: https://www.europeanreview.org/article/36864

--

Frequency-Dependent Antioxidant Responses in HT-1080 Human Fibrosarcoma Cells Exposed to Weak Radio Frequency Fields

Gurhan H, Barnes F. Frequency-Dependent Antioxidant Responses in HT-1080 Human Fibrosarcoma Cells Exposed to Weak Radio Frequency Fields. *Antioxidants*. 2024; 13(10):1237. doi: 10.3390/antiox13101237.

Abstract

This study explores the complex relationship between radio frequency (RF) exposure and cancer cells, focusing on the HT-1080 human fibrosarcoma cell line. We investigated the modulation of reactive oxygen species (ROS) and key antioxidant enzymes, including superoxide dismutase (SOD), peroxidase, and glutathione (GSH), as well as mitochondrial superoxide levels and cell viability. Exposure to RF fields in the 2–5 MHz range at very weak intensities (20 nT) over 4 days resulted in distinct, frequency-specific cellular effects. Significant increases in SOD and GSH levels were observed at 4 and 4.5 MHz, accompanied by reduced mitochondrial superoxide levels and enhanced cell viability, suggesting improved mitochondrial function. In contrast, lower frequencies like 2.5 MHz induced oxidative stress, evidenced by GSH depletion and increased mitochondrial superoxide levels. The findings demonstrate that cancer cells exhibit frequency-specific sensitivity to RF fields even at intensities significantly below current safety standards, highlighting the need to reassess exposure limits. Additionally, our analysis of the radical pair mechanism (RPM) offers deeper insight into RF-induced cellular responses. The modulation of ROS and antioxidant enzyme activities is significant for cancer treatment and has broader implications for age-related diseases, where oxidative stress is a central factor in cellular degeneration. The findings propose that RF fields may serve as a therapeutic tool to selectively modulate oxidative stress and mitochondrial function in cancer cells, with antioxidants playing a key role in mitigating potential adverse effects.

Open access paper: https://www.mdpi.com/2076-3921/13/10/1237

--

The influence of eyelashes on electric field distribution and absorbed power density in the cornea under millimeter-wave exposure

Foroughimehr N, Vilagosh Z, Yavari A, Wood A. The influence of eyelashes on electric field distribution and absorbed power density in the cornea under millimeter-wave exposure. Bioelectromagnetics. 2024 Oct 14. doi: 10.1002/bem.22526.

Abstract

As millimeter wave (MMW) technology, particularly in fifth-generation (5G) devices, gains prominence, there is a crucial need for comprehensive electromagnetic (EM) models of ocular tissues to understand and characterize EM exposure conditions accurately. This study employs numerical modeling to investigate the interaction between MMW and the cornea, aiming to characterize EM field distributions and absorption within an anatomically accurate eye model while considering the influence of eyelashes. Using the finite-difference time-domain (FDTD) method, we conduct simulations of EM radiation interactions from 20.0 to 100.0 GHz with a human eye model. Moreover, we analyze the temperature distribution increase within the eye model using a thermal sensor in XFdtd, employing a scheme based on the finite difference (FD) method. Our findings reveal a nonuniform distribution of the EM field, particularly intensified in corneal regions adjacent to eyelashes and eyelids. Despite similar EM field patterns, the presence or absence of eyelashes has minimal impact on temperature differences. However, the study highlights increased radiation absorption by the eyelid's epidermis at 100.0 GHz, reducing the rise in the cornea's temperature.

https://pubmed.ncbi.nlm.nih.gov/39402869/

--

Exploring the potential link between prostate cancer and magnetic fields

Dart DA, Koushyar S, Uysal-Onganer P. Exploring the potential link between prostate cancer and magnetic fields. Medical Hypotheses. Volume 189, 2024. doi: 10.1016/j.mehy.2024.111384.

Abstract

Prostate cancer is the most common solid cancer in men worldwide. Various lifestyle factors have been established as contributors to cancer risk, and prostate cancer is no exception. Elements such as advanced age, diet, obesity, smoking, alcohol consumption, circadian rhythm disruptions, Afro-Caribbean ethnicity, and sexual activity have all been linked to an increased risk of prostate cancer. Notably, famous male guitarists from bands spanning the last few decades are now entering the age demographic most closely associated with prostate cancer. It is not surprising that they may fall into the 'at-risk' category. Media speculation has hinted at a potentially higher incidence of prostate cancer among guitarists compared to their bandmates, although no tangible correlation has been established. This speculation piqued our curiosity and prompted an exploration into potential reasons why musicians, particularly guitarists, might be at an elevated risk of developing prostate cancer. We hypothesise the electromagnetic fields generated by the electric guitar may in part explain the possible increase in prostate cancer risk among electric guitarists. This study delves into these intriguing possibilities, shedding light on an area of research that remains speculative but warrants further investigation.

Highlights

Prostate cancer is the most common solid cancer in men globally.
Age, diet, ethnicity, sexual activity and circadian rhythm disruptions, are identified as contributors to prostate cancer risk.
Magnetic fields (MF) and electromagnetic fields (EMF) may amplify prostate cancer risks.
Magnetic fields may potentially affect the behaviour of quantum systems within biological molecules.

https://www.sciencedirect.com/science/article/pii/S0306987724001270

--

Electromagnetic Field Exposure in the Public Space of the Slovakian City

Trnka M, Gálik P, Kráľová E, Važan R. Electromagnetic Field Exposure in the Public Space of the Slovakian City. Scientific Letters of the University of Zilina 2023, 25(1):G1-G6. doi: 10.26552/com.C.2023.014

The main objective of our research was to map the exposure to electromagnetic smog in the frequented space of shopping centres in the city of Bratislava and to compare our results to the actual hygienic limits. The measurements of the low- and high-frequency electromagnetic fields were performed at different places in shopping centres. Our results did not exceed the Slovak current limits in any of the

measurements. However, almost all of them markedly exceed new permitted limits according to EUROPAEM. Based on our results, stricter limits in many European countries and increasing evidence on possible harmfulness of long-term exposures to artificial electromagnetic fields, preventive carefulness can be recommended - to support the research in this field, to prepare professional public education and possibly to prepare the stricter Slovak exposure limits.

Open access paper: https://komunikacie.uniza.sk/pdfs/csl/2023/01/08.pdf

--

Development of electromagnetic pollution maps utilizing Gaussian process spatial models

Kiouvrekis Y, Zikas S, Katis I, Tsilikas I, Filippopoulos I. Development of electromagnetic pollution maps utilizing Gaussian process spatial models. Sci Total Environ. 2024 Oct 21;955:176907. doi: 10.1016/j.scitotenv.2024.176907.

Abstract

The rapid proliferation of wireless technologies in everyday environments demands the quick and precise estimation of electromagnetic field distribution. This distribution is commonly depicted through the electric field strength across various geographical areas. The objective of this research is to determine the most effective geospatial model for generating a national-level electric field strength map within the 30 MHz-6 GHz frequency range. To achieve this, we employed five different methodologies for constructing the electric field strength map. Four of these methodologies are based on Gaussian process regression, while the fifth utilizes the classical weighted-average method of the nearest neighbor. Our study focused on a country with a total area of 9251 km^2, using a dataset comprising 3621 measurements. The findings reveal that Gaussian process spatial models, also known as Kriging models, generally outperform other methods when applied to spatial data. However, it was observed that, after excluding some outlier data points, the performance of the classical nearest neighbor models becomes comparable to that of the Gaussian process models. This indicates the potential for both approaches to be effective, depending on the data quality and the presence of outliers.

Excerpts

For the Total (Fig. 4 and Table 2), the mean stands at 4.78 V/m, with a maximum value of 37.99 V/m, a minimum of 0.28 V/m, and a median of 3.42 V/m....

The present research constitutes the first step of many that can be implemented in the future regarding the construction of systems for mapping EMR exposure. Initially, future research should be conducted in cities, exploring characteristics related to urban infrastructure, such as building materials, density, and height. This novel approach regards not only improving the model's performance but also developing models that consider these characteristics between measurement points to better capture the spatial distribution of EMR within urban and semi-urban areas. Additionally, it would be interesting to incorporate temporal patterns to account for the various varieties of EMR over time, considering factors such as daily, seasonal, and yearly variations. In a subsequent stage, the inclusion of data from environmental sources, such as weather conditions, levels of atmospheric pollution, and population

density, could be beneficial to enhance the predictive capability of the models. Furthermore, research into improving the interpretability of the models should be conducted to provide meaningful insights into the factors influencing EMR levels in different environments (urban, semi-urban, and rural). All of the above should be combined with conducting validation studies in collaboration with stakeholders for the effectiveness of the developed models in real-world scenarios.

https://pubmed.ncbi.nlm.nih.gov/39442732/

--

An Analysis of Radio Frequency Radiation Emitted by Smartphones

Manoharan S, Mahalakshmi B, Ananthi K, Sindhu MP. An Analysis of Radio Frequency Radiation Emitted by Smartphones. 2024 8th International Conference on I-SMAC (IoT in Social, Mobile, Analytics and Cloud) (I-SMAC), Kirtipur, Nepal, 2024, pp. 301-307, doi: 10.1109/I-SMAC61858.2024.10714888.

Abstract

The introduction of smartphones has transformed how we interact, access information, and entertains ourselves. Nevertheless, as the usage of smartphones continues to rise, there has been greater concern over the possible health implications of Radio Frequency radiation released by these devices. RF radiation, also known as radiofrequency radiation, is a kind of electromagnetic radiation utilized for wireless communication in devices like smartphones. This study seeks to offer a thorough analysis of RF radiation and its emissions from smartphones, as well as the potential health implications associated with their use and strategies to minimize exposure. This research study also explores the importance of Specific Absorption Rate (SAR) testing in the context of 5G technologies, including testing methodologies, regulatory frameworks, and the implications of 5G frequency bands on SAR measurements.

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=10714888&isnumber=10714582

--

The 5G-FR1 Signals: Beams of the Phased Antennas Array and Time-Recurrence of Emissions with Consequences on Human Exposure

Deaconescu DB, Miclaus S. The 5G-FR1 Signals: Beams of the Phased Antennas Array and Time-Recurrence of Emissions with Consequences on Human Exposure. *Electronics*. 2023; 12(2):297. doi: 10.3390/electronics12020297.

Abstract

The fifth generation (5G) of mobile communication technology poses lots of questions while introducing significant improvements compared with previous generations. The most sensitive question is related to the safety of human exposure. The aim of present work was to analyze, with a few chosen examples, two of the most significant features of 5G emissions: the extreme spatial variability of the exposure and the nonlinear dynamics characteristics of the temporal variability of the exposure. Two models of patch antenna arrays operating at 3.7 GHz with varying beam forming and beam steering capabilities were

considered for an analysis of the specific absorption rate of electromagnetic energy deposition in tissues of a head model. This allowed clear emphasis on the influence of the antenna geometry and feeding peculiarities on the spatial variability of exposure. The second approach implemented the original idea of following the nonlinear recurrence behavior of exposure in time, and underlined the time variability characteristics of emissions with a real-life mobile phone running different 5G applications. Time series of the emitted electric-field strengths were recorded by means a real-time spectrum analyzer and two near-field probes differently positioned in the beam. The presence of laminar emissions, chaotic emissions, determinism and recurrence in the exposures prove the potential for recurrence quantification in predicting time variability features of 5G exposure. Overall, the impact of 5G signals on living bodies, with the highest possible man-made spatial and temporal variability, may have very unpredictable bio-medical consequences.

Open access paper: https://www.mdpi.com/2079-9292/12/2/297

--

Preliminary Study on the Impact of 900MHz Radiation on Human Sperm: An In Vitro Molecular Approach

Keskin I, Karabulut S, Kaplan AA, Alagöz M, Akdeniz M, Tüfekci KK, Davis DL, Kaplan S. Preliminary Study on the Impact of 900MHz Radiation on Human Sperm: An In Vitro Molecular Approach. Reprod Toxicol. 2024 Nov 4:108744. doi: 10.1016/j.reprotox.2024.108744.

Abstract

The use of technologies that produce and emit electromagnetic fields (EMF) is growing exponentially worldwide. The biological effects of EMF-emitting equipment, such as mobile phones and other wireless devices, have been studied in the last decade using in vitro and in vivo methods. Infertility is a growing health problem, and nearly half of cases are because of male-factor. This study investigated the direct in vitro effects of 900MHz radiation exposure on sperm parameters, genetic status, apoptotic markers, and the PI3K/AKT signaling pathway in healthy normozoospermic men. Semen samples were divided into four groups, two control (30min and 1h) and two EMF exposure (30min and 1h). Sperm parameters (motility, progressive motility, acrosomal index, morphology), genetic status (DNA fragmentation and chromatin integrity), apoptotic markers (cytokine-c and caspase-3 expression) and the PI3K/AKT signaling pathway (phosphoinoitide 3-kinase-PI3K- and phosphorylated AKT- p-AKT-) were analysed. Sperm motility were significantly reduced in 30min EMF exposure while a significant increase in the expression of p-AKT were observed in 1h EMF exposure group. An increased vacuolisation, acrosomal defect, extension of subacrosomal space, uncondensed chromatin structure, apoptotic signs and disrupted axoneme were observed in both EMF groups which were not observed in the control group. Other sperm parameters (morphology and acrosomal index), genetic status, apoptotic markers and the PI3K expression rates had no significant change.

https://pubmed.ncbi.nlm.nih.gov/39505052/

--

Quantitative Assessment of Thermal Effects on the Auricle Region Caused by Mobile Phones Operating in Different Modes

Rok T, Kacprzyk A, Rokita E, Kantor D, and Taton G. Quantitative Assessment of Thermal Effects on the Auricle Region Caused by Mobile Phones Operating in Different Modes. *AIMS Biophys* 11, no. 4 (2024): 427–44. https://doi.org/10.3934/biophy.2024023.

Abstract

To analyze thermal effects caused by mobile phones on the human auricle region, we performed an experiment with controlled exposure to mobile phones operating in different modes for a group of 40 men. Temperature changes were measured with the use of infrared thermography. Thermograms were taken before and after a standardized 15-minute phone call when the mobile phone was placed lightly against the skin surface in the auricle region. The measurements were performed in three modes: OFF, ON, and FLIGHT. Statistically significant differences ($p = 0.03$) were observed between the experimental temperature increase of the auricle region in OFF mode (average temperature rise = 1.1 °C ± 0.2 °C) and in ON mode (average temperature rise = 1.9 °C ± 0.3 °C), while between FLIGHT (average temperature rise = 1.4 °C ± 0.2 °C) and ON modes, no statistical differences were observed ($p = 0.20$). Based on thermographic measurements and the model of heat transfer between the ear and the phone, it was shown that the human ear is the largest heat source in the system and that the increase in skin temperature is mainly caused by the handheld mobile phone restricting heat dissipation from the skin surface.

Excerpt for the assessment of near-field exposure, Received Signal Strength Indicator (RSSI) and Specific Absorption Rate (SAR) were used. During exposure sessions RSSI values were continuously measured and retrospectively controlled. All the participants received exposure at an arbitrary mean value of RSSI between −90 and −80 dBm. The SAR quantifies the absorption of RF-EMF in tissues, and for the human body, it depends on the dielectric properties of tissues due to the complexity of the human head anatomy its estimation is a subject of great challenge [21],[22] and for the phone used (Huawei P20 Lite), it was reported as 0.75 W/kg in the manual.

For the far-field exposure a personal exposure meter (ExpoM-RF, Zurich, Switzerland) performing a spectral analysis of RF-EMF within 16 different frequency bands from 87.5 MHz up to 5.875 GHz was used. Based on electric field strength, the power density was calculated as 45 µW. The uplink and downlink radiation remained on a comparable level throughout a simulated phone call.

In OFF mode, the smartphone was switched off, while in FLIGHT mode, there was no connection with the GSM network, so music was constantly playing. In ON mode, the mobile phone was fully connected to the GSM network and the smartphone was operating at a frequency of 1800 MHz.

Open access paper: http://www.aimspress.com/article/doi/10.3934/biophy.2024023

--

Rat brain and testicular tissue effects of radiofrequency radiation exposure: Histopathological, DNA damage of brain and qRT-PCR analysis

Yavas MC, Kilitci A, Çelik E, Yegin K, Sirav B, Varol S. Rat brain and testicular tissue effects of radiofrequency radiation exposure: Histopathological, DNA damage of brain and qRT-PCR analysis. INTERNATIONAL JOURNAL OF RADIATION RESEARCH. JUL 2024. 22(3):529-536. doi: 10.61186/ijrr.22.3.529

Abstract

We evaluate the effects of radiofrequency electromagnetic field (RF-EMF) on rat brain and testicular tissue using histopathology, comet assay, and real-time quantitative PCR techniques.

Materials and Methods: Two equal groups of fourteen rats one for sham-control and the other for exposure (n = seven) were created. For a duration of 14 days, the exposure group (2100 MHz, testicular tissue SAR values of 163 mW/kg for 10 g, brain tissue SAR values of 292 mW/ kg on average) was subjected to five hours of exposure per day. Evaluations were conducted on tissue gene expression levels, histopathology, and DNA damage to brain tissue.

Results: The histological examination of brain tissue from the exposed group revealed vascular alterations and significant edema ($p < 0.05$). It was determined that RF radiation-induced much more cellular damage in the exposed group (18.26% tail DNA) than in the control group (4.06% tail DNA). Signs of deterioration in spermatogenic cells in the testicular tissue of the exposed group also changed significantly ($p < 0.05$). The Bax and bcl-2 genes showed a significant difference ($p < 0.05$) in the mRNA level data, whereas the p53 genes showed no significant change ($p > 0.05$).

Conclusion: These findings suggest that it may cause some histopathological and cellular damage in brain and testis tissue.

Open access paper: https://ijrr.com/article-1-5507-en.html

--

Does Radiofrequency Radiation From Mobile Phones Affect the Formation of Parotid Gland Malignancy? An Experimental Study

Ozergin Coskun Z, Tumkaya L, Yilmaz A, Dursun E, Mercantepe T, Kalkan Y, Ersoz S. Does Radiofrequency Radiation From Mobile Phones Affect the Formation of Parotid Gland Malignancy? An Experimental Study. Ear Nose Throat J. 2024 Sep 27:1455613241287295. doi: 10.1177/01455613241287295.

Abstract

Objectives: The use of mobile phone is increasing around the world. Although it is beneficial in terms of communication, the electromagnetic radiations emitted by mobile phones may cause undesirable biological effects on the human body. In practical use, the tissue with which mobile phones come into most and are closest is the parotid gland. This study investigated the effects of the 1800 MHz electromagnetic field created by a generator on the parotid gland in rats.

Methods: A total of 21 Sprague-Dawley Albino rats were included in the study. The rats were randomly divided into three equal groups. To simulate a mobile phone in conversation mode, the first study group was exposed to an 1800-MHz electromagnetic field for 6 hours a day for 30 days, and the second study group was exposed to an 1800-MHz electromagnetic field for 12 hours a day for 30 days. After 30 days, rats were sacrificed, and histopathological and immunohistochemical methods were used to evaluate the effects on the parotid gland. The total antioxidant level and the total oxidant level were measured biochemically in homogenized parotid tissue.

Results: Histopathological results showed an increase in degeneration in rats exposed to electromagnetic fields for 6 and 12 hours a day, and immunohistochemical analysis showed an increase in the apoptotic index in both study groups ($P = .001$, $P < .001$). Intranuclear inclusions was observed during histopathological examination performed by electron microscopy.

Conclusions: This study observed that the 1800 MHz electromagnetic field caused undesirable adverse histopathological and biochemical effects on the parotid gland of rats. Histopathological and biochemical findings were detected with increasing contact and exposure time. This study will lead to other studies on this topic and contribute to the literature by completing other studies.

Excerpts

This study used a generator (Anritsu MG3670 B type, Japan) that produces microwave radiation at 1800 MHz radiofrequency to create exposure in mobile communion. The generator peak power was fixed at 2 W during exposure. In the digital signal generator used in this study, the carrier frequency was 1800 MHz, the modulation frequency was 217 Hz, there was a pulse of 577 µseconds, and the maximum peak power was 2 W. The average specific absorption rate (SAR) of the whole body was 0.117 W/kg. The study was carried out on a total of 21 rats divided in 3 groups (n = 7).

Groups 1, 2, and 3 were exposed to microwave radiation for 0 (control group), 6, and 12 hours, respectively. A generator with an external antenna was placed in the lower middle part of the cages.[5] The rats in the study group were exposed to microwave radiation for the specified hours in the mobile phone conversation mode....

The parotid tissue has been reported to absorb 40% of the electromagnetic energy emitted by mobile phones on phones held in place. This absorbed energy can cause thermal and nonthermal effects in the parotid tissue.[14] ....

Conclusion

In light of our knowledge of the literature, this is the first study to investigate the effects of an 1800-MHz electromagnetic field on the parotid gland of rats with biochemical tests, immunohistochemically in light microscopy and histopathologically in electron microscopy. This study observed that an 1800-MHz electromagnetic field causes undesirable adverse histopathological and biochemical effects on the parotid gland of rats. Histopathological and biochemical findings were detected with increasing contact and exposure time. This study will lead other studies on this topic and contribute to the literature in this area.

Open access paper: https://journals.sagepub.com/doi/10.1177/01455613241287295

--

Blueberry anthocyanins regulate SIRT1/FoxO1 pathway to inhibit oxidative stress and reduce testicular tissue damage induced by microwave radiation in rats

Pang Y, Men J, Li Y, Zhang J, Zhao L, et al. Blueberry anthocyanins regulate SIRT1/FoxO1 pathway to inhibit oxidative stress and reduce testicular tissue damage induced by microwave radiation in rats. Journal of Functional Foods. Vol. 122, 2024. doi: 10.1016/j.jff.2024.106523.

Abstract

Researches have shown that microwave radiation could cause oxidative stress injury in male reproductive system, and blueberry anthocyanins had excellent oxidation resistance. Our study aimed to investigate the protective effect of blueberry anthocyanins (100, 200 and 400 mg/kg/d) on testicular tissue damage in Wistar rats exposed to 2.856 GHz microwave and the optimal dose. We found that blueberry anthocyanins could ameliorate the decrease of sperm motility and sex hormone levels and testicular tissue structure damage caused by microwave radiation, increase SIRT1 expression and decrease FoxO1 expression, increase GSH/GSSG, SOD and inhibit MDA. The LDH, SDH and ATP synthase were increased, and Caspase-3 expression was decreased, and the high-dose of blueberry anthocyanins (400 mg/kg/d) had the best protective effect. These results suggested that blueberry anthocyanins could inhibit oxidative stress injury induced by 2.856 GHz microwave radiation in rat testicular tissue by regulating SIRT1/FoxO1 pathway, enhance energy metabolism and reduce cell apoptosis.

Open access paper: https://www.sciencedirect.com/science/article/pii/S1756464624005255

--

Effects of 4.9 GHz Radiofrequency Field Exposure on Brain Metabolomic and Proteomic Characterization in Mice

Wang X, Zhou G, Lin J, Zhang Z, Qin T, Guo L, Wang H, Huang Z, Ding G. Effects of 4.9 GHz Radiofrequency Field Exposure on Brain Metabolomic and Proteomic Characterization in Mice. Biology (Basel). 2024 Oct 10;13(10):806. doi: 10.3390/biology13100806.

Abstract

Electromagnetic exposure has become increasingly widespread, and its biological effects have received extensive attention. The purpose of this study was to explore changes in the metabolism profile of the brain and serum and to identify differentially expressed proteins in the brain after exposure to the 4.9 GHz radiofrequency (RF) field. C57BL/6 mice were randomly divided into a Sham group and an RF group, which were sham-exposed and continuously exposed to a 4.9 RF field for 35 d, 1 h/d, at an **average power density (PD) of 50 W/m^2**. After exposure, untargeted metabolomics and Tandem Mass Tags (TMT) quantitative proteomics were performed. We found 104 and 153 up- and down-regulated differentially expressed metabolites (DEMs) in the RF_Brain group and RF_Serum group, and the DEMs were significantly enriched in glycerophospholipid metabolism. Moreover, 10 up-regulated and 51 down-regulated differentially expressed proteins (DEPs) were discovered in the RF group. Functional correlation analysis showed that most DEMs and DEPs showed a significant correlation. These results suggested that 4.9 GHz exposure induced disturbance of metabolism in the brain and serum, and caused deregulation of proteins in the brain.

Simple Summary

The brain, as the central nervous system that controls the body's sensory, behavior, and mental symptoms, is sensitive to RF exposure, and lots of studies have explored the potential health hazards of RF-EMR with different frequencies to the brain. Our previous study found that 4.9 GHz radiofrequency radiation induced depression-like behavior in mice, but the mechanism of the behavioral changes was unclear. Studies have shown that changes in peripheral energy metabolism might affect brain lipid levels, and thereby cortical excitability, and a deregulated hippocampus proteome might influence the healthy functioning of the brain. Here, we provide evidence that 4.9 GHz RF exposure altered metabolite expression patterns in brain tissue and serum, especially glycerophospholipid metabolism. In addition, 4.9 GHz RF exposure induced an imbalance in the protein profile of brain tissue and may alter gap junction communication. Our results initially revealed the biological effects of 5G communication frequency exposure and provided a possible mechanism for electromagnetic radiation-induced behavioral changes from the perspective of metabolome and proteome.

Open access paper: https://www.mdpi.com/2079-7737/13/10/806

--

Cellular signaling pathways in the nervous system activated by various mechanical and electromagnetic stimuli

Kazuhito M, Ferguson AR, Aboubacar W, Youngjae R. Cellular signaling pathways in the nervous system activated by various mechanical and electromagnetic stimuli. Frontiers in Molecular Neuroscience. Vol. 17. 2024. doi: 10.3389/fnmol.2024.1427070.

Abstract

Mechanical stimuli, such as stretch, shear stress, or compression, activate a range of biomolecular responses through cellular mechanotransduction. In the nervous system, studies on mechanical stress have highlighted key pathophysiological mechanisms underlying traumatic injury and neurodegenerative diseases. However, the biomolecular pathways triggered by mechanical stimuli in the nervous system has not been fully explored, especially compared to other body systems. This gap in knowledge may be due to the wide variety of methods and definitions used in research. Additionally, as mechanical stimulation techniques such as ultrasound and electromagnetic stimulation are increasingly utilized in psychological and neurorehabilitation treatments, it is vital to understand the underlying biological mechanisms in order to develop accurate pathophysiological models and enhance therapeutic interventions. This review aims to summarize the cellular signaling pathways activated by various mechanical and electromagnetic stimuli with a particular focus on the mammalian nervous system. Furthermore, we briefly discuss potential cellular mechanosensors involved in these processes

Open access paper: https://www.frontiersin.org/journals/molecular-neuroscience/articles/10.3389/fnmol.2024.1427070

--

--

Neurobiological effects and mechanisms of magnetic fields: a review from 2000 to 2023

Wang X, Ye Y, Zuo H, Li Y. Neurobiological effects and mechanisms of magnetic fields: a review from 2000 to 2023. BMC Public Health. 2024 Nov 8;24(1):3094. doi: 10.1186/s12889-024-18987-9.

Abstract

Magnetic fields are widely used in medical diagnostics because of their superior non-invasive properties. In addition, with the widespread use of magnetic fields in transportation and other areas, their potential hazards to human health and the assessment of their safety have attracted considerable attention. The effects of magnetic fields on living organisms have a long history. The biological effects of magnetic field exposure in mice and rats depend on the magnetic field strength, exposure time, and direction; depending on these and potentially other factors, magnetic fields can cause a series of neurobiological effects. We reviewed global research on the neurobiological effects of magnetic fields from recent years to provide an overview and insights into the underlying mechanisms. This review focuses on the biological effects of static and dynamic magnetic fields of different frequencies and intensities on animals and nerve cells and their mechanisms of action.

Conclusion

In summary, this article reviews the progress in research on the neural effects of SMF and DMF at the level of animal models and nerve cells in recent years. The number of studies on the effects of MFs on learning memory, emotional behavior, nerve cells, and neurotransmitters is gradually increasing, but owing to the diversity of MF parameters, experimental subjects, and conditions, the conclusions are inconsistent. Certain conditions of MF exposure can lead to changes in emotional behavior and learning memory and cause or relieve anxiety-like and depressive behaviors, with or without significant effects. The biological effects of MFs on neurons and glial cells include alterations in cell proliferation, cell cycle distribution, and apoptosis. However, some problems remain unclear. Due to the unspecific nature of MFs, their neurobiological effects are difficult to target experimentally. Urgent problems to be solved by future research include how to establish proper experimental animal and neural cell models, and how to select the appropriate MF exposure intensity and time.

Open access paper: https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-024-18987-9

--

Cellular and Molecular Effects of Magnetic Fields

Tota M, Jonderko L, Witek J, Novickij V, Kulbacka J. Cellular and Molecular Effects of Magnetic Fields. Int J Mol Sci. 2024 Aug 17;25(16):8973. doi: 10.3390/ijms25168973.

Abstract

Recently, magnetic fields (MFs) have received major attention due to their potential therapeutic applications and biological effects. This review provides a comprehensive analysis of the cellular and molecular impacts of MFs, with a focus on both in vitro and in vivo studies. We investigate the mechanisms by which MFs influence cell behavior, including modifications in gene expression, protein synthesis, and cellular signaling pathways. The interaction of MFs with cellular components such as ion channels, membranes, and the cytoskeleton is analyzed, along with their effects on cellular processes like proliferation, differentiation, and apoptosis. Molecular insights are offered into how MFs modulate oxidative stress and inflammatory responses, which are pivotal in various pathological conditions.

Furthermore, we explore the therapeutic potential of MFs in regenerative medicine, cancer treatment, and neurodegenerative diseases. By synthesizing current findings, this article aims to elucidate the complex bioeffects of MFs, thereby facilitating their optimized application in medical and biotechnological fields.

Open access paper: https://www.mdpi.com/1422-0067/25/16/8973

--

Morphological, biochemical and genotoxic effects of non-ionizing radiation at 1800 MHz and 2400 MHz frequencies in Allium cepa L

Sharma A, Bahel S, Katnoria JK. Morphological, biochemical and genotoxic effects of non-ionizing radiation at 1800 MHz and 2400 MHz frequencies in Allium cepa L. Environ Sci Pollut Res Int. 2024 Oct 30. doi: 10.1007/s11356-024-35414-z.

Abstract

The frequent use of electronic devices in daily lives, predominantly reliant on non-ionizing radiation, has increased the prevalence of electromagnetic radiation (EMR) in natural environment. In light of this, effects of EMR at frequencies of 1800 MHz and 2400 MHz characterized by a power of 10.0 dBm (0.01 W), across varying exposure durations of 1 h/day, 2 h/day, 4 h/day, 6 h/day, and 8 h/day for 7 days, in Allium cepa L. were studied. The effects of the treatment on the morphological features (root length, fresh weight, and dry weight of roots) and biochemical characteristics (protein content and antioxidative enzymes, namely, ascorbate peroxidase (APX), glutathione reductase (GR), superoxide dismutase (SOD), glutathione-S-transferase (GST), guaiacol peroxidase (POD), catalase (CAT), and dehydroascorbate (DHAR)) were studied in roots and bulbs of Allium cepa L. Further genotoxicity for different exposure periods at both frequencies was also conducted. Prolonged exposure to electromagnetic radiation (EMR) at both frequencies was found to reduce root length, fresh weight, and dry weight of plant. Furthermore, significant effects were observed on protein content, indicating a reduction with prolonged exposure duration. Investigation into the activities of antioxidative enzymes such as APX, GR, GST, DHAR, CAT, SOD, and POD at a frequency of 1800 MHz and 2400 MHz in roots and bulbs demonstrated a significant enhancement in enzyme activity during 6 h/day and 8 h/day exposure periods. Additional investigation during genotoxicity studies demonstrated the induction of chromosomal aberrations in the root tip cells of the Allium cepa L. plant test system. The current study revealed the initiation of oxidative stress and genotoxicity resulting from long-term exposure to electromagnetic radiation in the studied plant test systems.

https://pubmed.ncbi.nlm.nih.gov/39472374/

--

Representations of 5G in the Chinese and British press: a corpus-assisted critical discourse analysis

Pei, J., Cheng, L. Representations of 5G in the Chinese and British press: a corpus-assisted critical discourse analysis. *Humanit Soc Sci Commun* **11**, 400 (2024). https://doi.org/10.1057/s41599-024-02896-8

Abstract

This study employs a corpus-assisted critical discourse analysis to demystify the dominant patterns of representations around 5G in the Chinese and British press. Keyword analyses identify four thematic categories around the representation of 5G: the nature of 5G, social actors in 5G discourse, actions around 5G and timing in 5G discourse. Findings suggest that the Chinese press tends to use positive evaluative expressions to depict 5G as a trustworthy and beneficial issue and stress China's strong support for 5G development. In contrast, the British press prefers to frame 5G as a product of geopolitical rivalry and an issue with scientific uncertainties and controversies by repeatedly employing negative language patterns related to 5G risks and conspiracy theories. Besides, a subtle but varying "self versus other" schema is constructed by the two presses. The Chinese press is inclined to use positive predication strategies to construct a positive self-representation, whereas the British press tends to adopt negative predication strategies to portray China as an outgroup and meanwhile use scapegoating strategies to profile Britain as a positive self by using recurrent patterns denoting the pressure exerted on Britain by the United States. Such differences could result from their journalistic ideologies and values and the contrasting socio-political contexts where the two presses are situated.

Open access paper: https://www.nature.com/articles/s41599-024-02896-8

--

Magnetic fields from indoor transformer stations and risk of cancer in adults: a cohort study

Juutilainen J, Khan MW, Naarala J, Roivainen P. Magnetic fields from indoor transformer stations and risk of cancer in adults: a cohort study. Occup Environ Med. 2024 Nov 18:oemed-2024-109466. doi: 10.1136/oemed-2024-109466.

Abstract

Objectives: Studies assessing the association of adult cancers with extremely low frequency (ELF) magnetic fields (MF) have provided inconclusive results, probably affected by limitations such as low exposure levels, confounding and various forms of bias. This study investigated the association between residential ELF MF exposure and adult cancer using a design that avoids the main limitations of previous studies.

Methods: Persons who have lived in buildings with indoor transformer stations during the period 1971-2016 formed the study cohort. Their MF exposure was assessed based on the location of their apartment in relation to the transformer room. Information on their cancer diagnoses was obtained from the Finnish Cancer Registry. SIR with 95% CI was calculated to investigate the association of MF exposure with overall cancer and specific cancers.

Results: The SIR for all primary sites was 1.01 (95% CI 0.93 to 1.09). An increased risk of digestive organ cancers was observed among the exposed persons, with a SIR of 1.23 (95% CI 1.03 to 1.46). The highest SIR was observed for gallbladder cancer (3.92, 95% CI 1.44 to 8.69). Increased risk of testicular cancer was observed among men exposed to MF during childhood, but this is likely to be due to confounding associated with living on the lowest floors. No other significant associations were observed for other primary cancer sites studied.

Conclusions: Overall cancer risk was not affected by residential MF exposure. The increased risk of digestive organ cancers among MF-exposed persons is a novel finding requiring confirmation in further studies.

WHAT IS ALREADY KNOWN ON THIS TOPIC

- Extremely low frequency magnetic fields (ELF MFs) have been classified as possibly carcinogenic. The evidence is strongest for increased risk of childhood leukaemia and more inconsistent for risks of adult cancers.

WHAT THIS STUDY ADDS

- This study was carried out using a unique database of residential buildings with indoor transformer stations. Many limitations of previous studies could be avoided. The study suggested an association between residential MF exposure and digestive organ cancers. The overall cancer risk was not affected by MF exposure.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- As there is little previous evidence of an association between digestive organ cancers and MF exposure, further research on this topic is required. Compared with previous studies, this study included higher exposure levels and thus provides a useful addition to the data available for health risk assessment and risk communication.

https://pubmed.ncbi.nlm.nih.gov/39557565/

--

Exploring Non-Thermal Mechanisms of Biological Reactions to Extremely Low-Frequency Magnetic Field Exposure

Radil R, Carnecka L, Judakova Z, Pobocikova I, Bajtos M, Janousek L. Exploring Non-Thermal Mechanisms of Biological Reactions to Extremely Low-Frequency Magnetic Field Exposure. *Applied Sciences*. 2024; 14(20):9409. https://doi.org/10.3390/app14209409

Abstract

The increasing evidence regarding biological effects of exposure to an extremely low frequency magnetic field is of utmost interest not only to the scientific community, but also to legislative bodies and the public. However, the research in this field is full of controversial and inconsistent results, originated from a lack of widely acceptable physical mechanisms that could sufficiently describe the principle of such a field's action. This experimental study addresses and points to possible sources of ambiguities via investigation of the ion parametric resonance mechanism at 50 Hz frequency. The chosen methodology incorporates exposure of the Saccharomyces cerevisiae yeast strain based on an established exposure protocol with special attention to the measurement of an applied time-varying magnetic field corresponding to the ion parametric resonance requirements. Subsequently, the differences in cell growth as a reaction to changes in magnetic flux density are evaluated and statistically

analyzed. It is found that fluctuations in the magnetic field within the exposure setup need to be addressed properly, since this could have an impact on replication of the experiments and reliability of the results. Furthermore, comparison of two independently performed sets of 10 experiments showed statistically significant effects even in conditions that did not fulfill the requirements of the resonance theory, putting the validity and practical application of the ion parametric resonance model into question.

Open access paper: https://www.mdpi.com/2076-3417/14/20/9409

--

Migratory birds can extract positional information from magnetic inclination and magnetic declination alone

Packmor F, Kishkinev D, Zechmeister T, Mouritsen H, Holland RA. Migratory birds can extract positional information from magnetic inclination and magnetic declination alone. Proc Biol Sci. 2024 Nov;291(2034):rspb20241363. doi: 10.1098/rspb.2024.1363.

Abstract

Migratory birds are able to navigate over great distances with remarkable accuracy. The mechanism they use to achieve this feat is thought to involve two distinct steps: locating their position (the 'map') and heading towards the direction determined (the 'compass'). For decades, this map-and-compass concept has shaped our perception of navigation in animals, although the nature of the map remains debated. However, some recent studies suggest the involvement of the Earth's magnetic field in the map step. Here, we tested whether migratory songbirds, Eurasian reed warblers (*Acrocephalus scirpaceus*), can determine their position based on two magnetic field components that are also associated with direction finding, i.e. magnetic inclination and magnetic declination. During a virtual magnetic displacement experiment, the birds were exposed to altered magnetic inclination and magnetic declination values that would indicate a displacement from their natural migratory corridor, but the total intensity of the field remained unchanged, creating a spatial mismatch between these components. The response was a change in the birds' migratory direction consistent with a compensatory re-orientation. This suggests that birds can extract positional as well as directional information from these cues, even when they are in conflict with another component of the magnetic field. It remains to be seen whether birds use the total intensity of Earth's magnetic field for navigation.

Open access paper: https://royalsocietypublishing.org/doi/10.1098/rspb.2024.1363

--

Indoor & outdoor artificial light-at-night (ALAN) & cancer risk: A systematic review & meta-analysis of multiple cancer sites with a critical appraisal of exposure assessment

Palomar-Cros A, Deprato A, Papantoniou K, Straif K, Lacy P, Maidstone R, Adan A, Haldar P, Moitra S, Navarro JF, Durrington H, Moitra S, Kogevinas M, Harding BN. Indoor and outdoor artificial light-at-night (ALAN) and cancer risk: A systematic review and meta-analysis of multiple cancer sites and with a critical appraisal of exposure assessment. Science of The Total Environment, Vol 955, 2024. doi: 10.1016/j.scitotenv.2024.177059.

Abstract

Exposure to artificial light-at-night (ALAN) has been linked to cancer risk. Few meta-analyses on this topic have reviewed only breast cancer. This study aimed to systematically review and meta-analyze existing studies on ALAN exposure and cancer incidence, thoroughly evaluating exposure assessment quality. We considered observational studies (cohort, case-control, cross-sectional) on ALAN exposure (indoor and outdoor) and cancer incidence, measured by relative risk, hazard ratio, and odds ratio. We searched six databases, two registries, and Google Scholar from inception until April 17, 2024. Quality of studies was assessed using the Joanna Briggs Institute (JBI) critical appraisal tools. Random-effects meta-analysis was used to estimate relative risks (RR) and 95 % confidence intervals (CI) for ALAN exposures. We identified 9835 studies and included 28 for qualitative synthesis with 2,508,807 individuals (15 cohort, 13 case-control). Out of the included studies, 20 studies on breast cancer (731,493 individuals) and 2 studies on prostate cancer (53,254 individuals) were used for quantitative synthesis. Higher levels of outdoor ALAN were associated with breast cancer risk (meta-estimate = 1.12, 95 % CI 1.03–1.23 (I2 = 69 %)). We observed a non-significant positive association between indoor ALAN levels and breast cancer risk (meta-estimate = 1.07, 0.95–1.21, I2 = 60 %), and no differences by menopausal status. The meta-analysis for prostate cancer suggested a non-statistically significant increased risk for higher levels of outdoor ALAN (meta-estimate = 1.43, 0.75–2.72, I2 = 90 %). In the qualitative synthesis, we observed positive associations with non-Hodgkin lymphoma and colorectal, pancreatic and thyroid cancer. We found an association between outdoor ALAN and breast cancer risk. However, most studies relied on satellite-images with a very low resolution (1 to 5 km, from the Defense Meteorological Program [DMSP]) and without information on color of light. Future studies with better exposure assessment should focus on investigating other cancer sites.

https://www.sciencedirect.com/science/article/pii/S0048969724072164

--

Effect of exposure to RF on cancer risk: A systematic review of human observational studies – Part I: Most researched outcomes (WHO SR 1A)

My note: See Biased WHO-commissioned review claims no cancer link

Karipidis K, Baaken D, Loney T, Blettner M, Brzozek C, Elwood M, Narh C, Orsini N, Röösli M, Paulo MS, Lagorio S. The effect of exposure to radiofrequency fields on cancer risk in the general and working population: A systematic review of human observational studies – Part I: Most researched outcomes. Environment International (2024), doi: 10.1016/j.envint.2024.108983. Open access paper: https://doi.org/10.1016/j.envint.2024.108983

Abstract

Background The objective of this review was to assess the quality and strength of the evidence provided by human observational studies for a causal association between exposure to radiofrequency electromagnetic fields (RF-EMF) and risk of the most investigated neoplastic diseases.

Methods Eligibility criteria: We included cohort and case-control studies of neoplasia risks in relation to

three types of exposure to RF-EMF: near-field, head-localized, exposure from wireless phone use (SR-A); far-field, whole body, environmental exposure from fixed-site transmitters (SR-B); near/far-field occupational exposures from use of hand-held transceivers or RF-emitting equipment in the workplace (SR-C). While no restrictions on tumour type were applied, in the current paper we focus on incidence-based studies of selected "critical" neoplasms of the central nervous system (brain, meninges, pituitary gland, acoustic nerve) and salivary gland tumours (SR-A); brain tumours and leukaemias (SR-B, SR-C). We focussed on investigations of specific neoplasms in relation to specific exposure sources (i.e. E-O pairs), noting that a single article may address multiple E-O pairs.

Information sources: Eligible studies were identified by literature searches through Medline, Embase, and EMF-Portal.

Risk-of-bias (RoB) assessment: We used a tailored version of the Office of Health Assessment and Translation (OHAT) RoB tool to evaluate each study's internal validity. At the summary RoB step, studies were classified into three tiers according to their overall potential for bias (low, moderate and high).

Data synthesis: We synthesized the study results using random effects restricted maximum likelihood (REML) models (overall and subgroup meta-analyses of dichotomous and categorical exposure variables), and weighted mixed effects models (dose–response meta-analyses of lifetime exposure intensity).

Evidence assessment: Confidence in evidence was assessed using the Grading of Recommendations, Assessment, Development and Evaluations (GRADE) approach.

Results We included 63 aetiological articles, published between 1994 and 2022, with participants from 22 countries, reporting on 119 different E-O pairs. RF-EMF exposure from mobile phones (ever or regular use vs no or non-regular use) was not associated with an increased risk of glioma [meta-estimate of the relative risk (mRR) = 1.01, 95 % CI = 0.89–1.13), meningioma (mRR = 0.92, 95 % CI = 0.82–1.02), acoustic neuroma (mRR = 1.03, 95 % CI = 0.85–1.24), pituitary tumours (mRR = 0.81, 95 % CI = 0.61–1.06), salivary gland tumours (mRR = 0.91, 95 % CI = 0.78–1.06), or paediatric (children, adolescents and young adults) brain tumours (mRR = 1.06, 95 % CI = 0.74–1.51), with variable degree of across-study heterogeneity (I2 = 0 %-62 %). There was no observable increase in mRRs for the most investigated neoplasms (glioma, meningioma, and acoustic neuroma) with increasing time since start (TSS) use of mobile phones, cumulative call time (CCT), or cumulative number of calls (CNC). Cordless phone use was not significantly associated with risks of glioma [mRR = 1.04, 95 % CI = 0.74–1.46; I2 = 74 %) meningioma, (mRR = 0.91, 95 % CI = 0.70–1.18; I2 = 59 %), or acoustic neuroma (mRR = 1.16; 95 % CI = 0.83–1.61; I2 = 63 %). Exposure from fixed-site transmitters (broadcasting antennas or base stations) was not associated with childhood leukaemia or paediatric brain tumour risks, independently of the level of the modelled RF exposure. Glioma risk was not significantly increased following occupational RF exposure (ever vs never), and no differences were detected between increasing categories of modelled cumulative exposure levels.

Discussion In the sensitivity analyses of glioma, meningioma, and acoustic neuroma risks in relation to mobile phone use (ever use, TSS, CCT, and CNC) the presented results were robust and not affected by

changes in study aggregation.

In a leave-one-out meta-analyses of glioma risk in relation to mobile phone use we identified one influential study. In subsequent meta-analyses performed after excluding this study, we observed a substantial reduction in the mRR and the heterogeneity between studies, for both the contrast Ever vs Never (regular) use (mRR = 0.96, 95 % CI = 0.87–1.07, I2 = 47 %), and in the analysis by increasing categories of TSS ("<5 years": mRR = 0.97, 95 % CI = 0.83–1.14, I2 = 41 %; "5-9 years ": mRR = 0.96, 95 % CI = 0.83–1.11, I2 = 34 %; "10+ years": mRR = 0.97, 95 % CI = 0.87–1.08, I2 = 10 %).

There was limited variation across studies in RoB for the priority domains (selection/attrition, exposure and outcome information), with the number of studies evenly classified as at low and moderate risk of bias (49 % tier-1 and 51 % tier-2), and no studies classified as at high risk of bias (tier-3). The impact of the biases on the study results (amount and direction) proved difficult to predict, and the RoB tool was inherently unable to account for the effect of competing biases. However, the sensitivity meta-analyses stratified on bias-tier, showed that the heterogeneity observed in our main meta-analyses across studies of glioma and acoustic neuroma in the upper TSS stratum (I2 = 77 % and 76 %), was explained by the summary RoB-tier. In the tier-1 study subgroup, the mRRs (95 % CI; I2) in long-term (10+ years) users were 0.95 (0.85–1.05; 5.5 %) for glioma, and 1.00 (0.78–1.29; 35 %) for acoustic neuroma.

The time-trend simulation studies, evaluated as complementary evidence in line with a triangulation approach for external validity, were consistent in showing that the increased risks observed in some case-control studies were incompatible with the actual incidence rates of glioma/brain cancer observed in several countries and over long periods. Three of these simulation studies consistently reported that RR estimates > 1.5 with a 10+ years induction period were definitely implausible, and could be used to set a "credibility benchmark". In the sensitivity meta-analyses of glioma risk in the upper category of TSS excluding five studies reporting implausible effect sizes, we observed strong reductions in both the mRR [mRR of 0.95 (95 % CI = 0.86–1.05)], and the degree of heterogeneity across studies (I2 = 3.6 %).

Conclusions Consistently with the published protocol, our final conclusions were formulated separately for each exposure-outcome combination, and primarily based on the line of evidence with the highest confidence, taking into account the ranking of RF sources by exposure level as inferred from dosimetric studies, and the external coherence with findings from time-trend simulation studies (limited to glioma in relation to mobile phone use).

For near field RF-EMF exposure to the head from mobile phone use, there was moderate certainty evidence that it likely does not increase the risk of glioma, meningioma, acoustic neuroma, pituitary tumours, and salivary gland tumours in adults, or of paediatric brain tumours.

For near field RF-EMF exposure to the head from cordless phone use, there was low certainty evidence that it may not increase the risk of glioma, meningioma or acoustic neuroma.

For whole-body far-field RF-EMF exposure from fixed-site transmitters (broadcasting antennas or base stations), there was moderate certainty evidence that it likely does not increase childhood leukaemia risk and low certainty evidence that it may not increase the risk of paediatric brain tumours. There were

no studies eligible for inclusion investigating RF-EMF exposure from fixed-site transmitters and critical tumours in adults.

For occupational RF-EMF exposure, there was low certainty evidence that it may not increase the risk of brain cancer/glioma, but there were no included studies of leukemias (the second critical outcome in SR-C).

The evidence rating regarding paediatric brain tumours in relation to environmental RF exposure from fixed-site transmitters should be interpreted with caution, due to the small number of studies. Similar interpretative cautions apply to the evidence rating of the relation between glioma/brain cancer and occupational RF exposure, due to differences in exposure sources and metrics across the few included studies.

Other This project was commissioned and partially funded by the World Health Organization (WHO). Co-financing was provided by the New Zealand Ministry of Health; the Istituto Superiore di Sanità in its capacity as a WHO Collaborating Centre for Radiation and Health; and ARPANSA as a WHO Collaborating Centre for Radiation Protection. Registration: PROSPERO CRD42021236798. Published protocol: [(Lagorio et al., 2021) DOI https://doi.org/10.1016/j.envint.2021.106828].

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0160412024005695?via%3Dihub

--

The effects of radiofrequency exposure on adverse female reproductive outcomes: A systematic review of human observational studies with dose–response meta-analysis (WHO SR 3 female)

My note: See WHO Radiofrequency EMF Health Risk Assessment Monograph (EHC series)

Johnson EE, Kenny RPW, Adesanya AM, Richmond C, Beyer F, Calderon C, Rankin J, Pearce MS, Toledano M, Craig D, Pearson F. The effects of radiofrequency exposure on adverse female reproductive outcomes: A systematic review of human observational studies with dose-response meta-analysis. Environ Int. 2024 Jun 12;190:108816. doi: 10.1016/j.envint.2024.108816.

Abstract

Background To inform radiofrequency electromagnetic field (RF-EMF) exposure guidelines the World Health Organization (WHO) is bringing together evidence on RF-EMF in relation to health outcomes prioritised for evaluation by experts in this field. Given this, a network of topic experts and methodologists have conducted a series of systematic reviews collecting, assessing, and synthesising data of relevance to these guidelines. Here we present a systematic review of the effect of RF-EMF exposure on adverse pregnancy outcomes in human observational studies which follows the WHO handbook for guideline development and the COSTER conduct guidelines.

Methods We conducted a broad, sensitive search for potentially relevant records within the following bibliographic databases: MEDLINE; Embase; and the EMF Portal. Grey literature searches were also conducted through relevant databases (including OpenGrey), organisational websites and via

consultation of RF-EMF experts. We included quantitative human observational studies on the effect of RF-EMF exposure in adults' preconception or pregnant women on pre-term birth, small for gestational age (SGA; associated with intrauterine growth restriction), miscarriage, stillbirth, low birth weight (LBW) and congenital anomalies. In blinded duplicate, titles and abstracts then full texts were screened against eligibility criteria. A third reviewer gave input when consensus was not reached. Citation chaining of included studies was completed. Two reviewers' data extracted and assessed included studies for risk of bias using the Office of Health Assessment and Translation (OHAT) tool. Random effects meta-analyses of the highest versus the lowest exposures and dose–response meta-analysis were conducted as appropriate and plausible. Two reviewers assessed the certainty in each body of evidence using the OHAT GRADE tool.

Results We identified 18 studies in this review; eight were general public studies (with the general public as the population of interest) and 10 were occupational studies (with the population of interest specific workers/workforces). General public studies. From pairwise meta-analyses of general public studies, the evidence is very uncertain about the effects of RF-EMF from mobile phone exposure on preterm birth risk (relative risk (RR) 1.14, 95% confidence interval (CI): 0.97–1.34, 95% prediction interval (PI): 0.83–1.57; 4 studies), LBW (RR 1.14, 95% CI: 0.96–1.36, 95% PI: 0.84–1.57; 4 studies) or SGA (RR 1.13, 95% CI: 1.02–1.24, 95% PI: 0.99–1.28; 2 studies) due to very low-certainty evidence. It was not feasible to meta-analyse studies reporting on the effect of RF-EMF from mobile phone exposure on congenital anomalies or miscarriage risk. The reported effects from the studies assessing these outcomes varied and the studies were at some risk of bias. No studies of the general public assessed the impact of RF-EMF exposure on stillbirth. Occupational studies. In occupational studies, based on dose–response meta-analyses, the evidence is very uncertain about the effects of RF-EMF amongst female physiotherapists using shortwave diathermy on miscarriage due to very low-certainty evidence (OR 1.02 95% CI 0.94–1.1; 2 studies). Amongst offspring of female physiotherapists using shortwave diathermy, the evidence is very uncertain about the effects of RF-EMF on the risk of congenital malformations due to very low-certainty evidence (OR 1.4, 95% CI 0.85 to 2.32; 2 studies). From pairwise meta-analyses, the evidence is very uncertain about the effects of RF-EMF on the risk of miscarriage (RR 1.06, 95% CI 0.96 to 1.18; very low-certainty evidence), pre-term births (RR 1.19, 95% CI 0.32 to 4.37; 3 studies; very low-certainty evidence), and low birth weight (RR 2.90, 95% CI: 0.69 to 12.23; 3 studies; very low-certainty evidence). Results for stillbirth and SGA could not be pooled in meta-analyses. The results from the studies reporting these outcomes were inconsistent and the studies were at some risk of bias.

Discussion Most of the evidence identified in this review was from general public studies assessing localised RF-EMF exposure from mobile phone use on female reproductive outcomes. In occupational settings, each study was of heterogenous whole-body RF-EMF exposure from radar, short or microwave diathermy, surveillance and welding equipment and its effect on female reproductive outcomes. Overall, the body of evidence is very uncertain about the effect of RF-EMF exposure on female reproductive outcomes. Further prospective studies conducted with greater rigour (particularly improved accuracy of exposure measurement and using appropriate statistical methods) are required to identify any potential effects of RF-EMF exposure on female reproductive outcomes of interest.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024004021

--

The effects of radiofrequency exposure on male fertility: A systematic review of human observational studies with dose-response meta-analysis (SR 3 male)

My note: See WHO Radiofrequency EMF Health Risk Assessment Monograph (EHC series)

Kenny RPW, Evelynne Johnson E, Adesanya AM, Richmond C, Beyer F, Calderon C, Rankin J, Pearce MS, Toledano M, Craig D, Pearson F. The effects of radiofrequency exposure on male fertility: A systematic review of human observational studies with dose-response meta-analysis. Environ Int. 2024 Jun 11;190:108817. doi: 10.1016/j.envint.2024.108817.

Abstract

Background: The World Health Organization (WHO) is bringing together evidence on radiofrequency electromagnetic field (RF-EMF) exposure in relation to health outcomes, previously identified as priorities for research and evaluation by experts in the field, to inform exposure guidelines. A suite of systematic reviews have been undertaken by a network of topic experts and methodologists to collect, assess and synthesise data relevant to these guidelines. Following the WHO handbook for guideline development and the COSTER conduct guidelines, we systematically reviewed the evidence on the potential effects of RF-EMF exposure on male fertility in human observational studies.

Methods: We conducted a broad and sensitive search for potentially relevant records within the following bibliographic databases: MEDLINE; Embase; Web of Science and EMF Portal. We also conducted searches of grey literature through relevant databases including OpenGrey, and organisational websites and consulted RF-EMF experts. We hand searched reference lists of included study records and for citations of these studies. We included quantitative human observational studies on the effect of RF-EMF exposure in adult male participants on infertility: sperm concentration; sperm morphology; sperm total motility; sperm progressive motility; total sperm count; and time to pregnancy. Titles and abstracts followed by full texts were screened in blinded duplicate against pre-set eligibility criteria with consensus input from a third reviewer as required. Data extraction from included studies was completed by two reviewers, as was risk of bias assessment using the Office of Health Assessment and Translation (OHAT) tool. We conducted a dose-response meta-analysis as possible and appropriate. Certainty of the evidence was assessed by two reviewers using the OHAT GRADE tool with input from a third reviewer as required.

Results: We identified nine studies in this review; seven were general public studies (with the general public as the population of interest) and two were occupational studies (with specific workers/workforces as the population of interest). General public studies. Duration of phone use: The evidence is very uncertain surrounding the effects of RF-EMF on sperm concentration (10/6 mL) (MD (mean difference) per hour of daily phone use 1.6 10^6/mL, 95 % CI -1.7 to 4.9; 3 studies), sperm morphology (MD 0.15 percentage points of deviation of normal forms per hour, 95 % CI -0.21 to 0.51; 3 studies), sperm progressive motility (MD -0.46 percentage points per hour, 95 % CI -1.04 to 0.13; 2 studies) and total sperm count (MD per hour -0.44 10^6/ejaculate, 95 % CI -2.59 to 1.7; 2 studies) due to very low-certainty evidence. Four additional studies reported on the effect of mobile phone use on sperm motility but were unsuitable for pooling; only one of these studies identified a statistically significant effect. All four studies were at risk of exposure characterisation and selection bias; two of confounding, selective reporting and attrition bias; three of outcome assessment bias and one used an inappropriate statistical method. Position of phone: There may be no or little effect of carrying a mobile phone in the front pocket on sperm concentration, total count, morphology, progressive motility or on

time to pregnancy. Of three studies reporting on the effect of mobile phone location on sperm total motility and, or, total motile count, one showed a statistically significant effect. All three studies were at risk of exposure characterisation and selection bias; two of confounding, selective reporting and attrition bias; three of outcome assessment bias and one used inappropriate statistical method. RF-EMF Source: One study indicates there may be little or no effect of computer or other electric device use on sperm concentration, total motility or total count. This study is at probably high risk of exposure characterisation bias and outcome assessment bias. Occupational studies. With only two studies of occupational exposure to RF-EMF and heterogeneity in the population and exposure source (technicians exposed to microwaves or seamen exposed to radar equipment), it was not plausible to statistically pool findings. One study was at probably or definitely high risk of bias across all domains, the other across domains for exposure characterisation bias, outcome assessment bias and confounding.

Discussion: The majority of evidence identified was assessing localised RF-EMF exposure from mobile phone use on male fertility with few studies assessing the impact of phone position. Overall, the evidence identified is very uncertain about the effect of RF-EMF exposure from mobile phones on sperm outcomes. One study assessed the impact of other RF-EMF sources on male fertility amongst the general public and two studies assessed the impact of RF-EMF exposure in occupational cohorts from different sources (radar or microwave) on male fertility. Further prospective studies conducted with greater rigour (in particular, improved accuracy of exposure measurement and appropriate statistical method use) would build the existing evidence base and are required to have greater certainty in any potential effects of RF-EMF on male reproductive outcomes. Prospero Registration: CRD42021265401 (SR3A).

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024004033

--

Response to letter from Bevington M., Electrosensitivity UK (WHO SR 8)

Bosch-Capblanch X, Esu E, Oringanje CM, Dongus S, Jalilian H, Eyers J, Auer C, Meremikwu M, Röösli M. Response to letter from Bevington M., Electrosensitivity UK. Environment International. 191, 2024. doi: 10.1016/j.envint.2024.108982.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024005683

--

Bevington Letter to Editor re: WHO systematic review of RF-EMF effects in human experimental studies of self-reported symptoms (WHO SR 8)

Bevington M. Letter to the Editor, Environment International 'Available evidence shows adverse symptoms from acute non-thermal RF-EMF exposure'. Comment on: Bosch-Capblanch X et al., The effects of radiofrequency electromagnetic fields exposure on human self-reported symptoms: A systematic review of human experimental studies, Envir Int. vol. 187, May 2024, 108612, Environment International, 2024, 108888, doi: 10.1016/j.envint.2024.108888.

No abstract

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024004744

--

The effects of radiofrequency electromagnetic field exposure on biomarkers of oxidative stress in vivo and in vitro: A systematic review of experimental studies (WHO SR 9)

My note: SR 9 is the eighth of the WHO SRs (systematic reviews) on RF-EMF effects to be published in *Environment International*.

Like its predecessors, the authors of this SR systematically excluded most of the relevant research. Most of the meta-analyses in this paper suffered from high levels of heterogeneity with little effort to understand the source of the varied results. Some meta-analyses were unreliable as they violated a key assumption of the random effects model, namely, independence of observations. I think a critique of this SR could build a strong case for retraction of this paper.

Meyer F, Bitsch A, Forman HJ, Fragoulis A, Ghezzi P, Henschenmacher B, Kellner R, Kuhne J, Ludwig T, Sachno D, Schmid G, Tsaioun K, Verbeek J, Wright R. The effects of radiofrequency electromagnetic field exposure on biomarkers of oxidative stress in vivo and in vitro: A systematic review of experimental studies. Environment International, 2024. doi: 10.1016/j.envint.2024.108940.

Highlights

The evidence for or against a relation between RF-EMF and biomarkers of oxidative stress is overall of very low certainty.

Inconsistent overall study results.

Need for drastic improvements of studies on RF-EMF and biomarkers of oxidative stress.

Abstract

Background

Oxidative stress is thought to be related to many diseases. Furthermore, it is hypothesized that radiofrequency electromagnetic fields (RF-EMF) may induce excessive oxidative stress in various cell types and thereby have the potential to compromise human and animal health. The objective of this systematic review (SR) is to summarize and evaluate the literature on the relation between the exposure to RF-EMF in the frequency range from 100 kHz to 300 GHz and biomarkers of oxidative stress.

Methods

The SR framework was developed following the guidelines established in the WHO Handbook for Guideline Development and NTP/OHAT's Handbook for Conducting a Literature-Based Health Assessment. We used the latter handbook's methodology for implementing the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach for environmental health assessments. We searched the following databases up until June 30, 2023: PubMed, Embase, Web of Science Core Collection, Scopus, and the EMF-Portal. The reference lists of included studies and retrieved review articles were also manually searched. We rated Risk of Bias (RoB) using the OHAT RoB Rating Tool and assessed publication bias using funnel plots of included studies. We assessed the certainty of the evidence (high, moderate, low, or inadequate) for an association between RF-EMF and

oxidative stress using an adapted version of the GRADE framework. Data were extracted according to a predefined set of forms developed in DistillerSR. Data were analysed after grouping them first as in vitro or in vivo and then according to outcome category, species category, and exposed tissue. We synthesized study results using a random effects meta-analysis when study characteristics were judged sufficiently similar to be combined and heterogeneity (I2) was lower than 75 %, otherwise we describe the findings narratively.

Results

Fifty-six (56) studies, 45 in vivo and 11 in vitro, exposed to frequencies in the range 800–2450 MHz, were included in the SR after eliminating 11,543 publications because they did not meet the criteria defined in the published protocol (Henschenmacher et al., 2022). Of 56 studies 51 studies with 168 individual results were included in the meta-analysis. Together, these studies examined six human in vitro samples and fifty animal samples, including rodents (mice, rats, hamsters, and guinea pigs, [n = 46]) and rabbits [n = 4]. RF-EMF were predominantly applied as continuous wave exposures in these studies. The outcome biomarkers for modified proteins and amino acids were measured in 30 studies, for oxidized DNA bases in 26 studies, for oxidized lipids in 2 studies and hydrogen peroxide production in 2 studies. Outcomes were mostly measured in the brain (n = 22), liver (n = 9), cells (n = 9), blood (n = 6), and testis (n = 2). RoB in studies was high, mainly due to biases in exposure and outcome assessment.

In vivo studies

Brain: The effect on biomarkers for oxidized DNA bases in the rodent brain (five studies, n = 98) had an inconsistent effect, varying from a large decrease with a standardized mean difference (SMD) of −3.40 (95 % CI [−5.15, −1.64]) to a large increase with an SMD of 2.2 (95 % CI [0.78, 3.62]). In the brain of rabbits (two studies, n = 44), the effect sizes also varied, from an SMD of −1.06 (95 % CI [−2.13, 0.00]) to an SMD of 5.94 (95 % CI [3.14, 8.73]). The effect on biomarkers for modified proteins and amino acids in the rodent brain (15 studies, n = 328) also varied from a large decrease with an SMD of −6.11 (95 % CI [−8.16, −4.06]) to a large increase with an SMD of 5.33 (95 % CI [2.49, 8.17]).

The effect on biomarkers for oxidized lipids in the brain of rodents (one study, n = 56) also varied from a large decrease with SMD = −4.10 (95 % CI [−5.48, −2.73]) to SMD = 1.27 (95 % CI [0.45, 2.10]).

Liver: The effect on biomarkers for oxidized DNA bases in the rodent liver (two studies, n = 26) was inconsistent with effect sizes in both directions: SMD = −0.71 (95 % CI [−1.80, 0.38]) and SMD = 1.56 (95 % CI [0.19, 2.92]). The effect on biomarkers for oxidized DNA bases in the rabbits' liver (two studies, n = 60) was medium with a pooled SMD of 0.39 (95 % CI [−0.79, 1.56]).

Biomarkers for modified proteins and amino acids in the liver of rodents (six studies, n = 159) increased with a pooled SMD of 0.55 (95 % CI [0.06, 1.05]).

Blood: The effect of RF-EMF on biomarkers for oxidized DNA bases in rodent blood (four studies, n = 104) was inconsistent, with SMDs ranging from −1.14 (95 % CI [−2.23, −0.06]) to 1.71 (95 % CI [−0.10, 3.53]).

RF-EMF had no effect on biomarkers for modified proteins and amino acids in rodent blood (three studies, n = 40), with a pooled SMD of −0.08 (95 % CI [−1.32, 1.16]).

There was a large increase in biomarkers for oxidized DNA bases in rodent plasma (two studies, n = 38) with a pooled SMD of 2.25 (95 % CI [1.27, 3.24]).

Gonads: There was an increase in biomarkers for oxidized DNA bases in the rodent testis (two studies, n = 24) with a pooled SMD of 1.60 (95 % CI [0.62, 2.59]).

The effect of RF-EMF on biomarkers for modified proteins and amino acids in the ovary of rodents (two studies, n = 52) was inconsistent with a small effect, SMD = 0.24 (95 % CI [−0.74, 1.23])) and a large effect (SMD = 2.08 (95 % CI [1.22, 2.94])).

Thymus: RF-EMF increased biomarkers for modified proteins and amino acids in the thymus of rodents (one study, n = 42) considerably with a pooled SMD of 6.16 (95 % CI [3.55, 8.76]).

Cells: RF-EMF increased oxidized DNA bases in rodent cells with SMD of 2.49 (95 % CI [1.30, 3.67]) (one study, n = 27). There was a small effect in oxidized lipids (one study, n = 18) but not statistically significant with an SMD of 0.34 (95 % CI [−0.62, 1.29]).

In vitro studies

In in vitro studies in human cells (three studies, n = 112), there were inconsistent increases in biomarkers for oxidized DNA bases, where the SMDs varied between 0.01 (95 % CI [−0.59, 0.62]) and 7.74 (95 % CI [2.24, 13.24]) in 4 results (2 of them statistically significant). In rodent cells (three studies, n = 24), there was a not statistically significant large effect in biomarkers for oxidized DNA bases with SMD = 2.07 (95 % CI [−1.38, 5.52]).

The RF-EMF biomarkers for modified proteins and amino acids in human cells (one study, n = 18) showed a large effect with SMD = 1.07 (95 % CI [−0.05, 2.19]). In rodent cells (two studies, n = 24) a medium effect of SMD = 0.56 (95 % CI [−0.29, 1.41]) was observed.

Discussion

The evidence on the relation between the exposure to RF-EMF and biomarkers of oxidative stress was of very low certainty, because a majority of the included studies were rated with a high RoB level and provided high heterogeneity. This is due to inaccurate measurements of exposure and/or of measurement of oxidative stress biomarkers and missing information on the blinding of the research personnel. There may be no or an inconsistent effect of RF-EMF on biomarkers of oxidative stress in the brain, liver, blood, plasma and serum, and in the female reproductive system in animal experiments but the evidence is of very low certainty. There may be an increase in biomarkers of oxidative stress in testes, serum and thymus of rodents but the evidence is of very low certainty. Future studies should improve experimental designs and characterization of exposure systems as well as the use of validated biomarker measurements with positive controls.

Other: This review was partially funded by the World Health Organization. The protocol for this review is registered in PROSPERO (https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021235573) and published in Environment International (https://doi.org/10.1016/j.envint.2021.106932) (Henschenmacher et al., 2022).

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024005269

--

Occupational exposure to radiofrequency electromagnetic fields and brain tumor risk: application of the INTEROCC job-exposure matrix

Maxime Turuban, Hans Kromhout, Javier Vila, Miquel Vallbona-Vistos, Frank de Vocht , Baldi, L Richardson, G Benke, D Krewski, E Parent, S Sadetzki, B Schlehofer, J Schuz, J Siemiatycki, M van Tongeren, A Woodward, E Cardis, M Turner. Occupational exposure to radiofrequency electromagnetic fields and brain tumor risk: application of the INTEROCC job-exposure matrix. International J Cancer. First published: 20 September 2024. doi: 10.1002/ijc.35182.

Abstract

Radiofrequency electromagnetic fields (RF-EMF, 100 kHz to 300 GHz) are classified by IARC as possibly carcinogenic to humans (Group 2B). This study evaluates the potential association between occupational RF-EMF exposure and brain tumor risk, utilizing for the first time, a RF-EMF job-exposure matrix (RF-JEM) developed in the multi-country INTEROCC case-control study. Cumulative and time-weighted average occupational RF-EMF exposures were estimated for study participants based on lifetime job histories linked to the RF-JEM using three different methods: 1) by considering RF-EMF intensity among all exposed jobs, 2) by considering RF-EMF intensity among jobs with an exposure prevalence ≥ the median exposure prevalence of all exposed jobs, and 3) by considering RF-EMF intensity of jobs of participants who reported RF-EMF source use. Stratified conditional logistic regression models were used, considering various lag periods and exposure time windows defined a priori. Generally, no clear associations were found for glioma or meningioma risk. However, some statistically significant positive associations were observed including in the highest exposure categories for glioma for cumulative and time-weighted average exposure in the 1-4 year time window for electric fields (E) in the first JEM application method (ORs = 1.36, 95% CI 1.08, 1.72 64 and 1.27, 95% CI 1.01, 1.59 respectively), as well as for meningioma for cumulative exposure in the 5-9 year time window for electric fields (E) in the third JEM application method (OR = 2.30, 95% CI 1.11, 4.78). We did not identify convincing associations between occupational RF-EMF exposure and risk of glioma or meningioma.

What's New?

Radiofrequency electromagnetic fields have been classified as possibly carcinogenic to humans. Here, the authors applied a novel job-exposure matrix to investigate the link between occupational radiofrequency electromagnetic field exposure and brain tumors within the multi-country INTEROCC case–control study. The findings revealed no clear associations, despite some positive findings for gliomas and meningiomas in the 1- to 4- and 5- to 9-year exposure windows. The study underlines the applicability of the job-exposure matrix in occupational research and the need to further investigate the association between brain tumors and occupational radiofrequency electromagnetic field exposure

Conclusions

The previous SEM-based and the current JEM-based analyses (with three ways of linking the JEM to participant job histories) produced similar results that fail to show clear trends of associations between various RF-EMF exposure measures and either glioma or meningioma risk. Nevertheless, interpretation of current findings remains challenging. The method of linking JEM estimates to a job only when RF source use was self-reported led to significantly lower exposure prevalences and similarly did not suggest clear evidence of increased risk from occupational IF-RF/RF-EMF exposures. Findings suggest that the RF-JEM could potentially be used in studies without self-reported information available on working with or near RF sources.

While we generally found no associations or inverse associations overall, a few positive associations between exposure to IF-RF/RF in cumulative and TWA exposure analyses were found with glioma in Method 1 in the highest exposure category in the 1- to 4-year time window of exposure, and with meningioma in Method 3 in the 5- to 9-year window prior to diagnosis/reference date. However, these statistically significant results, not consistent across analysis methods, could be due to chance, and will require verification in other independent studies. Integrating additional criteria when using JEM, such as exposure prevalence similar to Method 2 in this paper, when attributing exposure levels could be considered in further work. Future improvements of the RF-JEM, including by collecting and integrating personally measured exposure data could be considered.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/ijc.35182

--

Link between Wi-Fi, cordless devices, mobile phone usage patterns, and behavioral problems among Japanese children: A prospective cohort study

Ajmal A, Yamazaki K, Tamura N, Ait Bamai Y, Yoshikawa T, Hikage T, Ikeda A, Kishi R. Link between Wi-Fi, cordless devices, mobile phone usage patterns, and behavioral problems among Japanese children: A prospective cohort study. Environ Res. 2024 Aug 1:119715. doi: 10.1016/j.envres.2024.119715.

Highlights

- Persistence and change in behavioral problem over one year were assessed and grouped into normal, persistent, improved, and concurrent groups.
- Above 75th percentile mobile phone calls duration via Internet was associated with concurrent total difficulties.
- Cordless phone use at home was associated with improved total difficulties.
- Longer cordless phone call durations were less likely to lead to persistent problematic prosocial behavior.
- The observed findings may not be related to RF-EMF and could be affected by residual confounding and chance findings.

Abstract

Background: With the recent advent of technology, it is important to confirm the health and safety of the youth. This study aimed to prospectively evaluate the relationship between Wi-Fi, cordless phones,

and mobile phone usage patterns and behavioral problems.

Methods: This study involved 2,465 children aged 8-17 years from the Hokkaido Study on Environment and Children's Health from October 2020 to January 2021, with a follow-up from September 2021 to March 2022. The mother-child dyad provided information on the presence of residential Wi-Fi and cordless phones, cordless phone call duration, and mobile phone usage pattern (duration of calls using mobile network and internet, online audio streaming, online video streaming, and playing online games) via a baseline questionnaire. Based on the scores on Strength and Difficulties Questionnaire at baseline and follow-up,the children were categorized into four groups: normal, persistent, improved, and concurrent.

Results: No significant association was found between Wi-Fi, mobile phone calls via mobile networks, and behavioral problems. Cordless phone at home had higher odds for improvement in total difficulty scores, and cordless phone for calling more than 4 minutes per week had lower odds of persistent problematic prosocial behavior. Longer duration of mobile phone calling via the internet (>40 min/week) had higher odds of concurrent total difficulties. Mobile phone calling via mobile network for <5 minutes per week had higher odds for improved total difficulty scores. Audio streaming via mobile phones for 60-120 minutes had lower odds of persistent total difficulties.

Conclusion: Our results showed sporadic findings between residential RF-EMF indoor sources and mobile phone usage pattern. These observed findings could be affected by residual confounding and chance findings. Ongoing follow-up studies are necessary to further explore this association through detailed exposure assessment and addressing the potential limitations of our study.

https://pubmed.ncbi.nlm.nih.gov/39096992/

--

Evaluating the Effect on Heart Rate Variability of Adults Exposed to Radio-Frequency Electromagnetic Fields in Modern Office Environment

Dale S, Reiz R, Popa S, Ardelean-Dale A, Keller J, Geier JU. Evaluating the Effect on Heart Rate Variability of Adults Exposed to Radio-Frequency Electromagnetic Fields in Modern Office Environment. International Journal of Advanced Computer Science and Applications (IJACSA), 15(6), 2024. doi: 10.14569/IJACSA.2024.0150609.

Abstract

The objective of the study was to investigate whether heart rate variability (HRV) is an appropriate method to describe potential effects of RF-EMF on humans considering a modern office environment radiation level with the frequencies 1.8 GHz (DECT) and 2.45 GHz (Wi-Fi) and an exposure time of 10 min. The emitters were 1 m distant from the test subjects. The HRV parameters SDNN, RMSSD, LF and HF were recorded from 60 adults in three runs, totaling up to 154 recordings. Effects were evident for the parameter SDNN. In two runs, HRV changed from control to exposure phase, in one run from exposure phase to control. The cofactors smoking, coffee consumption, and the use of strong medications did not modulate EMF effects. HRV seems to be suitable to detect effects of radio-frequency electromagnetic fields on humans under certain conditions. In the future, prolonged exposure

and new frequencies (5G) should be included in order to provide a better description of RF-EMF effects in modern office environments.

http://dx.doi.org/10.14569/IJACSA.2024.0150609.

--

Hypersensitivity to man-made electromagnetic fields (EHS) correlates with immune responsivity to oxidative stress: a case report

Thoradit T, Chabi M, Aguida B, Baouz S, Stierle V, Pooam M, Tousaints S, Akpovi CD, Ahmad M. Hypersensitivity to man-made electromagnetic fields (EHS) correlates with immune responsivity to oxidative stress: a case report. Commun Integr Biol. 2024 Aug 4;17(1):2384874. doi: 10.1080/19420889.2024.2384874.

Abstract

There is increasing evidence that exposure to weak electromagnetic fields (EMFs) generated by modern telecommunications or household appliances has physiological consequences, including reports of electromagnetic field hypersensitivity (EHS) leading to adverse health effects. Although symptoms can be serious, no underlying mechanism for EHS is known and there is no general cure or effective therapy. Here, we present the case study of a self-reported EHS patient whose symptoms include severe headaches, generalized fatigue, cardiac arrhythmia, attention and memory deficit, and generalized systemic pain within minutes of exposure to telecommunications (Wifi, cellular phones), high tension lines and electronic devices. Tests for cerebral, cardiovascular, and other physiological anomalies proved negative, as did serological tests for inflammation, allergies, infections, auto-immune conditions, and hormonal imbalance. However, further investigation revealed deficits in cellular anti-oxidants and increased radical scavenging enzymes, indicative of systemic oxidative stress. Significantly, there was a large increase in circulating antibodies for oxidized Low-Density Lipoprotein (LDLox), byproducts of oxidative stress accumulating in membranes of vascular cells. Because a known primary effect of EMF exposure is to increase the concentration of cellular oxidants, we propose that pathology in this patient may be causally related to a resulting increase in LDLox synthesis. This in turn could trigger an exaggerated auto-immune response consistent with EHS symptoms. This case report thereby provides a testable mechanistic framework for EHS pathology with therapeutic implications for this debilitating and poorly understood condition.

Open access paper: https://www.tandfonline.com/doi/full/10.1080/19420889.2024.2384874

--

Effects of 5G radiofrequency electromagnetic radiation on indicators of vitality and DNA integrity of in vitro exposed boar semen

Butković I, Vince S, Lojkić M, Folnožić I, Tur SM, Vilić M, Malarić K, Berta V, Samardžija M, Kreszinger M, Žaja IŽ. Effects of 5G radiofrequency electromagnetic radiation on indicators of vitality and DNA integrity of in vitro exposed boar semen. Theriogenology. 2024 Sep 25;230:243-249. doi: 10.1016/j.theriogenology.2024.09.025.

Abstract

The effects of radiofrequency electromagnetic radiation (RF-EMR) on semen quality have been in the spotlight in recent years, though research results to date have been contradictory. The effects of RF-EMR amongst others depend upon frequency, and there is currently no literature concerning the influence of 5G frequencies on both DNA integrity and spermatozoa vitality in males. The aim of this study was to investigate the effect of 5G RF-EMR on sperm membrane integrity, mitochondrial potential, and DNA integrity of in vitro exposed semen of breeding boars. The study included semen samples of eight breeding boars of the Pietren breed and four breeding boars of the German Landrace breed, from 1.5 to 3.5 years in age. Freshly diluted semen of each boar was divided into a control (n = 12) and experimental group (n = 12). The samples of the experimental group were exposed for 2 hours to continuous RF-EMR at a single frequency (700 MHz, 2500 MHz and 3500 MHz) and an electromagnetic field strength of 10 V/m using a transverse gigahertz electromagnetic cell. Sperm DNA fragmentation was assessed using a Halomax® kit and sperm membrane integrity and mitochondrial potential was assessed using a PI/SYBR-14 LIVE/DEAD viability kit with JC-1. A significantly higher proportion of spermatozoa with DNA fragmentation was found in exposed semen samples for all frequencies compared to the control group. The highest DNA damage was recorded in semen samples exposed to 5G RF-EMR at 2500 MHz ($p < 0.01$) and 3500 MHz ($p < 0.05$) vs. control semen samples. A significantly higher proportion of spermatozoa with damaged cell membrane and good mitochondrial potential was recorded in semen samples exposed with 3500 MHz. In vitro exposure of breading boar semen to 5G RF-EMR significantly increases the proportion of DNA fragmentation. The harmful effect of 5G RF-EMR on the proportion of spermatozoa with damaged DNA was frequency dependent. The 3500 MHz frequency displayed the most harmful effects due to significant impacts on DNA integrity and spermatozoa vitality indicators.

https://pubmed.ncbi.nlm.nih.gov/39342826/

--

Study of genotoxic and cytotoxic effects induced in human fibroblasts by exposure to pulsed and continuous 1.6 GHz radiofrequency

Massaro L, De Sanctis S , Franchini V, Regalbuto E, Alfano G, Focaccetti C, Benvenuto M, Cifaldi L, Sgura A, Berardinelli F, Marinaccio J, Barbato F, Rossi E, Nardozi D, Masuelli L, Bei R, Lista F. Study of genotoxic and cytotoxic effects induced in human fibroblasts by exposure to pulsed and continuous 1.6 GHz radiofrequency. Frontiers in Public Health, Vol 12. 2024. doi: 10.3389/fpubh.2024.1419525.

Abstract

Background The widespread use of radiofrequency (RF) sources, ranging from household appliances to telecommunications devices and military equipment, raises concerns among people and regulatory agencies about the potential health risks of RF exposure. Consequently, several *in vitro* and *in vivo* studies have been done to investigate the biological effects, in particular non-thermal, of this non-ionizing radiation. To date, this issue is still being debated due to the controversial results that have been reported. Furthermore, the impact of different RF signal modulations on biological systems remains poorly investigated. The present *in vitro* study aims to evaluate the cytotoxicity and genotoxicity of continuous or pulsed 1.6 GHz RF in human dermal fibroblasts (HDF).

Methods HDF cultures were exposed to continuous and pulsed 1.6 GHz RF, for 2 h, with Specific Absorption Rate (SAR) of 0.4 W/kg. The potential biological effects of 1.6 GHz RF on HDF were assessed

with a multi-methodological approach, analyzing the effects on cell cycle, ultrastructure, protein expression, mitotic spindle, CREST stained micronuclei, chromosome segregation and γ-H2AX/53BP1 foci.

Results 1.6 GHz RF exposure modified proteins expression and morphology of HDF. Specifically, the expression of different heat-shock proteins (HSP) (i.e., HSP-90, HSP-60, and HSP-25) and phospho-AKT were affected. In addition, both continuous and pulsed RF modified the cytoskeletal organization in HDF and increased the number of lysosomes, while the formation of autophagosomes was observed only after pulsed RF exposure. Mitotic spindle anomalies were also found after exposure. However, no significant effect was observed on cell cycle, chromosome segregation, CREST-stained micronuclei and γ-H2AX/53BP1 foci.

Conclusion The results of the present study show the absence of genotoxic damage in 1.6 GHz RF exposed HDF and, although mitotic spindle alterations were observed, they did not have an aneugenic effect. On the other hand, changes in some proteins expression and cell ultrastructure in exposed HDF suggest that RF can potentially induce cell alterations at the morphological and molecular levels.

Excerpt

In this controversial scientific scenario, our results are in agreement with most *in vitro* findings on different cell models about the lack of genotoxic damage induced by RF, either clastogenic or aneugenic. Despite this, we find alterations of the mitotic spindle, with a significant increase in multipolar spindles following PW exposure and we also observed a tendency of RF to induce non-disjunction events with both signals. However, our study reveals an increase in these spindle abnormalities without a concomitant increase in MN-positive CREST, suggesting no aneuploidy effect probably due to spindle abnormalities reversion, as proposed by other authors (78). Nonetheless, this apparent discordant result requires further investigations.

The reported observations indeed highlight the complexity of cellular response to RF and emphasize the need for further investigations to clarify the overall biological effects of 1.6 GHz PW and CW RF, given the widespread and constant public RF-EMF exposure, mostly due to mobile communication devices.

Open access paper: https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2024.1419525

--

Pulsed High-Power Radio Frequency Energy Can Cause Non-Thermal Harmful Effects on the BRAIN

Yaghmazadeh O. Pulsed High-Power Radio Frequency Energy Can Cause Non-Thermal Harmful Effects on the BRAIN. IEEE Open Journal of Engineering in Medicine and Biology, 5:50-53, 2024. doi: 10.1109/OJEMB.2024.3355301.

Abstract

High-power microwave applications are growing for both military and civil purposes, yet they can induce brain-related risks and raise important public health concerns. High-power sub-millisecond radio frequency energy pulses have been demonstrated to be able to induce neurological and

neuropathological changes in the brain while being compliant with current regulatory guidelines' limits, highlighting the necessity of revising them.

Impact Statement

This commentary piece looks into the different mechanisms of interaction of pulsed versus continuous-wave RF energy radiation and the brain. It also highlights potential harmful effects of pulsed high-power RF energy on the brain.

Open access paper: https://ieeexplore.ieee.org/document/10402080

--

External RF-EMF alters cell number and ROS balance possibly via the regulation of NADPH metabolism and apoptosis

Chow S-C, Zhang Y, Ng RWM , Hui S-YR , Solov'yov IA , Lui W-Y. External RF-EMF alters cell number and ROS balance possibly via the regulation of NADPH metabolism and apoptosis. Frontiers in Public Health. 12 (2024). doi: 10.3389/fpubh.2024.1425023.

Abstract

The influence of weak radio-frequency electromagnetic field (RF-EMF) on living organisms raises new concern because of the Industrial, Scientific, and Medical (ISM) frequency band at 6.78 MHz being promoted by the AirFuel Alliance for mid-range wireless power transfer (WPT) applications and product development. Human exposure to the RF-EMF radiation is unavoidable. In this study, we employed *in vitro* cell culture and molecular biology approach coupled with integrated transcriptomic and proteomic analyses to uncover the effects of RF-EMF on cells at molecular and cellular levels. Our study has demonstrated that weak RF-EMF is sufficient to exert non-thermal effects on human umbilical vein endothelial cells (HUVEC). Exposure of weak RF-EMF promotes cell proliferation, inhibits apoptosis and deregulates ROS balance. Alteration of several signaling pathways and key enzymes involved in NADPH metabolism, cell proliferation and ferroptosis were identified. Our current study provide solid evidence for the first time that the present safety standards that solely considered the thermal effect of RF-EMF on cell tissue are inadequate, prompt response and modification of existing Guidelines, Standards and Regulation are warranted.

Open access paper: https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2024.1425023/full

--

No observable non-thermal effect of microwave radiation on the growth of microtubules

Hammarin G, Norder P, Harimoorthy R, Chen G, Berntsen P, Widlund PO, Stoij C, Rodilla H, Swenson J, Brändén G, Neutze R. No observable non-thermal effect of microwave radiation on the growth of microtubules. Sci Rep. 2024 Aug 7;14(1):18286. doi: 10.1038/s41598-024-68852-3.

Abstract

Despite widespread public interest in the health impact of exposure to microwave radiation, studies of the influence of microwave radiation on biological samples are often inconclusive or contradictory. Here we examine the influence of microwave radiation of frequencies 3.5 GHz, 20 GHz and 29 GHz on the growth of microtubules, which are biological nanotubes that perform diverse functions in eukaryotic cells. Since microtubules are highly polar and can extend several micrometres in length, they are predicted to be sensitive to non-ionizing radiation. Moreover, it has been speculated that tubulin dimers within microtubules might rapidly toggle between different conformations, potentially participating in computational or other cooperative processes. Our data show that exposure to microwave radiation yields a microtubule growth curve that is distorted relative to control studies utilizing a homogeneous temperature jump. However, this apparent effect of non-ionizing radiation is reproduced by control experiments using an infrared laser or hot air to heat the sample and thereby mimic the thermal history of samples exposed to microwaves. As such, no non-thermal effects of microwave radiation on microtubule growth can be assigned. Our results highlight the need for appropriate control experiments in biophysical studies that may impact on the sphere of public interest.

Open access paper: https://www.nature.com/articles/s41598-024-68852-3

--

Exploring edible bird nest's potential in mitigating Wi-Fi's impact on male reproductive health

Maluin SM, Jaffar FHF, Osman K, Zulkefli AF, Mat Ros MF, Ibrahim SF. Exploring edible bird nest's potential in mitigating Wi-Fi's impact on male reproductive health. Reprod Med Biol. 2024 Sep 11;23(1):e12606. doi: 10.1002/rmb2.12606.

Abstract

Purpose: This study aimed to evaluate the protective effects of edible bird nest (EBN) against the detrimental impact of Wi-Fi on male reproductive health. Specifically, it examines whether EBN can mitigate Wi-Fi-induced changes in male reproductive hormones, estrogen receptors (ER), spermatogenesis, and sperm parameters.

Methods: Thirty-six adult male rats were divided into six groups (n = 6): Control, Control EBN, Control E2, Wi-Fi, Wi-Fi+EBN, and Wi-Fi+E2. Control EBN and Wi-Fi+EBN groups received 250 mg/kg/day EBN, while Control E2 and Wi-Fi+E2 groups received 12 μg/kg/day E2 for 10 days. Wi-Fi exposure and EBN supplementation lasted eight weeks. Assessments included organ weight, hormone levels (FSH, LH, testosterone, and E2), ERα/ERβ mRNA and protein expression, spermatogenic markers (c-KIT and SCF), and sperm quality.

Results: Wi-Fi exposure led to decreased FSH, testosterone, ERα mRNA, and sperm quality (concentration, motility, and viability). EBN supplementation restored serum FSH and testosterone levels, increased serum LH levels, and the testosterone/E2 ratio, and normalized mRNA ERα expression. Additionally, EBN increased sperm concentration in Wi-Fi-exposed rats without affecting motility or viability.

Conclusions: EBN plays a crucial role in regulating male reproductive hormones and spermatogenesis,

leading to improved sperm concentration. This could notably benefit men experiencing oligospermia due to excessive Wi-Fi exposure.

Excerpts

Wi-Fi exposure setting: For Wi-Fi exposure, this study utilized the TP-LINK AC750 Wireless Dual Band Wi-Fi Router Archer C20 (Shenzhen, China). This router features three external antennas, emitting signals at a frequency of 2.45 GHz using the IEEE 802.11n standard. The router was positioned 20 cm away from the rat cages and constantly exchanged data with a Raspberry Pi device through a ping protocol.3 The router was selected for its minimal vibration and noise output, which is typical for standard Wi-Fi routers used in laboratory settings. Additionally, the rats were housed in a controlled environment designed to minimize external disturbances, ensuring the reliability of our findings related to Wi-Fi exposure effects on male reproductive health.

Conclusion: This study highlights the negative impact of eight weeks of Wi-Fi exposure on sperm quality, including decreased concentration, motility, and viability, which can be ascribed to changes in male reproductive hormones. EBN supplementation appears to be a preventive intervention, significantly increasing gonadotrophin and testosterone levels, as well as sperm concentration in Wi-Fi-exposed rats, but not influencing sperm motility or viability. Furthermore, it increases the T/E2 ratio and restores estrogenic activity in the testes, resulting in enhanced sperm concentration. The study also reveals a discrepancy between the impacts of EBN supplementation and E2 treatment on sperm concentration in a Wi-Fi-exposed setting, emphasizing EBN's unique protective properties without adverse effects on male reproduction. In conclusion, EBN supplementation effectively restores spermatogenesis capabilities that are affected by Wi-Fi-induced damage. This is achieved through the modulation of male reproductive hormones, with a primary influence on sperm concentration. Nevertheless, further research is necessary to fully understand the mechanisms and establish safe usage limits for maximizing the benefits of EBN while minimizing potential risks.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/rmb2.12606

--

Spots with Extremely High Radiofrequency Radiation After Deployment of 5G Base Stations in Stockholm, Sweden

Hardell L, Koppel T. Spots with Extremely High Radiofrequency Radiation After Deployment of 5G Base Stations in Stockholm, Sweden. Ann Clin Med Case Rep. 2024; V14(4): 1-8.

Abstract

During recent years there is an on-going deployment of 5G base stations for radiofrequency [RF] communication in Sweden as well in many other countries. This is made without investigations on risks to human health and the environment. Since 2016 we have made several measurements of environmental exposure to RF radiation in Stockholm, Sweden, including previous generations and now also 5G. In the current study, performed in October 2023, the broadband meter Narda-550 with the probe EF-1891 was used. It gives results in the root mean square [RMS] mode, thus not peak levels. Both

the International Commission on Non-Ionizing Radiation Protection [ICNIRP] and the US Federal Communications Commission [FCC] base their guidelines on RMS levels of RF radiation. Measurements were made in similar city areas as in our previous studies. Results show that the RF radiation has increased substantially. High maximum levels were found at e.g. the Central Railway Station [3 637 191 µW/m2], and popular walking streets such as Mäster Samuelsgatan [11 613 976 µW/m2], and Drottninggatan [5 271 555 µW/m2]. The peak values would be much higher. These levels are in the same magnitude or even higher than those measured in homes of persons that rapidly developed symptoms of the microwave syndrome after installation of 5G, usually in combination with 4G+, in the neighborhood. They are also substantially higher than levels that have previously been linked to cancer and the microwave syndrome in studies of people living near base stations and masts from previous generations of telecommunications.

Open access paper: https://acmcasereport.org/wp-content/uploads/2024/09/ACMCR-v14-2262-3-1.pdf

--

Mobile Phone Emissions in 5G FR1: Using Statistic Inferences and Deep Learning for Empiric Features Extraction

Miclaus S, Deaconescu DB, Vatamanu D, Buda AM. Mobile Phone Emissions in 5G FR1: Using Statistic Inferences and Deep Learning for Empiric Features Extraction*. 2024 IEEE International Symposium on Measurements & Networking (M&N)*, Rome, Italy, 2024, pp. 01-06, doi: 10.1109/MN60932.2024.10615263.

Abstract

As the primary source of human exposure to electromagnetic fields, the emission of a 5G mobile phone is quantified for several mobile applications use. Following the time variability of the exposure, its peak levels and their statistic distributions, together with the time-frequency change in the spectrograms, we emphasize the peculiarities of the exposure in connection to modulation scheme used and to the application. The results show notable differences in amplitude probability densities, in complementary cumulative density functions and in repeatability of time-frequency features of the emissions.

Excerpts

It has been recently proved that the main exposure source of a person is the mobile phone itself [5], [6]. The study results of [7] confirmed that the human exposure in a 5G network is dominated by the uplink, and can be ten times larger than the downlink exposure....

Conducting measurements in real-life situations contributes to true assessment of the field exposure impact. It is essential to interpret all the findings on the background of dosimetric determinations and based on present regulatory guidelines [16], [17]. However, studies like [18] - [20] clearly pointed out that “cell phones emit a succession of pulse trains of different durations, the instantaneous fields of such signals allow activation of molecular, electronic, and protonic components within cells that have different thresholds and relaxation times, enriching the non-thermal effects of radiofrequency fields”.

In the current work we aimed to analyze the exposure situation generated by a 5G mobile phone in a static position, while emitting signals in a 100 MHz bandwidth centered on 3.58 GHz. While using four different mobile applications running on the phone ...

An Iphone 14pro (model A 2890, Apple, Zhengzhou, China) was the source of the emissions. It was maintained in a stationary position and connected to the 5G mobile network - Orange Romania operator. The central frequency was f=3.58 GHz, the bandwidth was 100 MHz and Time Division Multiple Access (TDMA) scheme was used....

... the phone was situated at the cell edge and it received a good downlink signal.

The receiving system was composed of a signal analyzer model FSW from Rhode & Schwarz, with a real time bandwidth of 160 MHz connected to a receiving antenna model Aaronia Omnilog 30800 (Fig. 1)....

... video streaming conducted to the highest mean emitted powers, while upload provokes the highest peak powers. Crest factors are the largest when a file is uploaded, and they can exceed 20 dB....

Variations of 19 dB around mean power is equivalent to a factor of approximately 8 x 10**3. Such a large variability during so short time (ms), may pose challenging biological response. The trends in the excessive exceedance of the mean power depend at least on mobile application type and on the modulation scheme. From this perspective, the tail features of time-variability of the exposure level may trigger unexpected consequences....

CONCLUSIONS

A mobile phone connected to a 5G-NR network in FR1 band was investigated for its emitted EMF level and the associated time-variability characterization (during 250 ms), in four situations of usage: during file-upload, file-download, video call and Internet streaming. The measurement set-up consisted in a receiving antenna mostly planar and parallel to the phone’s display connected to a vector signal analyzer with a capability of analysing bandwidth larger than the communication bandwidth (100 MHz). The antenna collected the field of the phone at 30 cm distance from it, and had an omnidirectional pattern in azimuth....

All the results showed that the uplink EMF level is far below the safe limit based on thermal effects of human exposure. However, other important features evolved and they may be further linked to the non-thermal/specific effects of microwaves....

When it comes to the prevalence of peaks in the exposure, it resulted that crest factors varied mostly in function of the mobile application used but also, in a lower measure, in function on the modulation scheme. The largest crest factors were more than one order of magnitude higher than the smallest ones, overall. Mean and peak powers of emissions could deviate from each other with approximately two orders of magnitude, for the same mobile application. Such short and pronounced extremes should be carefully investigated from the biological endpoint of view.

--

Estimates and measurements of radiofrequency exposures in smart-connected homes

Joyner K, Milligan M, Knipe P. Estimates and measurements of radiofrequency exposures in smart-connected homes. Bioelectromagnetics. 2024 Jul 18. doi: 10.1002/bem.22518.

Abstract

The aim of this research was to quantify the levels of radiofrequency electromagnetic energy (RF-EME) in a residential home/apartment equipped with a range of wireless devices, often referred to as internet of things (IoT) devices or smart devices and subsequently develop a tool that could be useful for estimating the levels of RF-EME in a domestic environment. Over the course of 3 years measurements were performed in peoples' homes on a total of 43 devices across 16 device categories. Another 12 devices were measured in detail in a laboratory setup. In all a total of 55 individual devices across 23 device categories were measured. Based on this measurement data we developed predictive software that showed that even with a single device in 23 device categories operating near maximum they would, in total, produce exposures at a distance of 1 m of 0.17% of the ICNIRP (2020) public exposure limits. Measurements were also made in two separate smart apartments—one contained over 50 IoT devices and a second with over 100 IoT devices with the devices driven as hard as could reasonably be achieved. The respective 6-min average exposure level recorded were 0.0077% and 0.44% of the ICNIRP (2020) 30-min average public exposure limit.

Excerpts

.... Based on all of these measurements, we have developed predictive software that can be used to estimate exposure levels on a conservative basis by incorporating a 3 dB enhancement to produce a realistic upper bound for the exposure estimation. The RF estimator tool, which is available from the Mobile & Wireless Forum (MWF) website (www.mwfai.org), has a drop-down menu that allows the selection of multiple IoT devices and separation distances and returns an estimate of the exposure level that could be expected in the home environment. Devices in neighboring rooms can be included in the software by selecting the appropriate device and distance. However, because of the inverse square dependence with distance and the attenuation through walls, the majority contribution in a particular room is determined by the proximity of devices in that room. Future work will include the expansion of the device categories and the inclusion of wall/window attenuation to account for neighboring homes and apartments.

CONFLICT OF INTEREST STATEMENT This work was fully funded by the Mobile and Wireless Forum (www.mwfai.org) and that KJ and MM are contractored by the MWF.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.22518

--

Review Antenna Design for Modern Mobile Phones: A Review

Wang Y, Sun L, Du Z, Zhang Z. Review Antenna Design for Modern Mobile Phones: A Review. Electromagnetic Science, 2(2):1-36. June 2024, doi: 10.23919/emsci.2023.0052.

Abstract

Due to limited antenna space, high communication requirements, and strict regulatory constraints, the design of antennas for modern mobile phones has become an exceedingly challenging task. In recent years, numerous studies have been conducted in this area, leading to significant advancements. This review paper comprehensively summarizes recent progress made in antenna design for modern mobile phones. Firstly, the challenges faced in antenna design for modern mobile phones are described, including bandwidth enhancement, integration and decoupling techniques, mm-wave array antennas,

satellite communication antennas, as well as interactions between mobile antennas and the human body. Secondly, the basic antenna types (such as inverted-F, slot, loop, and planar inverted-F antennas) commonly used in modern metal-bezel mobile phones along with their key characteristics are briefly summarized. Thirdly, the commonly employed methods used in practical applications for designing wideband antennas within compact sizes and achieving decoupling among multiple antennas with wide bandwidths are collected. Fourthly, recent advances in the design of compact, wideband, and wide-angle scanning mm-wave arrays for modern mobile phones are summarized. Fifthly, recent progress made in satellite communication antenna designs for modern mobile phones, including broadside and end-fire radiation patterns, is presented. Sixthly, recent studies on the interaction between mobile antennas and the human body are briefly concluded. Finally, the future challenge of antenna design for mobile phones is briefly discussed. It is our hope that this comprehensive review will provide readers with a systematic understanding of antenna design principles applicable to modern mobile phones.

Excerpts

"Currently, with the improvement of communication requirements and the evolution of communication specifications, the number of antennas in smartphones has reached 20–30 (as depicted in Figure 3), including the following antennas:

- Four antennas for LB (low band): 698–960 MHz;
- Four antennas for MHB (mid- and high-band) MIMO operation: 1710–2690 MHz;
- Four antennas for 5G New Radio (NR) band MIMO operation: 3300–4200 MHz & 4400–5000 MHz;
- Two or more dual-band millimeter-wave AiP antennas: 24.25-29.50 GHz and 37.00-43.50 GHz;
- One or more satellite communication antennas: the operating band is the L- or S-band;
- Two GNSS (global navigation satellite system) antennas: GPS L5 at 1176 MHz and GPS L1 at 1575 MHz;
- Two to four tri-band WiFi (wireless fidelity) and BT (Bluetooth) antennas: 2400–2484 MHz, 5150–5350 MHz, and 5725–5825 MHz;
- One NFC (near field communication) antenna: 13.56 MHz;
- Three UWB (ultra-wideband) antennas: 6240–6740 MHz and 7750–8250 MHz."

"In [199], the interaction principle of antenna radiation fields with nearby human bodies are analyzed by using electromagnetic boundary conditions. Under these boundary conditions, the horizontal E-field penetrates human tissue more easily than the vertical E-field does [199], as shown in Figure 49. Thus, mobile antennas with a vertical E-field have a higher on-body efficiency and lower SAR. As shown in Figure 50, a smartwatch antenna with a vertical E-field has a weak E-field in the human body, thus, the performance of an antenna with a vertical E-field is approximately 4 dB greater than that of an antenna with a horizontal E-field [200], [201]. Additionally, the vertical E-field method can also be applied to ear-bar TWS earphones, which can improve the antenna performance from 19.1% to 30.9% with a vertical E-field [202]."

Open access paper: https://ieeexplore.ieee.org/document/10636175

--

High Frequency Electromagnetic Field Exposure in Paediatric and Female Patients with Implanted Cardiac Pacemaker

Bacova F, Benova M, Psenakova Z, Smetana M, Pacek M, Ochodnicky J. High Frequency Electromagnetic Field Exposure in Paediatric and Female Patients with Implanted Cardiac Pacemaker. *Applied Sciences*. 2024; 14(16):7198. doi: 10.3390/app14167198.

Abstract

This article investigates the effects of electromagnetic field (EMF) from mobile phones on human tissues and implanted medical devices. The intensity of the electric field (E) is evaluated based on simulations and measurements of various exposure scenarios. An area of interest is the case of a person with an implanted device (heart pacemaker) who may be affected by this exposure. Due to the rapid development of communication technologies and the growing awareness of the potential health risks of radio frequency (RF) EMF, the International Commission on Non-Ionizing Radiation Protection (ICNIRP) has established exposure limits within the European Union. Our study models and analyses EMF values in human tissues in an ideal environment, in a situation where a person uses a mobile phone in the DCS (Digital Cellular System) band, including the case of a person with an implanted pacemaker. Pilot simulations were verified by experimental measurements. Based on them, specific human models with the best matching results were selected for modelling other possible interactions of exogenous EMF and cardiac pacemaker in the same situations and locations

Excerpt

Based on our simulations and experimental measurements, we found that maintaining a minimum distance of 15 cm between the mobile phone and the pacemaker is not sufficient to ensure patient safety. Our results show that even at a distance of 30 cm, the E values for the models were: MM—54.96 V/m, for Child—65.34 V/m, for Laura—62.81 V/m, and measurements—52.72 In M. In the case of the Child and Laura complex models, the E values still exceeded the exposure limit of 58 V/m for a frequency of 1800 MHz. Our recommendation is to push this limit up to more than 30 cm.

Another recommendation is protective equipment that reduces the exposure of parts of the human body to EMF. However, these aids must have a certificate of satisfactory EM compatibility.

Open access paper: https://www.mdpi.com/2076-3417/14/16/7198

--

Importance of magnetic information for neuronal plasticity in desert ants

Grob R, Müller VL, Grübel K, Fleischmann PN. Importance of magnetic information for neuronal plasticity in desert ants. PNAS. 2024. 121 (8) e2320764121. doi: pnas.2320764121.

Significance

The Earth's magnetic field is an essential navigational cue for many animal species. However, where in the brain magnetic information is processed is still little understood. In this paper, we analyzed structural neuronal plasticity in the brain of Cataglyphis desert ants following sky-compass calibration under permanently manipulated magnetic field conditions. Our results demonstrate that information from the Earth's magnetic field is integrated into the ants' internal compass (central complex) and into

the learning and memory centers (mushroom bodies). Together with our behavioral analyses, the results show that the ants use magnetic information both as a navigational compass and as a reference system for visual compass calibration.

Abstract

Many animal species rely on the Earth's magnetic field during navigation, but where in the brain magnetic information is processed is still unknown. To unravel this, we manipulated the natural magnetic field at the nest entrance of Cataglyphis desert ants and investigated how this affects relevant brain regions during early compass calibration. We found that manipulating the Earth's magnetic field has profound effects on neuronal plasticity in two sensory integration centers. Magnetic field manipulations interfere with a typical look-back behavior during learning walks of naive ants. Most importantly, structural analyses in the ants' neuronal compass (central complex) and memory centers (mushroom bodies) demonstrate that magnetic information affects neuronal plasticity during early visual learning. This suggests that magnetic information does not only serve as a compass cue for navigation but also as a global reference system crucial for spatial memory formation. We propose a neural circuit for integration of magnetic information into visual guidance networks in the ant brain. Taken together, our results provide an insight into the neural substrate for magnetic navigation in insects.

https://www.pnas.org/doi/10.1073/pnas.2320764121

--

A critical appraisal of the WHO 2024 systematic review of the effects of RF-EMF exposure on tinnitus, migraine/headache, and non-specific symptoms (WHO SR 7)

Frank J, Melnick R, Moskowitz J, on behalf of the International Commission on the Biological Effects of Electromagnetic Fields (ICBE-EMF). A critical appraisal of the WHO 2024 systematic review of the effects of RF-EMF exposure on tinnitus, migraine/headache, and non-specific symptoms. Reviews on Environmental Health. 2024. doi: 10.1515/reveh-2024-0069.

Abstract

The World Health Organization (WHO) in 2012 initiated an expert consultation about research on the health effects of radio-frequency electromagnetic fields (RF-EMF) for a WHO monograph that was last updated in 1993. The project was abandoned over concerns about the quality of the commissioned review papers. The WHO restarted the project in 2019 by commissioning ten systematic reviews (SRs) of the research on RF-EMF exposure and adverse biological and health outcomes in laboratory animals, cell cultures, and human populations. The second of these SRs, published in 2024, addresses human observational studies of RF-EMF exposure and non-specific symptoms, including tinnitus, migraine/headache, and sleep disturbance. The present commentary is a critical appraisal of the scientific quality of this SR (SR7) employing criteria developed by the Oxford Centre for Evidence-Based Medicine. Based upon our review, we call for a retraction of SR7 and an impartial investigation by unconflicted experts of the currently available evidence and future research priorities.

Conclusion

To summarize, the way in which any epidemiologically unsophisticated reader is likely to be misled by this SR is clear. It appears to conclude unequivocally that the body of scientific evidence reviewed supports the safety of current (e.g. ICNIRP-based) population exposure limits for RF-EMF [10]. We reiterate that, on the contrary, this body of evidence is not adequate to either support or refute the safety of current exposure limits – largely due to the very small number and low methodological quality of the relevant primary studies to date, and the fundamental inappropriateness of meta-analysis for the handful of very heterogeneous primary studies identified by Röösli et al. [3] for each of the exposure/outcome combinations analysed.

We therefore call for a retraction of the SR by Röösli et al., and an impartial international investigation, by unconflicted experts, of both the currently available evidence base on these issues, as well as related research priorities for the future. That investigation should particularly address, above and beyond the topic of priority health outcomes to be researched (which was already assessed in the international expert consultation by WHO in 2018) [2], and the need for improved methods of accurately measuring RF-EMF exposures, suitable for large human observational studies in the general population – the Achilles heel of the current literature.

Open access paper: https://www.degruyter.com/document/doi/10.1515/reveh-2024-0069/html

--

WHO to build neglect of RF-EMF exposure hazards on flawed EHC reviews? Case study demonstrates how "no hazards" conclusion is drawn from data showing hazards (WHO SR 3)

Nordhagen E, Flydal E. WHO to build neglect of RF-EMF exposure hazards on flawed EHC reviews? Case study demonstrates how "no hazards" conclusion is drawn from data showing hazards. *Reviews on Environmental Health*. 2024. doi: 10.1515/reveh-2024-0089.

Abstract

We examined one of the first published of the several systematic reviews being part of WHO's renewed initiative to assess the evidence of associations between man-made radiofrequency electromagnetic radiation (RF-EMF) and adverse health effects in humans. The examined review addresses experimental studies of pregnancy and birth outcomes in non-human mammals. The review claims that the analyzed data did not provide conclusions certain enough to inform decisions at a regulatory level. Our objective was to assess the quality of this systematic review and evaluate the relevance of its conclusions to pregnant women and their offspring. The quality and relevance were checked on the review's own premises: e.g., we did not question the selection of papers, nor the chosen statistical methods. While the WHO systematic review presents itself as thorough, scientific, and relevant to human health, we identified numerous issues rendering the WHO review irrelevant and severely flawed. All flaws found skew the results in support of the review's conclusion that there is no conclusive evidence for nonthermal effects. We show that the underlying data, when relevant studies are cited correctly, support the opposite conclusion: There are clear indications of detrimental nonthermal effects from RF-EMF exposure. The many identified flaws uncover a pattern of systematic skewedness aiming for uncertainty hidden behind complex scientific rigor. The skewed methodology and low quality of this review is highly concerning, as it threatens to undermine the trustworthiness and professionalism of the WHO in the area of human health hazards from man-made RF-EMF.

Conclusions

The rigorous protocol and extensive analyses presented in EHC2023 and its protocol, convey an impression of serious science, credibility, and reliability. However, we have shown that this is not the case.

We found EHC2023 to be a massive work with a rigorous and complex protocol and extensive and complex statistical analyses. A consequence of the complexity is that it can be assumed that no average reader – not even professionals – will check the results of the review, if not for other reasons, because of the major effort needed. Thereby, scientific exchange, debate and control is impeded and reduced to a matter of trust.

We had the opportunity to spend time on an in-depth analysis of representative parts of EHC2023 to assess its quality based to the extent possible on the review's own premises – that is, independent of our opinion about the professional premises chosen.

We cannot prove that the flaws and omissions are deliberately added to reach wanted conclusions, as we have next to no information about the authors, neither of the process behind the authoring EHC2023 or its protocol. Anyhow, and whatever the cause, the EHC2023 review is clearly of such a low quality, also when evaluated within the thermal only tradition, that its conclusions are without scientific value.

Our findings show that the conclusion of EHC2023 is not well-founded, and therefore the final conclusions of EHC2023 that no conclusion can be drawn that are (EHC2023, p. 31) "certain enough to inform decisions at a regulatory level" cannot be trusted. The errors, flaws and omissions are grave enough to render EHC2023 unscientific and unethical, and it should therefore be retracted.

As it now stands, the conclusion of ECH2023 stands out as what appears to be a manufactured argument for current regulations being adequate to protect the health of human mothers and their offspring. Manipulating and skewing research results in order to manufacture a wanted conclusion is a well-known strategy to avoid stricter regulations [11], [12], [13], [14], [15]. Further investigations and better sources would be needed to prove such an assault on humanity to be the case as to EHC2023.

EHC2023 is just one of several studies commissioned by the same organization (WHO EHC no. 137) and states clearly that consistency has been assured in the protocols for these studies. Our analysis of EHC2023 may in this view be seen as a case study of the results of the entire WHO EHC undertaking: Since many of our concerns are related to core elements of the protocol, there are good reasons also to question the quality of all present and forthcoming results being part of the WHO EHC undertaking unless a thorough revision of its course is made.

Open access paper: https://www.degruyter.com/document/doi/10.1515/reveh-2024-0089/html

--

A comprehensive review of 5G NR RF-EMF exposure assessment technologies: fundamentals, advancements, challenges, niches, and implications

Korkmaz E, Aerts S, Coesoij R, Bhatt CR, Velghe M, Colussi L, Land D, Petroulakis N, Spirito M, Bolte J. A comprehensive review of 5G NR RF-EMF exposure assessment technologies: fundamentals, advancements, challenges, niches, and implications. Environ Res. 2024 Jul 6;260:119524. doi: 10.1016/j.envres.2024.119524.

Highlights

- Monitoring exposure to radiofrequency electromagnetic fields (RF-EMF) is crucial for environmental health and risk assessment
- A comprehensive review of the diverse landscape of RF-EMF assessment tools was missing.
- There is a definite need for cost-effective and long-lasting EMF sensors.
- Custom-developed RF-EMF measurement tools lack a standardized framework for comparison and validation.

Abstract

This review offers a detailed examination of the current landscape of radio frequency (RF) electromagnetic field (EMF) assessment tools, ranging from spectrum analyzers and broadband field meters to area monitors and custom-built devices. The discussion encompasses both standardized and non-standardized measurement protocols, shedding light on the various methods employed in this domain. Furthermore, the review highlights the prevalent use of mobile apps for characterizing 5G NR radio network data. A growing need for low-cost measurement devices is observed, commonly referred to as “sensors” or “sensor nodes”, that are capable of enduring diverse environmental conditions. These sensors play a crucial role in both microenvironmental surveys and individual exposures, enabling stationary, mobile, and personal exposure assessments based on body-worn sensors, across wider geographical areas. This review revealed a notable need for cost-effective and long-lasting sensors, whether for individual exposure assessments, mobile (vehicle-integrated) measurements, or incorporation into distributed sensor networks. However, there is a lack of comprehensive information on existing custom-developed RF-EMF measurement tools, especially in terms of measuring uncertainty. Additionally, there is a need for real-time, fast-sampling solutions to understand the highly irregular temporal variations EMF distribution in next-generation networks. Given the diversity of tools and methods, a comprehensive comparison is crucial to determine the necessary statistical tools for aggregating the available measurement data.

Conclusions

The objective of this review was to establish a groundwork for progress in the field of RF-EMF exposure assessment, ultimately contributing to a more thorough and efficient assessment. This review provides a comprehensive overview of the current state-of-the-art concerning RF-EMF measuring instruments. It covers a wide array of tools, such as spectrum analyzers, broadband field meters, area monitors, personal exposimeters, and custom-built instruments, as well as the existing measurement protocols, encompassing both standardized and non-standardized methods. In addition, we also have presented some of the most commonly used mobile apps for collecting 5G NR radio network data, which have also been used in RF-EMF exposure assessments. However, it is not yet clear on how accurate the measurement results of these apps are and how they compare among themselves and to more sophisticated tools.

Most importantly, this review revealed the need for cost-effective and long-lasting measurement devices or sensors that are capable of collecting data at a high time resolution in various frequency bands, as well as withstanding various environmental conditions. These sensors are essential for conducting stationary, mobile, and personal exposure assessments across larger geographical areas, time intervals, and populations than current capabilities allow. Additionally, it is important to recognize that the specific requirements for these sensors differ based on their intended usage, e.g., on-body measurement devices need to take into account the influence of the body, vehicle-integrated sensors the influence of the speed and the relative position of the sensor on the vehicle, and sensors on infrastructure the influence of the height and the building materials. Furthermore, there exists a demand for real-time, fast-sampling solutions to comprehend the highly irregular temporal variations in EMF distribution within next-generation networks.

Moreover, there is a notable absence of extensive information regarding currently employed custom-developed RF-EMF measurement tools, particularly with respect to measuring uncertainty. Considering the diversity of tools and methodologies in use, conducting a thorough comparison becomes crucial to identify the necessary statistical tools for aggregating the available measurement data.

A more in-depth discussion relating the current 5G NR assessment methods to measurement equipment is intended for a follow-up study, which will describe more in detail the requirements, opportunities, and priorities for new, low-cost, custom-built measurement equipment.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0013935124014294

--

The effect of exposure to radiofrequency electromagnetic fields on cognitive performance in human experimental studies: Systematic review and meta-analyses (SR 6)

Pophof B, Kuhne J, Schmid G, Weiser E, Dorn H, Henschenmacher B, Burns J, Danker-Hopfe H, Sauter C, Pophof B. The effect of exposure to radiofrequency electromagnetic fields on cognitive performance in human experimental studies: Systematic review and meta-analyses. Environment International (2024). doi: 10.1016/j.envint.2024.108899.

Abstract

Background The objective of this review is to evaluate the associations between short-term exposure to radiofrequency electromagnetic fields (RF-EMF) and cognitive performance in human experimental studies.

Methods Online databases (PubMed, Embase, Scopus, Web of Science and EMF-Portal) were searched for studies that evaluated effects of exposure to RF-EMF on seven domains of cognitive performance in human experimental studies. The assessment of study quality was based on the Risk of Bias (RoB) tool developed by the Office of Health Assessment and Translation (OHAT). Random effects meta-analyses of Hedges's g were conducted separately for accuracy- and speed-related performance measures of various cognitive domains, for which data from at least two studies were available. Finally, the certainty of evidence for each identified outcome was assessed according to Grading of Recommendations Assessment, Development, and Evaluation (GRADE).

Results 57,543 records were identified and 76 studies (80 reports) met the inclusion criteria. The included 76 studies with 3846 participants, consisting of humans of different age, sex and health status from 19 countries, were conducted between 1989 and 2021. Quantitative data from 50 studies (52 reports) with 2433 participants were included into the meta-analyses. These studies were performed in 15 countries between 2001 and 2021. The majority of the included studies used head exposure with GSM 900 uplink.

None of the meta-analyses observed a statistically significant effect of RF-EMF exposure compared to sham on cognitive performance as measured by the confidence interval surrounding the Hedges's g or the significance of the z-statistic.

For the domain Orientation and Attention, subclass Attention – Attentional Capacity RF-EMF exposure results in little to no difference in accuracy (Hedges's g 0.024, 95 % CI [-0.10; 0.15], I2 = 28 %, 473 participants).

For the domain Orientation and Attention, subclass Attention – Concentration / Focused Attention RF-EMF exposure results in little to no difference in speed (Hedges's g 0.005, 95 % CI [-0.17; 0.18], I2 = 7 %, 132 participants) and probably results in little to no difference in accuracy; it does not reduce accuracy (Hedges's g 0.097, 95 % CI [-0.05; 0.24], I2 = 0 %, 217 participants).

For the domain Orientation and Attention, subclass Attention – Vigilance RF-EMF exposure probably results in little to no difference in speed and does not reduce speed (Hedges's g 0.118, 95 % CI [-0.04; 0.28], I2 = 41 %, 247 participants) and results in little to no difference in accuracy (Hedges's g 0.042, 95 % CI, [-0.09; 0.18], I2 = 0 %, 199 participants).

For the domain Orientation and Attention, subclass Attention – Selective Attention RF-EMF exposure probably results in little to no difference in speed and does not reduce speed (Hedges's g 0.080, 95 % CI [-0.09; 0.25], I2 = 63 %, 452 participants); it may result in little to no difference in accuracy, but it probably does not reduce accuracy (Hedges's g 0.178, 95 % CI [-0.02; 0.38], I2 = 68 %, 480 participants).

For the domain Orientation and Attention, subclass Attention – Divided Attention RF-EMF exposure results in little to no difference in speed (Hedges's g −0.010, 95 % CI [-0.14; 0.12], I2 = 5 %, 307 participants) and may result in little to no difference in accuracy (Hedges's g −0.089, 95 % CI [-0.35; 0.18], I2 = 53 %, 167 participants).

For the domain Orientation and Attention, subclass Processing Speed − Simple Reaction Time Task RF-EMF exposure results in little to no difference in speed (Hedges's g 0.069, 95 % CI [-0.02; +0.16], I2 = 29 %, 820 participants).

For the domain Orientation and Attention, subclass Processing Speed – 2-Choice Reaction Time Task RF-EMF exposure results in little to no difference in speed (Hedges's g −0.023, 95 % CI [-0.13; 0.08], I2 = 0 %, 401 participants), and may result in little to no difference in accuracy (Hedges's g −0.063, 95 % CI [-0.38; 0.25], I2 = 63 %, 117 participants).

For the domain Orientation and Attention, subclass Processing Speed – >2-Choice Reaction Time Task RF-EMF exposure results in little to no difference in speed (Hedges's g −0.054, 95 % CI [-0.14; 0.03],

I2 = 0 %, 544 participants) and probably results in little to no difference in accuracy (Hedges's g −0.129, 95 % CI [-0.30; 0.04], I2 = 0 %, 131 participants).

For the domain Orientation and Attention, subclass Processing Speed – Other Tasks RF-EMF exposure probably results in little to no difference in speed and does not reduce speed (Hedges's g 0.067, 95 % CI [-0.12; 0.26], I2 = 38 %, 249 participants); it results in little to no difference in accuracy (Hedges's g 0.036, 95 % CI [-0.08; 0.15], I2 = 0 %, 354 participants).

For the domain Orientation and Attention, subclass Working Memory – n-back Task (0–3-back) we found Hedges's g ranging from −0.090, 95 % CI [-0.18; 0.01] to 0.060, 95 % CI [-0.06; 0.18], all I2 = 0 %, 237 to 474 participants, and conclude that RF-EMF exposure results in little to no difference in both speed and accuracy.

For the domain Orientation and Attention, subclass Working Memory – Mental Tracking RF-EMF exposure results in little to no difference in accuracy (Hedges's g −0.047, 95 % [CI −0.15; 0.05], I2 = 0 %, 438 participants). For the domain Perception, subclass Visual and Auditory Perception RF-EMF exposure may result in little to no difference in speed (Hedges's g −0.015, 95 % CI [-0.23; 0.195], I2 = 0 %, 84 participants) and probably results in little to no difference in accuracy (Hedges's g 0.035, 95 % CI [-0.13; 0.199], I2 = 0 %, 137 participants).

For the domain Memory, subclass Verbal and Visual Memory RF-EMF exposure probably results in little to no difference in speed and does not reduce speed (Hedges's g −0.042, 95 % CI [-0.15; 0.21], I2 = 0 %, 102 participants); it may result in little to no difference in accuracy (Hedges's g −0.087, 95 % CI [-0.38; 0.20], I2 = 85 %, 625 participants).

For the domain Verbal Functions and Language Skills, subclass Verbal Expression, a meta-analysis was not possible because one of the two included studies did not provide numerical values. Results of both studies did not indicate statistically significant effects of RF-EMF exposure on both speed and accuracy.

For the domain Construction and Motor Performance, subclass Motor Skills RF-EMF exposure may reduce speed, but the evidence is very uncertain (Hedges's g −0.919, 95 % CI [-3.09; 1.26], I2 = 96 %, 42 participants); it probably results in little to no difference in accuracy and does not reduce accuracy (Hedges's g 0.228, 95 % CI [-0.01; 0.46], I2 = 0 %, 109 participants).

For the domain Concept Formation and Reasoning, subclass Reasoning RF-EMF exposure results in little to no difference in speed (Hedges's g 0.010, 95 % CI [-0.11; 0.13], I2 = 0 %, 263 participants) and probably results in little to no difference in accuracy and does not reduce accuracy (Hedges's g 0.051, 95 % CI [-0.14; 0.25], I2 = 0 %, 100 participants).

For the domain Concept Formation and Reasoning, subclass Mathematical Procedures RF-EMF exposure results in little to no difference in speed (Hedges's g 0.033, 95 % CI [-0.12; 0.18], I2 = 0 %, 168 participants) and may result in little to no difference in accuracy but probably does not reduce accuracy (Hedges's g 0.232, 95 % CI [-0.12; +0.59], I2 = 86 %, 253 participants).

For the domain Executive Functions there were no studies.

Discussion Overall, the results from all domains and subclasses across their speed- and accuracy-related outcome measures according to GRADE provide high to low certainty of evidence that short-term RF-EMF exposure does not reduce cognitive performance in human experimental studies. For 16 out of 35

subdomains some uncertainty remains, because of limitations in the study quality, inconsistency in the results or imprecision of the combined effect size estimate. Future research should focus on construction and motor performance, elderly, and consideration of both sexes.

Other This review was partially funded by the WHO radioprotection programme. The protocol for this review was registered in Prospero reg. no. CRD42021236168 and published in Environment International (Pophof et al. 2021).

4.9. Implications of practice and policy

The present systematic review provides mostly moderate to high certainty of evidence that short-term RF-EMF exposure at SAR levels within the recommended limits (ICNIRP 2010) does not negatively affect the investigated domains of cognitive function.

4.10. Implications for research

Although the number of studies and participants is low for several domains / subclasses, the certainty of evidence for lack of an effect is very low only for the speed category of D5 *Construction and Motor Performance*. Only two small studies with a total number of 42 participants contribute to the result with substantial heterogeneity and the body of evidence suggests a large negative effect with very low certainty. Further studies are needed to confirm or refute this effect. In particular, all outcome measures used in both studies should be considered and reported numerically.

The number of studies that provide age-specific information on RF-EMF effects on cognitive performance is small. In particular, only five of the included studies were performed in children and only one in the elderly. Since children are generally considered to be a sensitive group, and the cognitive performance tends to decline with age, studies in children /adolescents and in elderly subjects are recommended to investigate whether these both age groups are more affected by RF-EMF exposure than young adults. Since the subgroup analyses (Supplementary Data 11) provide weak indications for a possible effect of sex, analyses stratified for sex of the participants are recommended.

In general, any new study should be designed in a way that any potential RoB is minimized. The criteria for definitely low RoB are summarized in Supplementary Data 7 of the protocol (Pophof et al. 2021). Furthermore, power calculations should be conducted prior to the final study design and study power should be sufficient for detecting even small effects. A comprehensive list of further issues that should be considered when conducting studies on possible effects of RF-EMF on cognitive performance, are published in Regel and Achermann (2011).

Authors of future studies are encouraged to make the original individual data publicly available in appropriate data repositories to avoid missing data due to incomplete reporting.

5.2. Deviations from the protocol

In the protocol, we did not describe how the ratings of the individual RoB questions for a given study would be used to define an overall quality rating for that study. This admission could be considered as a potential risk of bias for this systematic review if, for example, certain studies were selectively included/excluded from subgroup analyses based on the quality rating. However, given that the heterogeneity of effects across studies was low for most outcome measures, such selective analysis

would in principle have little to no impact. The GRADE assessment, however, strongly depends on the overall study quality assessment. Thus, in order to be as transparent as possible, we have explained and justified the chosen method for allocating studies to quality tiers in detail in Methods − Risk of Bias assessment.

We did not conduct separate analyses for individuals with and without IEI-EMF, because there were only seven studies including subjects with IEI. Only six of the seven studies provided numerical data (for only some domains / subclasses) to be used to calculate effect sizes. The data of these six studies do not allow a detailed analysis following the approach pursued in this paper.

Because there was only one study investigating females, heterogeneity could not be assessed for this subgroup. Therefore, we performed subgroup analyses stratified by studies investigating only males, and studies investigating males and females together. However, the explanatory power of this approach should not be overestimated.

We did not evaluate a potential dose–response relationship or perform subgroup analyses regarding exposure levels, because independently from the exposure level there was predominantly no effect of RF-EMF, and the exposure assessment in most studies did not provide spatially resolved exposure levels in the brain. For similar reasons we did not analyse the exposure category D separately from A, B and C (see Protocol, Pophof et al. 2021), but combined all exposure categories in analyses.

Due to the huge amount of data available for the domains D1 and to a lesser extent for D6, and due to the fact that some studies reported results for a large number of different subclasses of domains, a problem with the unit of analysis would have resulted in a huge amount of original data that could not be used for a *meta*-analysis if we had kept the outcome categories defined in the protocol. Therefore, we defined more detailed outcome categories according to the domain subclasses (Table 2) as defined by Lezak et al. (2012, Part II The Compendium of Tests and Assessment Techniques, p. 391 ff.). This enables a better differentiation of different cognitive abilities, and increases the number of possible *meta*-analyses and a higher proportion of original data included in the *meta*-analyses.

In order to explore possible sources of heterogeneity, we additionally included a stratification according to sample sizes (<30; ≥ 30). Studies with small sample sizes are likely to have a higher variability than studies with larger sample sizes.

Deviating from the protocol, we did not conduct subgroup analyses for the exposure-related categories of signals with different modulation, time course of exposure, different frequencies, for the nine outcome measures of subclasses with results with at least moderate heterogeneity (Table 3). This is because a large number of studies falls in the category “continuous GSM-modulated near-field exposure with frequency lower than 1 GHz” (in six out of these nine outcome measures at least half of results include this exposure category) and most of the outlier studies that strongly contribute to heterogeneity of the results also fall into this category (five studies).

5.3. Support

This review was partly funded by the World Health Organization (contracts 2020/1028371–0, 2022/1276784–0). WHO provided the basis for the protocol and methodological support throughout the review process.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: [The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. BP is member of the ICNIRP Scientific Expert Group (SEG) on environment, BfS observer in the working group SSK-A630 of the German Commission on Radiological Protection and was German delegate of European Cost Actions BM0704 and BM1309 “EMF-MED”. GS is member of the Committee “Non-Ionizing Radiation” (SSK-A6) and member of the working group SSK-A630 of the German Commission on Radiological Protection. GS is chair of the Austrian Standardization Sub-Committee TSK-EMV-EMF “Electromagnetic Fields”. HDHs research is entirely funded by public or not-for-profit foundations. She has served as advisor to a number of national and international public advisory groups concerning the potential health effects of exposure to non-ionizing radiation, including the World Health Organization, the German Commission on Radiological Protection (member of the committee “Non-Ionizing Radiation” (SSK-A6) and member of the working group 5G (SSK-A630)) and the Independent Expert Group of the Swedish Radiation Safety Authority. JK is member of the ICNIRP Scientific Expert Group (SEG) on ultrasound and BfS observer in the working group SSK-A630 of the German Commission on Radiological Protection].

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0160412024004859?via%3Dihub

--

Literature review: potential non-thermal molecular effects of external radiofrequency electromagnetic fields on cancer

Dieper A, Scheidegger S, Füchslin RM, Veltsista PD, Stein U, Weyland M, Gerster D, Beck M, Bengtsson O, Zips D, Ghadjar P. Literature review: potential non-thermal molecular effects of external radiofrequency electromagnetic fields on cancer. Int J Hyperthermia. 2024;41(1):2379992. doi: 10.1080/02656736.2024.2379992.

Abstract

Introduction: There is an ongoing scientific discussion, that anti-cancer effects induced by radiofrequency (RF)-hyperthermia might not be solely attributable to subsequent temperature elevations at the tumor site but also to non-temperature-induced effects. The exact molecular mechanisms behind said potential non-thermal RF effects remain largely elusive, however, limiting their therapeutical targetability.

Objective: Therefore, we aim to provide an overview of the current literature on potential non-temperature-induced molecular effects within cancer cells in response to RF-electromagnetic fields (RF-EMF).

Material and methods: This literature review was conducted following the PRISMA guidelines. For this purpose, a MeSH-term-defined literature search on MEDLINE (PubMed) and Scopus (Elsevier) was conducted on March 23rd, 2024. Essential criteria herein included the continuous wave RF-EMF nature

(3 kHz - 300 GHz) of the source, the securing of temperature-controlled circumstances within the trials, and the preclinical nature of the trials.

Results: Analysis of the data processed in this review suggests that RF-EMF radiation of various frequencies seems to be able to induce significant non-temperature-induced anti-cancer effects. These effects span from mitotic arrest and growth inhibition to cancer cell death in the form of autophagy and apoptosis and appear to be mostly exclusive to cancer cells. Several cellular mechanisms were identified through which RF-EMF radiation potentially imposes its anti-cancer effects. Among those, by reviewing the included publications, we identified RF-EMF-induced ion channel activation, altered gene expression, altered membrane potentials, membrane oscillations, and blebbing, as well as changes in cytoskeletal structure and cell morphology.

Conclusion: The existent literature points toward a yet untapped therapeutic potential of RF-EMF treatment, which might aid in damaging cancer cells through bio-electrical and electro-mechanical molecular mechanisms while minimizing adverse effects on healthy tissue cells. Further research is imperative to definitively confirm non-thermal EMF effects as well as to determine optimal cancer-type-specific RF-EMF frequencies, field intensities, and exposure intervals.

Open access paper: https://www.tandfonline.com/doi/full/10.1080/02656736.2024.2379992

--

Exposure to Electromagnetic Fields from Mobile Phones and Fructose Consumption Coalesce to Perturb Metabolic Regulators AMPK/SIRT1-UCP2/FOXO1 in Growing Rats

Tripathi R, Banerjee SK, Nirala JP, Mathur R. Exposure to Electromagnetic Fields from Mobile Phones and Fructose Consumption Coalesce to Perturb Metabolic Regulators AMPK/SIRT1-UCP2/FOXO1 in Growing Rats[J]. Biomedical and Environmental Sciences, 2023, 36(11): 1045-1058. doi: 10.3967/bes2023.134

Abstract

Objective In this study, the combined effect of two stressors, namely, electromagnetic fields (EMFs) from mobile phones and fructose consumption, on hypothalamic and hepatic master metabolic regulators of the AMPK/SIRT1-UCP2/FOXO1 pathway were elucidated to delineate the underlying molecular mechanisms of insulin resistance.

Methods Weaned Wistar rats (28 days old) were divided into 4 groups: Normal, Exposure Only (ExpO), Fructose Only (FruO), and Exposure and Fructose (EF). Each group was provided standard laboratory chow ad libitum for 8 weeks. Additionally, the control groups, namely, the Normal and FruO groups, had unrestricted access to drinking water and fructose solution (15%), respectively. Furthermore, the respective treatment groups, namely, the ExpO and EF groups, received EMF exposure (1,760 MHz, 2 h/day x 8 weeks). In early adulthood, mitochondrial function, insulin receptor signaling, and oxidative stress signals in hypothalamic and hepatic tissues were assessed using western blotting and biochemical analysis.

Result In the hypothalamic tissue of EF, SIRT1, FOXO 1, p-PI3K, p-AKT, Complex III, UCP2, MnSOD, and catalase expressions and OXPHOS and GSH activities were significantly decreased (P < 0.05) compared

to the Normal, ExpO, and FruO groups. In hepatic tissue of EF, the p-AMPKα, SIRT1, FOXO1, IRS1, p-PI3K, Complex I, II, III, IV, V, UCP2, and MnSOD expressions and the activity of OXPHOS, SOD, catalase, and GSH were significantly reduced compared to the Normal group (P < 0.05).

Conclusion The findings suggest that the combination of EMF exposure and fructose consumption during childhood and adolescence in Wistar rats disrupts the closely interlinked and multi-regulated crosstalk of insulin receptor signals, mitochondrial OXPHOS, and the antioxidant defense system in the hypothalamus and liver.

Open access paper: https://www.besjournal.com/en/article/doi/10.3967/bes2023.134

--

Upper bound for broadband radiofrequency field disruption of magnetic compass orientation in night-migratory songbirds

Leberecht B, Wong SY, Satish B, Döge S, Hindman J, Venkatraman L, Apte S, Haase K, Musielak I, Dautaj G, Solov'yov IA, Winklhofer M, Mouritsen H, Hore PJ. Upper bound for broadband radiofrequency field disruption of magnetic compass orientation in night-migratory songbirds. Proc Natl Acad Sci U S A. 2023 Jul 11;120(28):e2301153120. doi: 10.1073/pnas.2301153120. Epub 2023 Jul 3. PMID: 37399422; PMCID: PMC10334787.

Abstract

Night-migratory songbirds have a light-dependent magnetic compass sense, the mechanism of which is thought to depend on the photochemical formation of radical pairs in cryptochrome (Cry) proteins located in the retina. The finding that weak radiofrequency (RF) electromagnetic fields can prevent birds from orienting in the Earth's magnetic field has been regarded as a diagnostic test for this mechanism and as a potential source of information on the identities of the radicals. The maximum frequency that could cause such disorientation has been predicted to lie between 120 and 220 MHz for a flavin-tryptophan radical pair in Cry. Here we show that the magnetic orientation capabilities of Eurasian blackcaps (*Sylvia atricapilla*) are not affected by RF noise in the frequency bands 140 to 150 MHz and 235 to 245 MHz. From a consideration of its internal magnetic interactions, we argue that RF field effects on a flavin-containing radical-pair sensor should be approximately independent of frequency up to 116 MHz and that birds' sensitivity to RF disorientation should fall by about two orders of magnitude when the frequency exceeds 116 MHz. Taken together with our earlier finding that 75 to 85 MHz RF fields disrupt the magnetic orientation of blackcaps, these results provide compelling evidence that the magnetic compass of migratory birds operates by a radical pair mechanism.

Open access paper: https://www.pnas.org/doi/abs/10.1073/pnas.2301153120

--

Impact of high (1950 MHz) and extremely low (50 Hz) frequency electromagnetic fields on DNA damage caused by occupationally relevant exposures in human derived cell lines

Worel N, Mišík M, Kundi M, Ferk F, Hutter HP, Nersesyan A, Wultsch G, Krupitza G, Knasmueller S. Impact of high (1950 MHz) and extremely low (50 Hz) frequency electromagnetic fields on DNA damage caused by occupationally relevant exposures in human derived cell lines. Toxicol In Vitro. 2024 Jul 16:105902. doi: 10.1016/j.tiv.2024.105902.

Highlights

- Mobile phone-specific (HF) and low-frequency (LF) electromagnetic fields (EMF) may cause cancer.
- We investigated the impact of these fields on DNA damage in human-derived cells.
- The cells were exposed to the EMF in combination with occupationally related carcinogens.
- We found no evidence for synergistic effects.
- In neural cells, we observed a significant reduction of DNA damage after exposure to the HF field.

Abstract

Epidemiological studies indicate that electromagnetic fields are associated with cancer in humans. Exposure to mobile phone specific high frequency fields (HF-EMF) may lead to increased glioma risks, while low frequency radiation (LF-EMF) is associated with childhood leukemia. We studied the impact of HF-EMF (1950 MHz, UMTS signal) on DNA stability in an astrocytoma cell line (1321N1), and the effect of LF-EMF (50 Hz) in human derived lymphoma (Jurkat) cells. To find out if these fields affect chemically induced DNA damage, co-exposure experiments were performed. The cells were exposed to HF-EMF or LF-EMF and treated simultaneously and sequentially with mutagens. The compounds cause DNA damage via different molecular mechanisms, i.e. pyrimidine dimers which are characteristic for UV light (4-nitroquinoline 1-oxide, 4NQO), bulky base adducts (benzo[a]pyrene diolepoxide, BPDE), DNA-DNA and DNA-protein cross links and oxidative damage ($NiCl_2$, CrO_3). DNA damage was measured in single cell gel electrophoresis (comet) assays. We found a moderate reduction of basal and 4NQO-induced DNA damage in the astrocytoma line, but no significant alterations of chemically induced DNA migration by the HF and LF fields under all other experimental series. The biological consequences of the moderate reduction remain unclear, but our findings indicate that acute mobile phone and power line specific EMF exposures do not enhance genotoxic effects caused by occupationally relevant chemical exposures.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0887233324001322

--

Estimates and measurements of radiofrequency exposures in smart-connected homes

Joyner K, Milligan M, Knipe P. Estimates and measurements of radiofrequency exposures in smart-connected homes. Bioelectromagnetics. 2024 Jul 18. doi: 10.1002/bem.22518.

Abstract

The aim of this research was to quantify the levels of radiofrequency electromagnetic energy (RF-EME) in a residential home/apartment equipped with a range of wireless devices, often referred to as internet

of things (IoT) devices or smart devices and subsequently develop a tool that could be useful for estimating the levels of RF-EME in a domestic environment. Over the course of 3 years measurements were performed in peoples' homes on a total of 43 devices across 16 device categories. Another 12 devices were measured in detail in a laboratory setup. In all a total of 55 individual devices across 23 device categories were measured. Based on this measurement data we developed predictive software that showed that even with a single device in 23 device categories operating near maximum they would, in total, produce exposures at a distance of 1 m of 0.17% of the ICNIRP (2020) public exposure limits. Measurements were also made in two separate smart apartments-one contained over 50 IoT devices and a second with over 100 IoT devices with the devices driven as hard as could reasonably be achieved. The respective 6-min average exposure level recorded were 0.0077% and 0.44% of the ICNIRP (2020) 30-min average public exposure limit.

Conclusions

We have presented measurements of Wi-Fi signals from 55 individual IoT devices across 23 device categories, which showed that the respective 6-min average exposure levels recorded ranged from 0.0077% to 0.44% of the ICNIRP (2020) 30-min average public exposure limit. These results were corroborated by measurements made in two separate smart apartments—one contained over 50 IoT devices and a second with over 100 IoT devices with the devices driven as hard as could reasonably be achieved. Based on all of these measurements, we have developed predictive software that can be used to estimate exposure levels on a conservative basis by incorporating a 3 dB enhancement to produce a realistic upper bound for the exposure estimation. The RF estimator tool, which is available from the Mobile & Wireless Forum (MWF) website (www.mwfai.org), has a drop-down menu that allows the selection of multiple IoT devices and separation distances and returns an estimate of the exposure level that could be expected in the home environment. Devices in neighboring rooms can be included in the software by selecting the appropriate device and distance. However, because of the inverse square dependence with distance and the attenuation through walls, the majority contribution in a particular room is determined by the proximity of devices in that room. Future work will include the expansion of the device categories and the inclusion of wall/window attenuation to account for neighboring homes and apartments.

CONFLICT OF INTEREST STATEMENT

This work was fully funded by the Mobile and Wireless Forum (www.mwfai.org) and that KJ and MM are contractored by the MWF.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.22518

--

Maternal linalool treatment protects against radiofrequency wave-induced deteriorations in adolescent rats: A behavioral and electrophysiological study

Azimzadeh M, Noorbakhshnia M. Maternal linalool treatment protects against radiofrequency wave-induced deteriorations in adolescent rats: A behavioral and electrophysiological study. Sci Rep. 2024 Jul 27;14(1):17257. doi: 10.1038/s41598-024-68103-5.

Abstract

Recent years, the rapid advancement of technology has raised concerns. We studied the effects of prenatal exposure to 900 MHz radiofrequency (RF) from mobile phones and the protective effects of linalool on learning and memory, and anxiety in adolescent male and female offspring rats. Pregnant rats were divided into four groups: control, wave, wave + linalool, and linalool. Rats received linalool (25mg/kg) by gavage for 21 days. Irradiation was conducted from day 0 to day 21 of pregnancy. Offsprings underwent behavioral and electrophysiological tests on days 50 and 60 after birth. Exposure to RF during pregnancy caused anxiety-like behavior in the EPM test and impairment of learning and memory in the Morris water maze and shuttle box tests. Electrophysiological properties and synaptic plasticity of the dorsal hippocampal CA3-CA1 synapse showed a decrease in fEPSP amplitude and slope. The trace element levels in both male and female offspring were consistent across all groups compared to their respective controls. In the hippocampus tissue, the levels of Fe, Cu, and Mn, as well as the Cu/Zn ratio, were significantly higher in the exposed groups (wave groups) compared to their controls. Moreover, Zn levels were significantly lower in the hippocampus tissue of the exposed groups. Linalool administration mitigated the excessive increase in Fe, Cu, Mn, and Cu/Zn ratio and normalized the disrupted levels of trace elements, except for Zn levels in both male and female offspring. Sex differences were observed in the EPM and shuttle box tests, females were more sensitive than males. In summary, our study demonstrates that prenatal exposure to mobile phone radiation induces stress-like behaviors, disrupts learning and memory, alters hippocampal electrophysiological properties and trace element balance in offspring. Treatment with linalool mitigates these deleterious effects, highlighting its potential as a therapeutic intervention. These findings contribute to our understanding of the impact of prenatal environmental exposures on neurodevelopment and offer insights into potential strategies for neuroprotection.

https://www.nature.com/articles/s41598-024-68103-5

--

Sulforaphane Effects on Neuronal-like Cells and Peripheral Blood Mononuclear Cells Exposed to 2.45 GHz Electromagnetic Radiation

Bertuccio MP, Saija C, Acri G, Ientile R, Caccamo D, Currò M. Sulforaphane Effects on Neuronal-like Cells and Peripheral Blood Mononuclear Cells Exposed to 2.45 GHz Electromagnetic Radiation. Int J Mol Sci. 2024 Jul 18;25(14):7872. doi: 10.3390/ijms25147872.

Abstract

Exposure to 2.45 GHz electromagnetic radiation (EMR) emitted from commonly used devices has been reported to induce oxidative stress in several experimental models. Our study aims to evaluate the efficacy of sulforaphane, a well-known natural product, in preventing radiation-induced toxic effects caused by a 24 h exposure of SH-SY5Y neuronal-like cells and peripheral blood mononuclear cells (PBMCs) to 2.45 GHz EMR. Cells were exposed to radiation for 24 h in the presence or absence of sulforaphane at different concentrations (5-10-25 µg/mL). Cell viability, mitochondrial activity alterations, the transcription and protein levels of redox markers, and apoptosis-related genes were investigated. Our data showed a reduction in cell viability of both neuronal-like cells and PBMCs caused by EMR exposure and a protective effect of 5 µg/mL sulforaphane. The lowest sulforaphane

concentration decreased ROS production and increased the Mitochondrial Transmembrane Potential ($\Delta\psi m$) and the NAD+/NADH ratio, which were altered by radiation exposure. Sulforaphane at higher concentrations displayed harmful effects. The hormetic behavior of sulforaphane was also evident after evaluating the expression of genes coding for Nrf2, SOD2, and changes in apoptosis markers. Our study underlined the vulnerability of neuronal-like cells to mitochondrial dysfunction and oxidative stress and the possibility of mitigating these effects by supplementation with sulforaphane. To our knowledge, there are no previous studies about the effects of SFN on these cells when exposed to 2.45 GHz electromagnetic radiation.

Open access paper: https://www.mdpi.com/1422-0067/25/14/7872

--

Ubiquitous extremely low frequency electromagnetic fields induces anxiety-like behavior: mechanistic perspectives

Hosseini E. Ubiquitous extremely low frequency electromagnetic fields induces anxiety-like behavior: mechanistic perspectives. Electromagn Biol Med. 2024 Jul 29:1-16. doi: 10.1080/15368378.2024.2380305.

Abstract

Anxiety is an adaptive condition characterized by heightened uneasiness, which in the long term can cause complications such as reducing the quality of life and problems related to the mental and physical health. Concerns have been raised regarding the potential dangers of extremely low frequency electromagnetic fields (ELF-EMF) ranging from 3 to 3000 Hz, which are omnipresent in our daily lives and there have been studies about the anxiogenic effects of these fields. Studies conducted in this specific area has revealed that ELF-EMF can have an impact on various brain regions, such as the hippocampus. In conclusion, studies have shown that ELF-EMF can interfere with hippocampus-prefrontal cortex pathway, inducing anxiety behavior. Also, ELF-EMF may initiate anxiety behavior by generating oxidative stress in hypothalamus and hippocampus. Moreover, ELF-EMF may induce anxiety behavior by reducing hippocampus neuroplasticity and increasing the $NMDA2_A$ receptor expression in the hippocampus. Furthermore, supplementation with antioxidants could serve as an effective protective measure against the adverse effects of ELF-EMF in relation to anxiety behavior.

Conclusion

In the present study, it was found out that:

1. ELF-EMF by creating oxidative stress in different regions of brain involved in anxiety behavior, results in an increase of stress hormones and also excitability subsequently.
2. ELF-EMF by decreasing in neuroplasticity and modulation of some receptors in the hippocampus may have a key role in initiating anxiety behavior.

https://pubmed.ncbi.nlm.nih.gov/39074042/

--

Characterizing parameters and incorporating action potentials via the Hodgkin-Huxley model in a novel electric model for living cells

Bougandoura O, Achour Y, Zaoui A, Starzyński J. Characterizing parameters and incorporating action potentials via the Hodgkin-Huxley model in a novel electric model for living cells. Electromagn Biol Med. 2024 Jul 2;43(3):187-203. doi: 10.1080/15368378.2024.2372107.

Abstract

To enhance our understanding of electroporation and optimize the pulses used within the frequency range of 1 kHz to 100 MHz, with the aim of minimizing side effects such as muscle contraction, we introduce a novel electrical model, structured as a 2D representation employing exclusively lumped elements. This model adeptly encapsulates the intricate dynamics of living cells' impedance variation. A distinguishing attribute of the proposed model lies in its capacity to decipher the distribution of transmembrane potential across various orientations within living cells. This aspect bears critical importance, particularly in contexts such as electroporation and cellular stimulation, where precise knowledge of potential gradients is pivotal. Furthermore, the augmentation of the proposed electrical model with the Hodgkin-Huxley (HH) model introduces an additional dimension. This integration augments the model's capabilities, specifically enabling the exploration of muscle cell stimulation and the generation of action potentials. This broader scope enhances the model's utility, facilitating comprehensive investigations into intricate cellular behaviors under the influence of external electric fields.

https://pubmed.ncbi.nlm.nih.gov/38990565/

--

Risk Assessment for Workers with Wearable Medical Devices Exposed to Electromagnetic Fields

Vivarelli C, Censi F, Calcagnini G, Falsaperla R, Mattei E. Risk Assessment for Workers with Wearable Medical Devices Exposed to Electromagnetic Fields. Health Physics 127(2):p 269-275, August 2024. doi: 10.1097/HP.0000000000001798.

Abstract

The exponential diffusion of wearable medical devices (WMD) in recent years has involved people of all ages, including workers. Workers who use WMDs should be considered at a particular risk from electromagnetic fields, and in accordance with EU Directive 2013/35/EU, they require an individual risk assessment. Currently, there is no international standard that provides specific guidance on how to perform such a risk assessment. This paper focuses on the effects of electromagnetic fields on WMDs and does not consider the direct effects on human body tissues. It aims to offer practical recommendations to employers and/or health physicists for the risk assessment of workers with WMDs. Focusing on EU countries, we first describe the requirements outlined by the technical standard for the electromagnetic compatibility (EMC) of medical electrical equipment EN 60601-1-2. Then, some general guidelines on how to perform the risk assessment are provided. The assessment can be conducted by comparing the field values measured in the workplace with the immunity test levels specified in the technical standards of medical electrical equipment. If the measured values are lower than the immunity test levels indicated in the standard and the distance from the electromagnetic source is

greater than the distance used by the manufacturer during the EMC (electromagnetic compatibility) tests (typically 30 cm), the risk for the worker may be considered acceptable. However, if the measured values exceed the immunity test levels or the distance criteria, a specific evaluation based on a case-by-case analysis is required.

https://journals.lww.com/health-physics/abstract/2024/08000/risk_assessment_for_workers_with_wearable_medical.1.aspx

--

Health risks for medical personnel due to magnetic fields in magnetic resonance imaging

König AM, Pöschke A, Mahnken AH. Health risks for medical personnel due to magnetic fields in magnetic resonance imaging. Rofo. 2024 Jul 19. English, German. doi: 10.1055/a-2296-3860.

Abstract

The current state of medical and scientific knowledge on the effects of exposure to electromagnetic fields on workers in the field of clinical magnetic resonance imaging (MRI) is summarized here.A systematic literature search was conducted to analyze the health risks to medical personnel from magnetic fields in MRI. A total of 7273 sources were identified, with 7139 being excluded after screening of the title and abstract. After full-text screening, 34 sources remained and were included in this paper. There are a number of scientific publications on the occurrence of short-term sensory effects such as vertigo, metallic taste, phosphenes as well as on the occurrence of neurocognitive and neurobehavioral effects. For example, short-term exposure to clinical magnetic fields has been reported to result in a 4% reduction in speed and precision and a 16% reduction in visual contrast sensitivity at close range. Both eye-hand precision and coordination speed are affected. The long-term studies concern, among other things, the influence of magnetic fields on sleep quality, which could be linked to an increased risk of accidents. The data on the exposure of healthcare workers to magnetic fields during pregnancy is consistently outdated. However, it has been concluded that there are no particular deviations with regard to the duration of pregnancy, premature births, miscarriages, and birth weight. Epidemiological studies are lacking. With a focus on healthcare personnel, there is a considerable need for high-quality data, particularly on the consequences of long-term exposure to electromagnetic fields from clinical MRI and the effects on pregnancy.

Key points

- Short-term sensory effects such as vertigo, metallic taste, phosphenes as well as neurocognitive and neurological behavioral effects may occur upon exposure to magnetic fields.
- Long-term effects mainly concern quality of sleep, which can be associated with an increased risk of accidents.
- When pregnant women were exposed to magnetic fields, no particular deviations were found with regard to the duration of pregnancy, premature births, miscarriages, and birth weight.

Open access paper: https://www.thieme-connect.com/products/ejournals/html/10.1055/a-2296-3860?articleLanguage=en

--

A Reexamination of Peto's Paradox: Insights Gained from Human Adaptation to Varied Levels of Ionizing and Non-ionizing Radiation

Mortazavi SMJ, Zare O, Ghasemi L, Taghizadeh P, Faghani P, Arshadi M, Mortazavi SAR, Sihver L. A Reexamination of Peto's Paradox: Insights Gained from Human Adaptation to Varied Levels of Ionizing and Non-ionizing Radiation. J Biomed Phys Eng. 2024 Jun 1;14(3):309-314. doi: 10.31661/jbpe.v0i0.2402-1729.

Abstract

Humans have generally evolved some adaptations to protect against UV and different levels of background ionizing radiation. Similarly, elephants and whales have evolved adaptations to protect against cancer, such as multiple copies of the tumor suppressor gene p53, due to their large size and long lifespan. The difference in cancer protection strategies between humans and elephants/whales depends on genetics, lifestyle, environmental exposures, and evolutionary pressures. In this paper, we discuss how the differences in evolutionary adaptations between humans and elephants could explain why elephants have evolved a protective mechanism against cancer, whereas humans have not. Humans living in regions with high levels of background radiation, e.g. in Ramsar, Iran where exposure rates exceed those on the surface of Mars, seem to have developed some kind of protection against the ionizing radiation. However, humans in general have not developed cancer-fighting adaptations, so they instead rely on medical technologies and interventions. The difference in cancer protection strategies between humans and elephants/whales depends on genetics, lifestyle, environmental exposures, and evolutionary pressures. In this paper, we discuss how the differences in evolutionary adaptations between humans and elephants could explain why elephants have evolved a protective mechanism against cancer, whereas humans have not. Studying elephant adaptations may provide insights into new cancer prevention and treatment strategies for humans, but further research is required to fully understand the evolutionary disparities.

Open access paper: https://jbpe.sums.ac.ir/article_50037.html

--

A review on point mutations via proton transfer in DNA base pairs in the absence and presence of electric fields

Pushkaran AC, Arabi AA. A review on point mutations via proton transfer in DNA base pairs in the absence and presence of electric fields. Int J Biol Macromol. 2024 Jul 26:134051. doi: 10.1016/j.ijbiomac.2024.134051.

Abstract

This is a comprehensive review that focuses on spontaneous mutations that may occur during DNA replication, the fundamental process responsible for transferring genetic information. In 1963, Löwdin postulated that these mutations are primarily a result of proton transfer reactions within the hydrogen-bonded DNA base pairs. The single and double proton transfer reactions within the base pairs result in

zwitterions and rare tautomers, respectively. For persistent mutations, these products must be generated at high rates and should be thermodynamically stable. This review covers the proton transfer reactions studied experimentally and computationally. The review also examines the influence of externally applied electric fields on the thermodynamics and kinetics of proton transfer reactions within DNA base pairs, and their biological implications.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0141813024048566

--

Corrigendum to "The effects of radiofrequency electromagnetic fields exposure on human self-reported symptoms: A systematic review of human experimental studies"

Bosch-Capblanch X, Esu E, Moses Oringanje C, Dongus S, Jalilian H, Eyers J, Auer C, Meremikwu M, Röösli M. Corrigendum to "The effects of radiofrequency electromagnetic fields exposure on human self-reported symptoms: A systematic review of human experimental studies" [Environ. Int. 187 (2024) 108612]. Environ Int. 2024 Jul 20:108892. doi: 10.1016/j.envint.2024.108892. Epub ahead of print. Erratum for: Environ Int. 2024 May;187:108612. doi: 10.1016/j.envint.2024.108612. PMID: 39034214.

No abstract

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0160412024004781?via%3Dihub

--

Investigating the impact of anthropogenic radiofrequency electromagnetic fields on animals and plants in the environment: analysis from a systematic map

Brzozek C, Mate R, Bhatt CR, Loughran S, Wood AW, Karipidis K. (2024). Investigating the impact of anthropogenic radiofrequency electromagnetic fields on animals and plants in the environment: analysis from a systematic map. *International Journal of Environmental Studies*, 1–16. doi: 10.1080/00207233.2024.2375861.

Abstract

Radiofrequency electromagnetic fields (RF EMF) from anthropogenic sources have become ubiquitous in the environment. International guidelines have been set by expert scientific bodies to protect human health from excessive exposure. There are no specifically established exposure guidelines to protect plants and animals. A previous systematic map identified 334 studies investigating the effects of RF EMF exposure from anthropogenic sources on animals and plants. This study aimed to analyse the effect of RF EMF and its relationship to different exposure parameters and the quality of those studies. Statistically significant inverse relationships were found between effect size and exposure-level parameters for studies investigating effects in animals. The results indicated that quality score is more indicative of the magnitude of the effect size than exposure-level parameters or exposure duration. This highlights the need for further, high-quality research on the effects of RF EMF exposure on plants and animals.

Quality score

A risk of bias assessment for each study was not conducted as this is more appropriate for a systematic review rather than a systematic map [Citation22]. Instead an assessment of the quality of the studies included in the systematic map was conducted with separate methods used for assessing experimental and observational studies. This was considered important to describe the distribution of quality of evidence across the research in this field. For experimental studies, the method of Vijayalaxmi and Prihoda was used which assessed five quality criteria including adequate dosimetry, use of controls, use of positive controls, use of blinding, use of temperature monitoring [Citation24]. For observational studies, the five quality criteria used in the assessment were as follows: appropriate exposure assessment, appropriate subject selection/comparison groups, consideration of confounders, follow-up assessment, and appropriate outcome assessment. A quality score (QS) was given for each study by two assessors independently, with the final quality score being the average of the two scores. Pearson correlation coefficients were used to assess the reliability and consistency between assessors. The QS was derived by scoring each of the five quality criteria; a score of 1 was awarded when the criterion was adequately addressed, 0.5 awarded when it was partially addressed and a score of 0 when the criterion was not addressed. This resulted in the QS for each study ranging between 0 and 5 with a higher value indicating higher quality for that study. The quality score results were further categorised as studies with QS ≤ 2 being 'poor quality', studies with QS > 2 - < 3.5 as 'average quality' and studies with QS ≥ 3.5 as 'good quality'.

Effect size

Studies investigating the impact of RF EMF on animals and plants have reported a range in the magnitude of effects including no effect and from small effects to large effects. It is therefore important to map the distribution of the effect size (ES) across the different studies, noting that ES = 0 denotes no statistically significant effect and ES > 0 denotes an effect size with a larger number corresponding to a larger effect. We extracted the ES for every study in the systematic map where this could be gleaned from the information provided. The ES was estimated using a number of different methods available in Thalheimer and Cook and according to the type of data provided by each study [Citation23]. Thalheimer and Cook include calculating the effect size from t-test (with and without the standard deviation or standard error) and from F-tests (with and without the mean squared error). Studies that used other statistics did not receive an effect size score. In many of the included studies more than one frequency and exposure intensity was investigated, and in some studies, several endpoints were investigated (and occasionally, more than one animal/plant). In these cases, the largest statistically significant effect was selected to represent that study. For effects that were not statistically significant the ES was set to zero.

Conclusion

The analysis of the systematic map dataset has identified that quality score is more indicative of the magnitude of the effect size than exposure-level parameters or exposure duration in fauna. Similar trends are seen in flora studies; but, these results were not found to be statistically significant. This highlights the need for further, high-quality research.

An investigation of the effect size in relation to different exposure characteristics, as well as the quality of the studies, raises doubts on whether animals and plants are truly affected at levels below human

exposure limits. There are distinct evidence clusters on insect and bird reproduction, development and behaviour for animals, and grain and legume germination and growth for plants that would benefit from specific systematic reviews, with appropriate risk of bias assessments and pooling of the data in meta-analyses. Our review also highlights the need for studies of additional animal and plant species and additional effects that are yet to be investigated. Further, there is a particular need for more observational studies and for an improvement in the quality of all studies.

Disclosure statement

KK, CBr, CBh, RM and SL as part of their employment at ARPANSA provide advice to the Australian Government, Australian States and Territories and the general public on the effects and risks of exposure to ionising and non-ionising radiation. KK and SL are also members of the main commission and scientific expert group, respectively, of the International Commission on Non-Ionising Radiation Protection where they contribute in the development and dissemination of science-based advice on limiting exposure to non-ionising radiation. KK, AW and CBr are members of systematic review teams into the WHO assessment of health effects of exposure to radiofrequency electromagnetic fields. SL is a member of the WHO Task Group on Radiofrequency Fields and Health Risks. No member of the mapping team screened or extracted data from any study in which he or she is an author.

https://www.tandfonline.com/doi/full/10.1080/00207233.2024.2375861

--

Electrostatic pollination by butterflies and moths

England SJ, Robert D. Electrostatic pollination by butterflies and moths. J R Soc Interface. 2024 Jul;21(216):20240156. doi: 10.1098/rsif.2024.0156.

Abstract

Animals, most notably insects, generally seem to accumulate electrostatic charge in nature. These electrostatic charges will exert forces on other charges in these animals' environments and therefore have the potential to attract or repel other objects, for example, pollen from flowers. Here, we show that butterflies and moths (Lepidoptera) accumulate electrostatic charge while in flight. Then, using finite element analysis, we demonstrate that when within millimetres of a flower, the electrostatic charge of a lepidopteran generates an electric field in excess of 5 kV m^{-1}, and that an electric field of this magnitude is sufficient to elicit contactless pollen transfer from flowers across air gaps onto the body of a butterfly or moth. Furthermore, we see that phylogenetic variations exist in the magnitude and polarity of net charge between different species and families and Lepidoptera. These phylogenetic variations in electrostatic charging correlate with morphological, biogeographical and ecological differences between different clades. Such correlations with biogeographical and ecological differences may reflect evolutionary adaptations towards maximizing or minimizing charge accumulation, in relation to pollination, predation and parasitism, and thus we introduce the idea that electrostatic charging may be a trait upon which evolution can act.

Open access paper: https://royalsocietypublishing.org/doi/10.1098/rsif.2024.0156

--

COSMOS. A methodologically-flawed cohort study of the health effects from exposure to radiofrequency radiation from mobile phone use

Moskowitz JM, Frank JW, Melnick RL, Hardell L, Belyaev I, Héroux P, Kelley E, Lai H, Maisch D, Mallery-Blythe E, Philips A; International Commission on the Biological Effects of Electromagnetic Fields. COSMOS: A methodologically-flawed cohort study of the health effects from exposure to radiofrequency radiation from mobile phone use. Environ Int. 2024 Jun 21;190:108807. doi: 10.1016/j.envint.2024.108807.

No abstract

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024003933

--

Response to the letter to the editor regarding "Mobile phone use and brain tumour risk – COSMOS, a prospective cohort study"

Feychting M, Schüz J, Toledano MB, Vermeulen R, Auvinen A, Poulsen AH, Deltour I, Smith RB, Heller J, Kromhout H, Huss A, Johansen C, Tettamanti G, Elliott P. Response to the Letter to the Editor regarding "Mobile phone use and brain tumour risk - COSMOS, a prospective cohort study". Environ Int. 2024 May;187:108664. doi: 10.1016/j.envint.2024.108664.

No abstract

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024003945

--

Summary of seven Swedish case reports on the microwave syndrome associated with 5G radiofrequency radiation

Hardell L, Nilsson M. Summary of seven Swedish case reports on the microwave syndrome associated with 5G radiofrequency radiation. Rev Environ Health. 2024 Jun 19. doi: 10.1515/reveh-2024-0017.

Abstract

The fifth generation, 5G, for wireless communication is currently deployed in Sweden since 2019/2020, as well as in many other countries. We have previously published seven case reports that include a total of 16 persons aged between 4 and 83 years that developed the microwave syndrome within a short time after being exposed to 5G base stations close to their dwellings. In all cases high radiofrequency (RF) radiation from 4G/5G was measured with a broadband meter. RF radiation reached >2,500,000 to >3,180,000 μW/m^2 in peak maximum value in three of the studies. In total 41 different health issues were assessed for each person graded 0 (no complaint) to 10 (worst symptoms). Most prevalent and severe were sleeping difficulty (insomnia, waking night time, early wake-up), headache, fatigue, irritability, concentration problems, loss of immediate memory, emotional distress, depression tendency, anxiety/panic, dysesthesia (unusual touched based sensations), burning and lancinating skin, cardiovascular symptoms (transitory high or irregular pulse), dyspnea, and pain in muscles and joints. Balance disorder and tinnitus were less prevalent. All these symptoms are included in the microwave syndrome. In most cases the symptoms declined and disappeared within a short time period after the studied persons had moved to a place with no 5G. These case histories are classical examples of

provocation studies. They reinforce the urgency to inhibit the deployment of 5G until more safety studies have been performed.

Open access paper: https://www.degruyter.com/document/doi/10.1515/reveh-2024-0017/html

--

Follow-Up Research on NTP's Clear Evidence on RF Causing Malignant Tumors in Rats

Lin JC. Follow-Up Research on NTP's Clear Evidence on RF Causing Malignant Tumors in Rats [Health Matters]. *IEEE Microwave Magazine*. 25(6):16-18. June 2024. doi: 10.1109/MMM.2024.3378608.

Abstract

The recent disclosure of the U.S. National Toxicology Program (NTP) closing down its RF radiation research program on how RF radiation causes cancer was a surprise [1], [2], [3]. NTP is a part of the National Institutes of Health and reports to the U.S. Department of Health and Human Services. NTP's mandates are to provide the scientific basis for programs, activities, and policies that promote health or lead to the prevention of disease.

Excerpts

In 2018, NTP published the final report on its US$30 million laboratory research showing "clear evidence" that lifelong exposure to low-level RF radiation caused cancers in rats [6], [7], [8]. The statistically significant findings showed that both GSM- and CDMA-modulated 900-MHz RF radiation had led to the development of malignant schwannoma, a rare form of tumor in the hearts of male rats. Furthermore, an independent analysis of the NTP data for overall cancer incidence detected in any organ or tissue inside the animal showed that rats exposed to GSM and CDMA cellphone RF radiation had significantly higher overall or total primary tumor incidence than the concurrent controls....

The NTP findings of clear evidence that RF radiation can cause cancer was already reaffirmed by the Ramazzini Institute study from Italy [10]....

In 2019, NTP announced a new genotoxicology project designed to examine how RF radiation causes cancer....

Apparently, now, close to five years later, the new project has never progressed beyond the feasibility studies with the compact exposure chambers. Moreover, the NTP located in Research Triangle Park, NC, USA, no longer plans to study biological effects from cellphone RF radiation. NTP has decided that "the research was technically challenging and more resource-intensive than expected." This effectively marks the end of the nonmilitary governmental project in the United States to study the biological effects of cellphone or RF radiation. Any future NTP publications on the topic would come from the feasibility studies only [3].

Unfortunately, whether the U.S. civilian government halts doing the important research on mechanistic causes for cancer from low-level chronic exposure to RF radiation or not, nearly 100% of all Americans are being cast as subjects in a massive health experiment without a formal protocol. Furthermore, billions of people worldwide, including the young and older adults, are being exposed to unnecessary levels of human-made wireless RF radiation.

The obvious question is, what happened? Should we all forget about NTP's findings of clear scientific evidence that lifelong exposure to low-level RF radiation causes cancer from a federal government health agency that prides itself in telling the nation that its product is "science you can depend on for decisions that matter"?

Postscript: It is interesting to recall that the recently announced termination of NTP's RF research program on how RF radiation causes cancer practically halts most, if not all, biological research of RF radiation supported by the civilian U.S. government. In contrast, the military arm of the U.S. government through its Defense Advanced Research Projects Agency (DARPA) has initiated a new research program on the roles RF radiation may have in biological systems [11]. DARPA's historical and contemporary contributions to scientific and technological advancements are widely known. Thus, it poses the question, what does DARPA know about RF radiation on biological systems?

https://ieeexplore.ieee.org/document/10523054

--

Epidemiological exploration of the impact of bluetooth headset usage on thyroid nodules using Shapley additive explanations method

Zhou N, Qin W, Zhang JJ, Wang Y, Wen JS, Lim YM. Epidemiological exploration of the impact of bluetooth headset usage on thyroid nodules using Shapley additive explanations method. Sci Rep. 2024 Jun 21;14(1):14354. doi: 10.1038/s41598-024-63653-0.

Abstract

With an increasing prevalence of thyroid nodules globally, this study investigates the potential correlation between the use of Bluetooth headsets and the incidence of thyroid nodules, considering the cumulative effects of non-ionizing radiation (NIR) emitted by these devices. In this study, we analyzed 600 valid questionnaires from the WenJuanXing platform using Propensity Score Matching (PSM) and the XGBOOST model, supplemented by SHAP analysis, to assess the risk of thyroid nodules. PSM was utilized to balance baseline characteristic differences, thereby reducing bias. The XGBOOST model was then employed to predict risk factors, with model efficacy measured by the area under the Receiver Operating Characteristic (ROC) curve (AUC). SHAP analysis helped quantify and explain the impact of each feature on the prediction outcomes, identifying key risk factors. Initially, 600 valid questionnaires from the WenJuanXing platform underwent PSM processing, resulting in a matched dataset of 96 cases for modeling analysis. The AUC value of the XGBOOST model reached 0.95, demonstrating high accuracy in differentiating thyroid nodule risks. SHAP analysis revealed age and daily Bluetooth headset usage duration as the two most significant factors affecting thyroid nodule risk. Specifically, longer daily usage durations of Bluetooth headsets were strongly linked to an increased risk of developing thyroid nodules, as indicated by the SHAP analysis outcomes. Our study highlighted a significant impact relationship between prolonged Bluetooth headset use and increased thyroid nodule risk, emphasizing the importance of considering health impacts in the use of modern technology, especially for devices like Bluetooth headsets that are frequently used daily. Through precise model predictions and variable importance analysis, our research provides a scientific basis for the formulation of public health policies and personal health habit choices, suggesting that attention should be paid to the duration of Bluetooth headset use in daily life to reduce the potential risk of thyroid nodules. Future

research should further investigate the biological mechanisms of this relationship and consider additional potential influencing factors to offer more comprehensive health guidance and preventive measures.

Open access paper: https://www.nature.com/articles/s41598-024-63653-0

--

The effects of radiofrequency exposure on male fertility: A systematic review of human observational studies with dose–response meta-analysis (SR 3--male)

Kenny RPW, Johnson EE, Adesanya AM, Richmond C, Beyer F, Calderon C, Rankin J, Pearce MS, Toledano M, Craig D, Pearson F. The effects of radiofrequency exposure on male fertility: A systematic review of human observational studies with dose–response meta-analysis. Environment International, 2024. doi: 10.1016/j.envint.2024.108817.

Abstract

Background The World Health Organization (WHO) is bringing together evidence on radiofrequency electromagnetic field (RF-EMF) exposure in relation to health outcomes, previously identified as priorities for research and evaluation by experts in the field, to inform exposure guidelines. A suite of systematic reviews have been undertaken by a network of topic experts and methodologists to collect, assess and synthesise data relevant to these guidelines. Following the WHO handbook for guideline development and the COSTER conduct guidelines, we systematically reviewed the evidence on the potential effects of RF-EMF exposure on male fertility in human observational studies.

Methods We conducted a broad and sensitive search for potentially relevant records within the following bibliographic databases: MEDLINE; Embase; Web of Science and EMF Portal. We also conducted searches of grey literature through relevant databases including OpenGrey, and organisational websites and consulted RF-EMF experts. We hand searched reference lists of included study records and for citations of these studies. We included quantitative human observational studies on the effect of RF-EMF exposure in adult male participants on infertility: sperm concentration; sperm morphology; sperm total motility; sperm progressive motility; total sperm count; and time to pregnancy. Titles and abstracts followed by full texts were screened in blinded duplicate against pre-set eligibility criteria with consensus input from a third reviewer as required. Data extraction from included studies was completed by two reviewers, as was risk of bias assessment using the Office of Health Assessment and Translation (OHAT) tool. We conducted a dose–response meta-analysis as possible and appropriate. Certainty of the evidence was assessed by two reviewers using the OHAT GRADE tool with input from a third reviewer as required.

Results We identified nine studies in this review; seven were general public studies (with the general public as the population of interest) and two were occupational studies (with specific workers/workforces as the population of interest). General public studies. Duration of phone use: The evidence is very uncertain surrounding the effects of RF-EMF on sperm concentration (10/6 mL) (MD (mean difference) per hour of daily phone use 1.6 106/mL, 95 % CI −1.7 to 4.9; 3 studies), sperm morphology (MD 0.15 percentage points of deviation of normal forms per hour, 95 % CI −0.21 to 0.51; 3 studies), sperm progressive motility (MD −0.46 percentage points per hour, 95 % CI −1.04 to 0.13; 2 studies) and total sperm count (MD per hour −0.44 106/ejaculate, 95 % CI −2.59 to 1.7; 2 studies) due to

very low-certainty evidence. Four additional studies reported on the effect of mobile phone use on sperm motility but were unsuitable for pooling; only one of these studies identified a statistically significant effect. All four studies were at risk of exposure characterisation and selection bias; two of confounding, selective reporting and attrition bias; three of outcome assessment bias and one used an inappropriate statistical method. Position of phone: There may be no or little effect of carrying a mobile phone in the front pocket on sperm concentration, total count, morphology, progressive motility or on time to pregnancy. Of three studies reporting on the effect of mobile phone location on sperm total motility and, or, total motile count, one showed a statistically significant effect. All three studies were at risk of exposure characterisation and selection bias; two of confounding, selective reporting and attrition bias; three of outcome assessment bias and one used inappropriate statistical method. RF-EMF Source: One study indicates there may be little or no effect of computer or other electric device use on sperm concentration, total motility or total count. This study is at probably high risk of exposure characterisation bias and outcome assessment bias. Occupational studies. With only two studies of occupational exposure to RF-EMF and heterogeneity in the population and exposure source (technicians exposed to microwaves or seamen exposed to radar equipment), it was not plausible to statistically pool findings. One study was at probably or definitely high risk of bias across all domains, the other across domains for exposure characterisation bias, outcome assessment bias and confounding.

Discussion

The majority of evidence identified was assessing localised RF-EMF exposure from mobile phone use on male fertility with few studies assessing the impact of phone position. Overall, the evidence identified is very uncertain about the effect of RF-EMF exposure from mobile phones on sperm outcomes. One study assessed the impact of other RF-EMF sources on male fertility amongst the general public and two studies assessed the impact of RF-EMF exposure in occupational cohorts from different sources (radar or microwave) on male fertility. Further prospective studies conducted with greater rigour (in particular, improved accuracy of exposure measurement and appropriate statistical method use) would build the existing evidence base and are required to have greater certainty in any potential effects of RF-EMF on male reproductive outcomes.

Prospero Registration: CRD42021265401 (SR3A)

Excerpts

Conclusions Overall, the evidence is very uncertain about the effect of RF-EMF on male fertility outcomes. Where dose–response *meta*-analysis was possible, the evidence was rated as very low-certainty on OHAT GRADE, while the majority of included studies were at risk of bias and only a small number of studies reported on each outcome of interest. Given this, we cannot be confident in what the current body of research concludes about the effect of RF-EMF on male fertility.

Further prospective studies conducted with greater rigour (in particular, improved accuracy of exposure measurement and appropriate statistical method use) are required to build on the existing evidence base and provide greater certainty in any potential effects of RF-EMF on male reproductive outcomes.

Competing interests Carolina Calderon was involved in both MOBI-Kids (risk of brain cancer from exposure to radiofrequency fields in childhood and adolescence) and GERoNIMO, Tsarna et al. 2019 was

one of the outcomes of this project. Although Carolina was not involved in the publication, with her involvement being in the intermediate frequency exposure assessment.

Mireille Toledano has been involved in funded research assessing mobile phone and other wireless technologies usage on health outcomes. The SCAMP (study cognition adolescents and mobile phones) prospective cohort study which is currently ongoing (2015–2021). The COSMOS (cohort study of mobile phone use and health) a longitudinal cohort study which is completed (2019).

Declaration of competing interest The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements We would like to thank Anastasija Jadrevska, Claudia Preuss, and Susanna Lagorio for providing translations. Additionally, we would like to thank Yangzihan Wang for translation and data extraction of studies in Chinese. We would like to thank Jos Verbeek, Emilie Van Deventer and Maria Feychting for their expertise and support throughout the systematic review.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024004033

--

The effects of radiofrequency exposure on adverse female reproductive outcomes: A systematic review of human observational studies with dose–response meta-analysis (SR 3--female)

Johnson EE, Kenny RPW, Adesanya AM, Richmond C, Beyer F, Calderon C, Rankin J, Pearce MS, Toledano M, Craig D, Pearson F. The effects of radiofrequency exposure on adverse female reproductive outcomes: A systematic review of human observational studies with dose–response meta-analysis. Environment International, 2024. doi: 10.1016/j.envint.2024.108816.

Abstract

Background To inform radiofrequency electromagnetic field (RF-EMF) exposure guidelines the World Health Organization (WHO) is bringing together evidence on RF-EMF in relation to health outcomes prioritised for evaluation by experts in this field. Given this, a network of topic experts and methodologists have conducted a series of systematic reviews collecting, assessing, and synthesising data of relevance to these guidelines. Here we present a systematic review of the effect of RF-EMF exposure on adverse pregnancy outcomes in human observational studies which follows the WHO handbook for guideline development and the COSTER conduct guidelines.

Methods We conducted a broad, sensitive search for potentially relevant records within the following bibliographic databases: MEDLINE; Embase; and the EMF Portal. Grey literature searches were also conducted through relevant databases (including OpenGrey), organisational websites and via consultation of RF-EMF experts. We included quantitative human observational studies on the effect of RF-EMF exposure in adults' preconception or pregnant women on pre-term birth, small for gestational age (SGA; associated with intrauterine growth restriction), miscarriage, stillbirth, low birth weight (LBW) and congenital anomalies. In blinded duplicate, titles and abstracts then full texts were screened against eligibility criteria. A third reviewer gave input when consensus was not reached. Citation chaining of included studies was completed. Two reviewers' data extracted and assessed included studies for risk of

bias using the Office of Health Assessment and Translation (OHAT) tool. Random effects meta-analyses of the highest versus the lowest exposures and dose–response meta-analysis were conducted as appropriate and plausible. Two reviewers assessed the certainty in each body of evidence using the OHAT GRADE tool.

Results We identified 18 studies in this review; eight were general public studies (with the general public as the population of interest) and 10 were occupational studies (with the population of interest specific workers/workforces). General public studies. From pairwise meta-analyses of general public studies, the evidence is very uncertain about the effects of RF-EMF from mobile phone exposure on preterm birth risk (relative risk (RR) 1.14, 95% confidence interval (CI): 0.97–1.34, 95% prediction interval (PI): 0.83–1.57; 4 studies), LBW (RR 1.14, 95% CI: 0.96–1.36, 95% PI: 0.84–1.57; 4 studies) or SGA (RR 1.13, 95% CI: 1.02–1.24, 95% PI: 0.99–1.28; 2 studies) due to very low-certainty evidence. It was not feasible to meta-analyse studies reporting on the effect of RF-EMF from mobile phone exposure on congenital anomalies or miscarriage risk. The reported effects from the studies assessing these outcomes varied and the studies were at some risk of bias. No studies of the general public assessed the impact of RF-EMF exposure on stillbirth. Occupational studies. In occupational studies, based on dose–response meta-analyses, the evidence is very uncertain about the effects of RF-EMF amongst female physiotherapists using shortwave diathermy on miscarriage due to very low-certainty evidence (OR 1.02 95% CI 0.94–1.1; 2 studies). Amongst offspring of female physiotherapists using shortwave diathermy, the evidence is very uncertain about the effects of RF-EMF on the risk of congenital malformations due to very low-certainty evidence (OR 1.4, 95% CI 0.85 to 2.32; 2 studies). From pairwise meta-analyses, the evidence is very uncertain about the effects of RF-EMF on the risk of miscarriage (RR 1.06, 95% CI 0.96 to 1.18; very low-certainty evidence), pre-term births (RR 1.19, 95% CI 0.32 to 4.37; 3 studies; very low-certainty evidence), and low birth weight (RR 2.90, 95% CI: 0.69 to 12.23; 3 studies; very low-certainty evidence). Results for stillbirth and SGA could not be pooled in meta-analyses. The results from the studies reporting these outcomes were inconsistent and the studies were at some risk of bias.

Discussion Most of the evidence identified in this review was from general public studies assessing localised RF-EMF exposure from mobile phone use on female reproductive outcomes. In occupational settings, each study was of heterogenous whole-body RF-EMF exposure from radar, short or microwave diathermy, surveillance and welding equipment and its effect on female reproductive outcomes. Overall, the body of evidence is very uncertain about the effect of RF-EMF exposure on female reproductive outcomes. Further prospective studies conducted with greater rigour (particularly improved accuracy of exposure measurement and using appropriate statistical methods) are required to identify any potential effects of RF-EMF exposure on female reproductive outcomes of interest.

Excerpts

Summary of the evidence and interpretation of the results In total, 18 studies were identified for this review: in eight the general public was the population of interest and in 10 the populations of interest were occupational. Within the general population, the evidence is very uncertain about the effects of RF-EMF on pre-term birth, SGA and low birth weight. It was not possible to conduct meta-analyses for miscarriage and congenital anomalies due to significant heterogeneity between studies. Within the occupational studies, the evidence is very uncertain about the effects of RF-EMF on miscarriage and the effects of maternal RF-EMF exposure on pre-term birth, congenital anomalies and low birth weight.

Limitations of the evidence The are multiple limitations with the evidence base of human observational studies assessing the effect of localised and whole-body RF-EMF exposure on pre-term birth; SGA; miscarriage; still birth and, or, congenital anomalies compared to no or low level exposure in preconception or pregnant adults.

Risk of bias was often apparent in both general public and occupational studies; all but one of the included studies were rated as probably or definitely high risk for at least two domains on the OHAT risk of bias tool. Nine studies were at probably or definitely high risk of exposure characterisation bias (Allam, 2016, Baste et al., 2012, Cromie et al., 2002, Källén et al., 1982, Karuserci et al., 2019, Lu et al., 2017, Taskinen et al., 1990, Tsarna et al., 2019, Zhao et al., 2021), while another nine were at risk of outcome assessment bias (Baste et al., 2012, Cromie et al., 2002, Källén et al., 1982, Karuserci et al., 2019, Lu et al., 2017, Mahmoudabadi et al., 2015, Tsarna et al., 2019, Xu et al., 2016, Zhao et al., 2021). In 10 studies, issues surrounding the identification and handling of confounders was present (Allam, 2016, Baste et al., 2012, Baste et al., 2015, Brizzi and Marinelli, 2018, Cromie et al., 2002, Källén et al., 1982, Karuserci et al., 2019, Kolmodin-Hedman et al., 1988, Mahmoudabadi et al., 2015, Xu et al., 2016), while three were also at risk of selective reporting (Cromie et al., 2002, Källén et al., 1982, Kolmodin-Hedman et al., 1988).

More generally, reporting of exposures across studies was often inconsistent and lacking in detail (see Table 1, Table 2). Most studies in the general population used a proxy exposure, such as time spent on mobile phone or mobile phone usage, with heterogeneity between how these were measured.

One of the main issues of the evidence base is a lack of confounding assessment in studies (Baste et al., 2015, Allam, 2016, Brizzi and Marinelli, 2018, Karuserci et al., 2019, Kolmodin-Hedman et al., 1988, Baste et al., 2012, Källén et al., 1982, Cromie et al., 2002, Mahmoudabadi et al., 2015, Xu et al., 2016) and inconsistency regarding the method of exposure measurement. All analyses were downgraded to either low-certainty or very low-certainty evidence on OHAT GRADE, with risk of bias, indirectness and imprecision causing concerns. Indirectness was an issue for occupational studies measuring congenital anomalies. Many of the studies only stated that they were assessing the risk of any congenital anomalies rather than identifying and assessing the risk of specific congenital anomalies (e.g. the effect of RF-EMF could potentially affect different congenital anomalies in different ways). This limits our ability to assess the risk of different kinds of congenital anomalies following RF-EMF exposure.

Conclusions Overall, the majority of evidence suggest that there is little to no effect of RF-EF on female reproductive outcomes. The evidence was rated as low to very low certainty, was at risk of bias and only a small number of studies reported on each outcome of interest. Given this we cannot be confident in what the body of research is indicating about the effect of RF-EMF on female reproductive outcomes. The *meta*-analyses that were possible suggest no increased relative risk for female reproductive outcomes due to RF-EMF exposure in the general public or occupational settings. This is further supported by the dose–response *meta*-analysis in female physiotherapists using SWD. For studies that were not *meta*-analysed, there was variation in effect. Most included studies were at risk of bias.

Overall, while we observe a lack of effect, further prospective studies conducted with greater rigour would build the existing evidence base and are required to have greater certainty in any potential effects of RF-EMF on female reproductive outcomes.

Registration and protocol The protocol was published in Environment International (Kenny et al., 2022) and an abridged version is also available on PROSPERO (CRD42021265401; referred to as SR3B).

Declaration of competing interest The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: [Mireille Toledano has been involved in funded research assessing mobile phone and other wireless technologies usage on health outcomes: the SCAMP (study cognition adolescents and mobile phones) prospective cohort study which is currently ongoing (2015–2021), and the COSMOS (cohort study of mobile phone use and health) a longitudinal cohort study which is completed (2019). Carolina Calderon has been involved in MOBI-Kids (risk of brain cancer from exposure to radiofrequency fields in childhood adolescence) and GERoNIMO, of which the Tsarna 2019 study was one of the outcomes. However, they were not directly involved in the Tsarna 2019 paper and was not involved in the selection, data extraction or risk of bias assessment for this study.].

Acknowledgements We would like to thank Jos Verbeek, Emilie Van Deventer and Maria Feychting for their expertise and support throughout the systematic review.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024004021

--

Detrimental impact of cell phone radiation on sperm DNA integrity

Koohestanidehaghi Y, Khalili MA, Dehghanpour F, Seify M. Detrimental impact of cell phone radiation on sperm DNA integrity. Clinical and Experimental Reproductive Medicine 2024;51(1):13-19.

Abstract

Radiofrequency electromagnetic radiation (RF-EMR) from various sources may impact health due to the generation of frequency bands. Broad pulses emitted within frequency bands can be absorbed by cells, influencing their function. Numerous laboratory studies have demonstrated that mobile phones—generally the most widely used devices—can have harmful effects on sex cells, such as sperm and oocytes, by producing RF-EMR. Moreover, some research has indicated that RF-EMR generated by mobile phones can influence sperm parameters, including motility, morphology, viability, and (most critically) DNA structure. Consequently, RF-EMR can disrupt both sperm function and fertilization. However, other studies have reported that exposure of spermatozoa to RF-EMR does not affect the functional parameters or genetic structure of sperm. These conflicting results likely stem from differences among studies in the duration and exposure distance, as well as the species of animal used. This report was undertaken to review the existing research discussing the effects of RF-EMR on the DNA integrity of mammalian spermatozoa.

Conclusion

EMW can induce oxidative stress, which subsequently leads to disorders such as reduced mobility, morphological changes, acrosome disturbances, and ultimately, damage to the nucleus and genetic material. This oxidative damage to DNA can result in the breakdown of both single-stranded and double-stranded DNA structures, culminating in fragmentation. If the DNA is not repaired and the damage accumulates, the sperm may undergo apoptosis. Damage to the sperm genome can ultimately impact

fertility, potentially leading to infertility. Therefore, it is advisable to limit daily exposure to these sources to prevent irreversible damage caused by EMWs. Many men carry their cell phones in their trouser pockets or clipped to their belts, and the use of Bluetooth can increase their susceptibility to RF-EMR exposure. This exposure can induce changes in sperm quality through oxidative stress, potentially leading to infertility. Agarwal et al. [11] suggested that carrying a cell phone in a pocket could lead to a decline in sperm quality. However, it is important to note that the phone and male reproductive organs are separated by multiple tissue layers. Therefore, extrapolating these in vitro effects to real-life conditions requires further studies [11].

In July 2021, the European Parliament commissioned a research report titled “Health impact of 5G.” The report concluded that the commonly used RF-EMFs are likely carcinogenic to humans and have a definitive impact on male fertility. It also suggested potential adverse effects on the development of embryos, fetuses, and newborns. To mitigate these adverse effects, the organization proposed several strategies. These include favoring non-wireless connections, increasing distance from the source of RF-EMFs, switching off devices when not in use, and practicing safe phone usage [55].

Open access paper: https://ecerm.org/journal/view.php?doi=10.5653/cerm.2023.06121

--

A comprehensive review of 5G NR RF-EMF exposure assessment technologies: fundamentals, advancements, challenges, niches, and implications

Korkmaz E, Aerts S, Coesoij R, Bhatt CR, Velghe M, Colussi L, Land D, Petroulakis N, Spirito M, Bolte J. A comprehensive review of 5G NR RF-EMF exposure assessment technologies: fundamentals, advancements, challenges, niches, and implications. Environ Res. 2024 Jul 5:119524. doi: 10.1016/j.envres.2024.119524.

Abstract

This review offers a detailed examination of the current landscape of radio frequency (RF) electromagnetic field (EMF) assessment tools, ranging from spectrum analyzers and broadband field meters to area monitors and custom-built devices. The discussion encompasses both standardized and non-standardized measurement protocols, shedding light on the various methods employed in this domain. Furthermore, the review highlights the prevalent use of mobile apps for characterizing 5G-NR radio network data. A growing need for low-cost measurement devices is observed, commonly referred to as ”sensors” or ”sensor nodes,” that are capable of enduring diverse environmental conditions. These sensors play a crucial role in both microenvironmental surveys and individual exposures, enabling stationary, mobile, and personal exposure assessments based on body-worn sensors, across wider geographical areas. This review revealed a notable need for cost-effective and long-lasting sensors, whether for individual exposure assessments, mobile (vehicle-integrated) measurements, or incorporation into distributed sensor networks. However, there is a lack of comprehensive information on existing custom-developed RF-EMF measurement tools, especially in terms of measuring uncertainty. Additionally, there is a need for real-time, fast-sampling solutions to understand the highly irregular temporal variations EMF distribution in next-generation networks.

Given the diversity of tools and methods, a comprehensive comparison is crucial to determine the necessary statistical tools for aggregating the available measurement data.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0013935124014294

--

Under the Hood of Electromagnetic Field Estimation and Evaluation in 5G Networks

Estrada-Jimenez JC, Pardo E, Roth U, L. Selmane L, Faye S. Under the Hood of Electromagnetic Field Estimation and Evaluation in 5G Networks. *IEEE Access*, doi: 10.1109/ACCESS.2024.3418301.

Abstract

The estimation of Electromagnetic Field (EMF) exposure is critical for evaluating the potential health risks associated with wireless network implementation. With the advent of new communication technologies, such as 5G and 6G, new methodologies are required. Indeed, higher frequency ranges are being studied, necessitating an increased number of base stations for service provision. In this paper, we introduce a novel methodology for EMF estimation in 5G mobile networks. We also evaluate common scenarios that mobile network operators frequently encounter, resulting in additional exposure, such as interference, overpropagation, and the presence of other users. Our approach is validated through a combination of simulated values and real data collected from sensors deployed in the field. Finally, a digital twin approach is proposed to combine the simulated values with the real data obtained from radio frequency sensors. The scenario is deployed in Belval, Luxembourg.

Open access paper: https://ieeexplore.ieee.org/document/10568919

--

Demonstration of Safe Electromagnetic Radiation Emitted by 5G Active Antenna Systems

Kumar S, Sheemar CK, Astro A, Querol J, Chatzinotas S. Demonstration of Safe Electromagnetic Radiation Emitted by 5G Active Antenna Systems. arXiv:2406.07910v1 [cs.ET] 12 Jun 2024.

Abstract

The careful planning and safe deployment of 5G technologies will bring enormous benefits to society and economy. Higher frequency, beamforming and small-cells are key technologies that will provide unmatched throughput and seamless connectivity to the 5G users. Superficial knowledge of these technologies has raised concerns among the general public about the harmful effects of radiation. Several standardization bodies are active to put limits on the emissions which are based on a defined set of radiation measurement methodologies. However, due to the peculiarity of 5G such as dynamicity of the beams, network densification, Time Division Duplexing mode of operation, etc, using existing EMF measurement methods may provide inaccurate results. In this context, we discuss our experimental studies aimed towards the measurement of radiation caused by beam-based transmissions from 5G base-station equipped with an Active Antenna System (AAS). We elaborate on the shortcomings of current measurement methodologies and address several open questions. Next, we demonstrate that using user-specific downlink beamforming, not only better performance is achieved compared to non-beamformed downlink, but also the radiation in the vicinity of the intended user is significantly

decreased. Further, we show that under weak reception conditions, an uplink transmission can cause significantly high radiation in the vicinity of the user-equipment. We believe that our work will help in clearing several misleading concepts about the 5G EMF radiation effects. We conclude the work by providing guidelines to improve the methodology of EMF measurement by considering the spatio-temporal dynamicity of the 5G transmission.

Open access paper: https://arxiv.org/html/2406.07910v1

--

Smartphone-Based Methodology Applied to Electromagnetic Field Exposure Assessment

López-Espí PL, Sánchez-Montero R, Guillén-Pina J, Castro-Sanz R, Chocano-Del-Cerro R, Martínez-Rojas JA. Smartphone-Based Methodology Applied to Electromagnetic Field Exposure Assessment. Sensors (Basel). 2024 May 31;24(11):3561. doi: 10.3390/s24113561.

Abstract

This study presents the measurements of exposure to electromagnetic fields, carried out comparatively following standard methods from fixed sites using a broadband meter and using a smartphone on which an App designed for this purpose has been installed. The results of two measurement campaigns carried out on the campus of the University of Alcalá over an area of 1.9 km^2 are presented. To characterize the exposure, 20 fixed points were measured in the first case and 860 points along the route made with a bicycle in the last case. The results obtained indicate that there is proportionality between the two methods, making it possible to use the smartphone for comparative measurements. The presented methodology makes it possible to characterize the exposure in the area under study in four times less time than that required with the traditional methodology

Open access paper: https://www.mdpi.com/1424-8220/24/11/3561

--

A boundary element method of bidomain modeling for predicting cellular responses to electromagnetic fields

Czerwonky DM, Aberra AS, Gomez LJ. A boundary element method of bidomain modeling for predicting cellular responses to electromagnetic fields. J Neural Eng. 2024 Jun 11. doi: 10.1088/1741-2552/ad5704.

Abstract

Objective: Commonly used cable equation approaches for simulating the effects of electromagnetic fields on excitable cells make several simplifying assumptions that could limit their predictive power. Bidomain or “whole” finite element methods have been developed to fully couple cells and electric fields for more realistic neuron modeling. Here, we introduce a novel bidomain integral equation designed for determining the full electromagnetic coupling between stimulation devices and the intracellular, membrane, and extracellular regions of neurons.

Methods: Our proposed boundary element formulation offers a solution to an integral equation that connects the device, tissue inhomogeneity, and cell membrane-induced E-fields. We solve this integral equation using first-order nodal elements and an unconditionally stable Crank-Nicholson time-stepping

scheme. To validate and demonstrate our approach, we simulated cylindrical Hodgkin-Huxley axons and spherical cells in multiple brain stimulation scenarios.

Main Results: Comparison studies show that a boundary element approach produces accurate results for both electric and magnetic stimulation. Unlike bidomain finite element methods, the bidomain boundary element method does not require volume meshes containing features at multiple scales. As a result, modeling cells, or tightly packed populations of cells, with microscale features embedded in a macroscale head model, is simplified, and the relative placement of devices and cells can be varied without the need to generate a new mesh.

Significance: Device-induced electromagnetic fields are commonly used to modulate brain activity for research and therapeutic applications. Bidomain solvers allow for the full incorporation of realistic cell geometries, device E-fields, and neuron populations. Thus, multi-cell studies of advanced neuronal mechanisms would greatly benefit from the development of fast-bidomain solvers to ensure scalability and the practical execution of neural network simulations with realistic neuron morphologies.

Conclusion

We introduced a novel bidomain integral equation for modeling the electric response of neuron cells to device-induced E-fields. Our study includes several canonical test cases with unmyelinated cells, including scenarios with multiple cells, transverse polarization, DBS electrodes at varying proximity to multiple axon geometries, and TMS with a spherical head model. The study results indicate that (1) the hybrid cable-equation approach is a sufficient choice for most simulations, (2) longitudinal stimulation serves as the primary activation mechanism for electromagnetic brain stimulation, and (3) multi-cell studies of advanced mechanisms would greatly benefit from further development of fast-bidomain or hybrid cable-bidomain solvers. Our future efforts will focus on developing fast bidomain solvers to fully incorporate realistic neuron morphologies.

Acknowledgements

Preliminary results for this work have been presented at the Applied Computational Electromagnetics Society (ACES) 2023 conference (March 2023, Monterey, California, USA)[67], the 2023 IEEE MTTS International Conference on Numerical Electromagnetic and Multiphysics Modeling and Optimization (NEMO'2023, June 2023, Winnipeg, Canada), the IEEE International Symposium on Antennas and Propagation Society and USNC-URSI Radio Science Meeting (IEEE-AP-S 2023, July 2023, Portland, Oregon, USA), and the Brain and Human Body Modeling conference 2023 (BHBM 2023, August 2023, Boston, Massachusetts, USA). Additionally, a preprint of this manuscript has been uploaded to https://www.biorxiv.org.

Research reported in this publication was supported by the National Institute of Mental Health of the National Institutes of Health under Award Number R00MH120046. The content of the current research is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Additionally, this effort was sponsored by the **Central Intelligence Agency (CIA)**, through CIA Federal Labs. The U.S. Government is authorized to reproduce and distribute reprints for Governmental

purposes notwithstanding any copyright notation thereon. The views and conclusions contained herein are those of the authors and should not be interpreted as necessarily representing the official policies or endorsements, either expressed or implied, of the U.S. Government or of the Central Intelligence Agency.

Open access paper: https://iopscience.iop.org/article/10.1088/1741-2552/ad5704

--

(In)accuracy and convergent validity of daily end-of-day and single-time self-reported estimations of smartphone use among adolescents

Tkaczyk M, Tancoš M, Smahel D, Elavsky S, Plhák J. (In)accuracy and convergent validity of daily end-of-day and single-time self-reported estimations of smartphone use among adolescents. Computers in Human Behavior. Vol. 158, 2024, doi: 10.1016/j.chb.2024.108281.

Abstract

Understanding the measurement inaccuracy and bias introduced by self-reports of smartphone use is essential for making meaningful inferences about smartphone use and its effects. Evidence for the self-reports of smartphone use in intensive longitudinal studies is largely missing. Based on self-reported and digital trace data from 137 Czech adolescents (41% girls, Mage = 14.95 years), this study examined the accuracy, directional bias, and convergent validity of daily end-of-day and single-time reports of screen time and phone-checking behavior. Overall, the study found considerable discrepancies between self-reported smartphone use and digital trace and low between-person convergent validity for all self-reports considered for the study. Respondents usually reported shorter screen time and lower frequency of phone-checking behavior as compared to digital trace, both in daily and single-time self-reports. The within-person convergent validity between daily reports and digital tracking was low, indicating poor self-reports ability to capture the actual day-to-day fluctuations in smartphone use. This study adds to the existing evidence showing that self-reports based insights into how people use smartphones differ considerably from digital trace data and shows that both person and situational levels contribute to explaining the discrepancy between digital trace and self-report data among adolescents.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0747563224001493

--

Fuzzy-Based Bioengineering System for Predicting and Diagnosing Diseases of the Nervous System Triggered by the Interaction of Industrial Frequency Electromagnetic Fields

Korenevskiy NA, Al-Kasasbeh RT, Krikunova EA, Rodionova SN, Shaqdan A, Al-Habahbeh OM, Filist S, Alshamasin MS, Khrisat MS, Ilyash M. Fuzzy-Based Bioengineering System for Predicting and Diagnosing Diseases of the Nervous System Triggered by the Interaction of Industrial Frequency Electromagnetic Fields. Crit Rev Biomed Eng. 2024;52(5):1-16. doi: 10.1615/CritRevBiomedEng.2024053240.

Abstract

The study aims to enhance the standard of medical care for individuals working in the electric power industry who are exposed to industrial frequency electromagnetic fields and other relevant risk factors. This enhancement is sought through the integration of fuzzy mathematical models with contemporary information and intellectual technologies. The study addresses the challenges of forecasting and diagnosing illnesses within a specific demographic characterized by a combination of poorly formalized issues with interconnected conditions. To tackle this complexity, a methodological framework was developed for synthesizing hybrid fuzzy decision rules. This approach combines clinical expertise with artificial intelligence methodologies to promote innovative problem-solving strategies. Additionally, the researchers devised an original method to evaluate the body's protective capacity, which was integrated into these decision rules to enhance the precision and efficacy of medical decision-making processes. The research findings indicate that industrial frequency electromagnetic fields contribute to illnesses of societal significance. Additionally, it highlights that these effects are worsened by other risk factors such as adverse microclimates, noise, vibration, chemical exposure, and psychological stress. Diseases of the neurological, immunological, cardiovascular, genitourinary, respiratory, and digestive systems are caused by these variables in conjunction with unique physical traits. The development of mathematical models in this study makes it possible to detect and diagnose disorders in workers exposed to electromagnetic fields early on, especially those pertaining to the autonomic nervous system and heart rhythm regulation. The results can be used in clinical practice to treat personnel in the electric power industry since expert evaluation and modeling showed high confidence levels in decision-making accuracy.

https://pubmed.ncbi.nlm.nih.gov/38884210/

--

Exposure assessment and cytogenetic biomonitoring study of workers occupationally exposed to extremely low-frequency magnetic fields

Nguyen H, Vandewalle G, Mertens B, Collard JF, Hinsenkamp M, Verschaeve L, Feipel V, Magne I, Souques M, Beauvois V, Ledent M. Exposure assessment and cytogenetic biomonitoring study of workers occupationally exposed to extremely low-frequency magnetic fields. Bioelectromagnetics. 2024 Jun 11. doi: 10.1002/bem.22506.

Abstract

Human cytogenetic biomonitoring (HCB) has long been used to evaluate the potential effects of work environments on the DNA integrity of workers. However, HCB studies on the genotoxic effects of occupational exposure to extremely low-frequency electromagnetic fields (ELF-MFs) were limited by the quality of the exposure assessment. More specifically, concerns were raised regarding the method of exposure assessment, the selection of exposure metrics, and the definition of exposure group. In this study, genotoxic effects of occupational exposure to ELF-MFs were assessed on peripheral blood lymphocytes of 88 workers from the electrical sector using the comet and cytokinesis-block micronucleus assay, considering workers' actual exposure over three consecutive days. Different methods were applied to define exposure groups. Overall, the summarized ELF-MF data indicated a low exposure level in the whole study population. It also showed that relying solely on job titles might misclassify 12 workers into exposure groups. We proposed combining hierarchical agglomerative clustering on personal exposure data and job titles to define exposure groups. The final results showed

that occupational MF exposure did not significantly induce more genetic damage. Other factors such as age or past smoking rather than ELF-MF exposure could affect the cytogenetic test outcomes.

Highlights

- The study results indicated that exposure assessment had better rely on the workers' actual exposure data.
- Electrical workers in this study were exposed on average to quite low levels of 50 Hz magnetic fields at work. But there was wide variability between workers. This highlights the value of using cluster analysis on workers' actual exposure data to define exposure groups.
- Occupational exposure to 50 Hz magnetic fields did not significantly increase the level of DNA or chromosomal damage on workers' blood samples.

Open access paper: https://pubmed.ncbi.nlm.nih.gov/38862415/

--

Thoughts on the entanglement of electromagnetism and life: A theoretical study

Zsarnovszky A. Thoughts on the entanglement of electromagnetism and life: A theoretical study. Acta Vet Hung. 2024 Jun 18. doi: 10.1556/004.2023.00976. Epub ahead of print. PMID: 38888962.

Abstract

Dissection of the matter into its constituents leads us to the smallest particles that we know. These particles form a material structure that is determined by the electromagnetic field generated and carried by those particles. Changes in any of the two major constituents leads to changes in that material system, be it a living organism or a lifeless object. The latter statement carries the mystery of life that is born from a continuous and programmed series of system changes fuelled by an energy source with a yet unknown functioning mechanism. The present work is a theoretical approach towards the understanding and potential discovery of the aforementioned, not-yet-known cellular energetic mechanism. Understanding the energetic basis of intracellular biochemistry is equally important in human and animal therapeutics. Additionally, as all such discoveries offer novel solutions in various fields of the global industry, the final outcome of this theoretical work also brings about the idea of a new discovery in electronics industry.

Open access paper: https://akjournals.com/view/journals/004/aop/article-10.1556-004.2023.00976/article-10.1556-004.2023.00976.xml

--

Investigation of genetic stress parameters in brain tissues of rats exposed to 1.8 GHz cell phone radiofrequency electromagnetic field

Arslan B, Aras N, Yaman S, Comelekoglu U. Investigation of genetic stress parameters in brain tissues of rats exposed to 1.8 GHz cell phone radiofrequency electromagnetic field. Med Science. 2024;13(1):78-82.

Abstract

Heat Shock Proteins (HSPs) may induce various cellular processes, including replication, apoptosis, cell-cycle progression. Mitogen-activated protein kinase (MAPK) cascades are the primary mechanism that mediates the cellular stress response to extracellular stimuli and regulates transcriptional activity. It has been shown that mobile phone exposure can stimulate the Hsp27/p38MAPK stress pathway. In this study, twenty-seven mature female Wistar albino rats were exposed to 1.8 GHz radiofrequency electromagnetic field (RF-EMF) 2h/day for 8 weeks (SAR: 0.06 W/kg). Hsp27 and p38MAPK gene expressions were investigated in rat brains. Rats were divided into groups sham-exposed, cage control, and 1.8 GHz RF-EMF exposed. Hsp27 and p38MAPK gene expression levels were investigated from the brain. p38MAPK expression was found to be upregulated in RF-EMF exposed group (p=0.018) Hsp27 expressions were not altered (p=0.897). In conclusion, long-term exposure to 1.8 GHz cell phone radiation can activate the Hsp27/p38MAPK stress pathway. It may cause several cellular disorders and can affect brain function.

Open access paper: https://www.medicinescience.org/article/4261

--

Critical role of model organism selection in assessing weak urban electromagnetic field effects: Implications for human health

Sincak M, Adamkova P, Demeckova V, Smelko M, Lipovsky P, Oravec M, Luptakova A, Sedlakova-Kadukova J. Critical role of model organism selection in assessing weak urban electromagnetic field effects: Implications for human health. Bioelectrochemistry. 2024 Jun 29;160:108756. doi: 10.1016/j.bioelechem.2024.108756.

Highlights

- Real urban electromagnetic field present a combination 50 and 150 Hz in ratio 4:1.
- Macrophages shifted from M2 to M1 phenotype in 2 h in urban magnetic field.
- Urban magnetic fields can induce pro-inflammatory macrophage phenotype.
- Yeast showed only one change: 17 h exponential growth phase extension.
- Is crucial to choose proper organism for study of urban magnetic field perception.

Abstract

The impact of electromagnetic fields on human health has been investigated in recent years using various model organisms, yet the findings remain unclear. In our work, we examined the effect of less-explored, weak electromagnetic fields commonly found in the urban environments we inhabit. We studied different impacts of electromagnetic fields with a frequency of 50 Hz and a combination of 50 Hz and 150 Hz, on both yeasts (Saccharomyces cerevisiae) and human macrophages. We determined growth, survival, and protein composition (SDS-PAGE) (Saccharomyces cerevisiae) and morphology of macrophages (human monocytic cell line). In yeast, the sole observed change after 24 h of exposure was the extension of the exponential growth phase by 17 h. Conversely, macrophages exhibited morphological transformations from the anti-inflammatory to the pro-inflammatory type within just 2 h of exposure to the electromagnetic field. Our results suggest that effects of electromagnetic field largely depend on the model organism. The selection of an appropriate model organism proves essential for the study of the specific impacts of electromagnetic fields. The potential risk associated with the presence of

pro-inflammatory M1 macrophages in everyday urban environments primarily arises from the continual promotion of inflammatory reactions within a healthy organism and deserves further investigation.

https://pubmed.ncbi.nlm.nih.gov/38959750/

Conclusion

Based on our observations, macrophages appear to be more responsive to the presence of an electromagnetic field, making them a more proper model organism for describing the influence of the urban electromagnetic environment on humans compared to yeasts. Our results demonstrate that while yeasts did not exhibit changes in growth, pH, and redox potential within 24 h under low-intensity fields (20 µT and 50 or 50/150 Hz), immune system cells detected a magnetic field alteration within just 2 h, manifested in a shift in their phenotype.

Though these changes were temporary, the transformation of macrophages into pro-inflammatory M1 macrophages raises concerns about the potential risks associated with prolonged exposure to the weak urban electromagnetic fields that surround us daily. Continuous exposure to electromagnetic fields in our daily lives may have adverse effects on our health, as demonstrated through the use of macrophages as model organisms.

--

On the origin of the biological effects of time varying magnetic fields: quantitative insights

Ravikumar K, Roy Choudury AN, Dubey AK, Kumaran V, Basu B. On the origin of the biological effects of time varying magnetic fields: quantitative insights. J Mater Chem B. 2024 Jul 3. doi:10.1039/d4tb00362d.

Abstract

In a number of recently published experimental studies from our research group, the positive impact of magnetic stimuli (static/pulsed) on cell functionality modulation or bactericidal effects, *in vitro*, has been established. In order to develop a theoretical understanding of such magnetobiological effects, the present study aimed to present two quantitative models to determine magnetic Maxwell stresses as well as pressure acting on the cell membrane, under the influence of a time varying magnetic field. The model predicts that magnetic field-induced stress on the cell/bacteria is dependent on the conductivity properties of the extracellular region, which is determined to be too low to cause any significant effect. However, the force on the cell/bacteria due to the induced electric field is more influential than that of the magnetic field, which has been used to determine the membrane tension that can cause membrane poration. With a known critical membrane tension for cells, the field parameters necessary to cause membrane rupture have been estimated. Based on the experimental results and theoretically predicted values, the field parameters can be classified into three regimes, wherein the magnetic fields cause no effect or result in biophysical stimulation or induce cell death due to membrane damage. Taken together, this work provides some quantitative insights into the impact of magnetic fields on biological systems.

https://pubmed.ncbi.nlm.nih.gov/38958687/

--

Do non-ionizing radiation concerns affect people's choice between hybrid and traditional cars?

Anat Tchetchik, Sigal Kaplan, Orit Rotem-Mindali O. Do non-ionizing radiation concerns affect people's choice between hybrid and traditional cars? Transportation Research Part D: Transport and Environment, Volume 131, 2024, doi: 10.1016/j.trd.2024.104226.

Abstract

The growing market for hybrid electric vehicles (HEV) has raised concerns about the long-term impacts of non-ionizing radiation (NIR) exposure. This study is the first to address the impact of NIR on consumer choice between HEV and internal combustion engine (ICE) vehicles. We explore the hypothesis that NIR is associated with a lower probability of HEV choice in the presence of NIR information and the relative effect of NIR-health concerns versus environmental attitudes and driving norms. The data are collected from a stated choice experiment and estimated via a hybrid choice model. The results show that i NIR is associated with a lower choice probability of HEV, ii NIR-dread is associated with a higher probability of choosing ICE vehicles, while skepticism about NIR is associated with a higher probability of choosing HEV, iii prompting positively or negatively framed information about NIR discourages HEV choice compared to providing no information.

Conclusions and policy recommendations

The results show the effect of NIR-associated barriers on the choice of HEV versus ICE and highlight the following policy recommendations.

First, the massive production of EVs combined with the lack of regulatory frameworks can lead to the introduction of low-cost car models with low NIR safety standards (Trentadue et al., 2020). The European Union recommends a clear regulatory framework and international standards to promote the transition toward EVs. This study showed that NIR levels negatively affect the choice of HEV, signaling to car manufacturers and policymakers that consumers are concerned about NIR levels. Accordingly, setting NIR safety standards and maintaining low NIR levels are important goals for the transition toward autonomous, connected, electric vehicles.

Second, this study showed that while NIR dread was a discouraging factor, NIR skepticism was a strong choice motivator. Thus, perceived occurrence probability is as important as NIR risk dread. As with other health issues, prevalence across the population is an important decision-making factor that, in the absence of information, may lead to self-exemption beliefs. Scientific evidence from large-scale studies regarding both short- and long-term NIR effects and their prevalence in the population and among risk groups will enable informed decision-making, help mitigate NIR dread, and establish meaningful guidelines for in-vehicle NIR levels. With climate goals requiring the transition toward EV by 2030 and with the rapid technological advancement of autonomous, connected, electric vehicles, establishing the prevalence of NIR short- and long-term health effects is important for the future of the industry.

Third, better information quality strengthens the relationship between the depiction of new vehicle technologies and perceived purchase value (Zhang et al., 2022). Our study showed that both positive and negative framing can lead to a lower choice probability when an NIR safety threshold is provided. In this study, the information that “Studies show that long-term exposure to NIR levels below 4 mG is safe”

was associated with lower choice probabilities, similar to the case of negative framing, “Studies show that long-term exposure to NIR levels below 4 mG increases the health risks to health concerns.” Policymakers and manufacturers must consider information quality in terms of accuracy, clarity, ambiguity, and potential sources of confusion and decision bias. In this study, consumers used the provided threshold of 4 mG as a decision anchor, which means that consumers in some cultural contexts seek clear, “fast and frugal” evaluation criteria without engaging in complex exposure evaluations.

Finally, the model shows that travel with children is negatively associated with HEV leasing. Nevertheless, while NIR dread is negatively associated with HEV leasing, an additional interaction effect between NIR levels and travel with children was not statistically significant. These results indicate that while NIR dread is important, there is no additional health concerns particularly asso- ciated with travel with children. Hence, the decrease in the HEV leasing propensity when traveling with children may be associated with other reasons, such as vehicle reliability or other concerns that were not investigated in the current study. Notably, previous studies found a particular concern for children’s health-related to NIR from mobile phones and cellular stations. Leach and Bromwich (2018) found that two-thirds of the participants believed that mobile technology use should be restricted due to possible health risks to children’s health. P¨olzl (2011) added that 30 % of the population had strong or considerable concerns regarding NIR health risks to children, and noted that adults can be motivated to adjust their behavior to protect their children. Further research is important in other regions and contexts, to understand more thoroughly the issue of HEV leasing or purchase when traveling with children.

https://www.sciencedirect.com/science/article/pii/S1361920924001834

--

50 Hz magnetic field influences caspase-3 activity and cell cycle distribution in ionizing radiation exposed SH-SY5Y neuroblastoma cells

Nieminen V, Martikainen MV, Kalliomäki S, Virén T, Seppälä J, Juutilainen J, Naarala J, Luukkonen J. 50 Hz magnetic field influences caspase-3 activity and cell cycle distribution in ionizing radiation exposed SH-SY5Y neuroblastoma cells. Int J Radiat Biol. 2024 Jun 26:1-10. doi: 10.1080/09553002.2024.2369105.

Abstract

Purpose: Earlier evidence suggests that extremely low frequency magnetic fields (ELF MFs) can modify the effects of carcinogenic agents. However, the studies conducted so far with ionizing radiation as the co-exposure agent are sparse and have provided inconclusive results. We investigated whether 50 Hz MFs alone, or in combination with ionizing radiation alter cell biological variables relevant to cancer and the biological effects of ionizing radiation.

Materials and methods: Human SH-SY5Y neuroblastoma cells were sham exposed or exposed to 100 or 500 µT MF for 24 h either before or after ionizing radiation exposure (0, 0.4 or 2 Gy). After the exposures, cells were assayed for viability, clonogenicity, reactive oxygen species, caspase-3 activity, and cell cycle distribution. Cell cycle distribution was assayed with propidium iodide staining followed by flow cytometry analysis and ROS levels were assayed together with cell viability by double staining with DeepRed and Sytox Blue followed by flow cytometry analysis.

Results: Increased caspase-3 activity was observed in cells exposed to 500 µT MF before or after ionizing radiation. Furthermore, exposure to the 500 µT MF after the ionizing radiation decreased the percentage of cells in S-phase. No changes in the ROS levels, clonogenicity, or viability of the cells were observed in the MF exposed groups compared to the corresponding sham exposed groups, and no MF effects were observed in cells exposed at 100 µT.

Conclusions: Only the 500 µT magnetic flux density affected SH-SY5Y cells significantly. The effects were small but may nevertheless help to understand how MFs modify the effects of ionizing radiation. The increase in caspase-3 activity may not reflect effects on apoptosis, as no changes were observed in the subG1 phase of the cell cycle. In contrast to some earlier findings, 50 Hz MF exposure after ionizing radiation was not less effective than MF treatment given prior to ionizing radiation.

https://pubmed.ncbi.nlm.nih.gov/38924721/

--

Electromagnetic field-induced adaptive response in Schwann cells through DNA methylation, histone deacetylation, and oxidative stress

Colciago A, Mohamed T, Colleoni D, Melfi V, Magnaghi V. Electromagnetic field-induced adaptive response in Schwann cells through DNA methylation, histone deacetylation, and oxidative stress. J Cell Physiol. 2024 Jun 30:e31365. doi: 10.1002/jcp.31365.

Abstract

Schwannomas are benign tumors of the peripheral nervous system arising from the transformation of Schwann cells (SCs). On the whole, these tumors are related to alterations of the neurofibromin type 2 gene, coding for the oncosuppressor merlin, a cytoskeleton-associated protein belonging to the ezrin-radixin-moesin family. However, the underlying mechanisms of schwannoma onset and progression are not fully elucidated, whereas one of the challenges might be the environment. In this light, the exposure to electromagnetic field (EMF), generated by the use of common electrical devices, has been defiantly suggested as the cause of SCs transformation even if the evidence was mostly epidemiologic. Indeed, insubstantial mechanisms have been so far identified to explain SCs oncotransformation. Recently, some in vitro evidence pointed out alterations in proliferation and migration abilities in SCs exposed to EMF (0.1 T, 50 Hz, 10 min). Here, we used the same experimental paradigma to discuss the involvement of putative epigenetic mechanisms in SCs adaptation to EMF and to explain the occurrence of hypoxic alterations after the exposure. Our findings indicate a set of environmental-induced changes in SCs, toward a less-physiological state, which may be pathologically relevant for the SCs differentiation and the schwannoma development.

https://pubmed.ncbi.nlm.nih.gov/38946084/

--

The Slowest Shared Resonance: A Review of Electromagnetic Field Oscillations Between Central and Peripheral Nervous Systems

Young A, Hunt T, Ericson M. The Slowest Shared Resonance: A Review of Electromagnetic Field Oscillations Between Central and Peripheral Nervous Systems. Front Hum Neurosci. 2022 Feb 16;15:796455. doi: 10.3389/fnhum.2021.796455.

Abstract

Electromagnetic field oscillations produced by the brain are increasingly being viewed as causal drivers of consciousness. Recent research has highlighted the importance of the body's various endogenous rhythms in organizing these brain-generated fields through various types of entrainment. We expand this approach by examining evidence of extracerebral shared oscillations between the brain and other parts of the body, in both humans and animals. We then examine the degree to which these data support one of General Resonance Theory's (GRT) principles: the Slowest Shared Resonance (SSR) principle, which states that the combination of micro- to macro-consciousness in coupled field systems is a function of the slowest common denominator frequency or resonance. This principle may be utilized to develop a spatiotemporal hierarchy of brain-body shared resonance systems. It is predicted that a system's SSR decreases with distance between the brain and various resonating structures in the body. The various resonance relationships examined, including between the brain and gastric neurons, brain and sensory organs, and brain and spinal cord, generally match the predicted SSR relationships, empirically supporting this principle of GRT.

Open access paper:
https://www.frontiersin.org/journals/human-neuroscience/articles/10.3389/fnhum.2021.796455/full

--

Brain structure alterations following neonatal exposure to low-frequency electromagnetic fields: A histological analysis

Sissons SM, Dotta BT. Brain structure alterations following neonatal exposure to low-frequency electromagnetic fields: A histological analysis. Int J Dev Neurosci. 2024 Jul 5. doi: 10.1002/jdn.10361.

Abstract

Nitric oxide (NO) and electromagnetic fields (EMF) have been extensively studied for their roles in neurobiology, particularly in regulating cerebral functions and synaptic plasticity. This study investigates the impact of EMFs on NO modulation and its subsequent effects on neurodevelopment, building upon prior research examining EMF exposure's consequences on Wistar albino rats. Rats were exposed perinatally to either tap water, 1 g/L of L-arginine (LA) or 0.5 g/L of N-methylarginine (NMA). Half of the rats in each group were also exposed to a 7-Hz square-wave EMF at three separate intensities (5, 50 and 500 nT) for 2-14 days following birth. Animals were allowed to develop, and their brains were harvested later in adulthood (mean age = 568.17 days, SD = 162.73). Histological analyses were used to elucidate structural changes in key brain regions. All brains were stained with Toluidine Blue O (TBO), enabling the visualization of neurons. Neuronal counts were then conducted in specific regions of interest (e.g. hippocampus, cortices, amygdala and hypothalamus). Histological analyses revealed significant alterations in neuronal density in specific brain regions, particularly in response to EMF exposure and pharmacological interventions. Notable findings include a main EMF exposure effect where increased neuronal counts were observed in the secondary somatosensory cortex under low EMF intensities ($p <$

0.001) and sex-specific responses in the hippocampus, where a significant increase in neuronal counts was observed in the left CA3 region in female rats exposed to EMF compared to unexposed females ($t(18) = 2.371$, $p = 0.029$). Additionally, a significant increase in neuronal counts in the right entorhinal cortex was seen in male rats exposed to EMF compared to unexposed males ($t(18) = 2.216$, $p = 0.040$). These findings emphasize the complex interaction among sex, EMF exposure and pharmacological agents on neuronal dynamics across brain regions, highlighting the need for further research to identify underlying mechanisms and potential implications for cognitive function and neurological health in clinical and environmental contexts.

https://pubmed.ncbi.nlm.nih.gov/38967459/

--

Electromagnetic Exposure Levels of Electric Vehicle Drive Motors to Passenger Wearing Cardiac Pacemakers

Dong X, Qian Y, Lu M. Electromagnetic Exposure Levels of Electric Vehicle Drive Motors to Passenger Wearing Cardiac Pacemakers. *Sensors*. 2024; 24(13):4395. doi: 10.3390/s24134395

Abstract

The number of individuals wearing cardiac pacemakers is gradually increasing as the population ages and cardiovascular disease becomes highly prevalent. The safety of pacemaker wearers is of significant concern because they must ensure that the device properly functions in various life scenarios. Electric vehicles have become one of the most frequently used travel tools due to the gradual promotion of low-carbon travel policies in various countries. The electromagnetic environment inside the vehicle is highly complex during driving due to the integration of numerous high-power electrical devices inside the vehicle. In order to ensure the safety of this group, the paper takes passengers wearing cardiac pacemakers as the object and the electric vehicle drive motors as the exposure source. Calculation models, with the vehicle body, human body, heart, and cardiac pacemaker, are built. The induced electric field, specific absorption rate, and temperature changes in the passenger's body and heart are calculated by using the finite element method. Results show that the maximum value of the induced electric field of the passenger occurs at the ankle of the body, which is 60.3 mV/m. The value of the induced electric field of the heart is greater than that of the human trunk, and the maximum value (283 mV/m) is around the pacemaker electrode. The maximum specific absorption rate of the human body is 1.08×10^{-6} W/kg, and that of heart positioned near the electrode is 2.76×10^{-5} W/kg. In addition, the maximum temperature increases of the human torso, heart, and pacemaker are 0.16×10^{-5} °C, 0.4×10^{-6} °C, and 0.44×10^{-6} °C within 30 min, respectively. Accordingly, the induced electric field, specific absorption rate, and temperature rise in the human body and heart are less than the safety limits specified in the ICNIRP. The electric field intensity at the pacemaker electrode and the temperature rise of the pacemaker meet the requirements of the medical device standards of ICNIRP and ISO 14708-2. Consequently, the electromagnetic radiation from the motor operation in the electric vehicle does not pose a safety risk to the health of passengers wearing cardiac pacemakers in this paper. This study also contributes to advancing research on the electromagnetic environment of electric vehicles and provides guidance for ensuring the safe travel of individuals wearing cardiac pacemakers.

Open access paper: https://www.mdpi.com/1424-8220/24/13/4395

--

Personal light exposure patterns and incidence of type 2 diabetes: analysis of 13 million hours of light sensor data and 670,000 person-years of prospective observation

Daniel P. Windred, Angus C. Burns, Martin K. Rutter, Chris Ho Ching Yeung, Jacqueline M. Lane, Qian Xiao, Richa Saxena, Sean W. Cain, Andrew J. K. Phillips. Personal light exposure patterns and incidence of type 2 diabetes: analysis of 13 million hours of light sensor data and 670,000 person-years of prospective observation. Lancet. 2024. doi: 10.1016/j.lanepe.2024.100943.

Abstract

Background Light at night disrupts circadian rhythms, and circadian disruption is a risk factor for type 2 diabetes. Whether personal light exposure predicts diabetes risk has not been demonstrated in a large prospective cohort. We therefore assessed whether personal light exposure patterns predicted risk of incident type 2 diabetes in UK Biobank participants, using ∼13 million hours of light sensor data.

Methods Participants (N = 84,790, age (M ± SD) = 62.3 ± 7.9 years, 58% female) wore light sensors for one week, recording day and night light exposure. Circadian amplitude and phase were modeled from weekly light data. Incident type 2 diabetes was recorded (1997 cases; 7.9 ± 1.2 years follow-up; excluding diabetes cases prior to light-tracking). Risk of incident type 2 diabetes was assessed as a function of day and night light, circadian phase, and circadian amplitude, adjusting for age, sex, ethnicity, socioeconomic and lifestyle factors, and polygenic risk.

Findings Compared to people with dark nights (0–50th percentiles), diabetes risk was incrementally higher across brighter night light exposure percentiles (50–70th: multivariable-adjusted HR = 1.29 [1.14–1.46]; 70–90th: 1.39 [1.24–1.57]; and 90–100th: 1.53 [1.32–1.77]). Diabetes risk was higher in people with lower modeled circadian amplitude (aHR = 1.07 [1.03–1.10] per SD), and with early or late circadian phase (aHR range: 1.06–1.26). Night light and polygenic risk independently predicted higher diabetes risk. The difference in diabetes risk between people with bright and dark nights was similar to the difference between people with low and moderate genetic risk.
Interpretation Type 2 diabetes risk was higher in people exposed to brighter night light, and in people exposed to light patterns that may disrupt circadian rhythms. Avoidance of light at night could be a simple and cost-effective recommendation that mitigates risk of diabetes, even in those with high genetic risk.

Open access paper: https://www.thelancet.com/pdfs/journals/lanepe/PIIS2666-7762(24)00110-8.pdf

--

IEC/IEEE Draft International Standard - Assessment methods of the human exposure to electric and magnetic fields from wireless power transfer systems

IEC/IEEE Draft International Standard - Assessment methods of the human exposure to electric and magnetic fields from wireless power transfer systems - models, instrumentation, measurement and numerical methods and procedures (Frequency range of 1 kHz to 30 MHz). *IEC/IEEE P63184/FDIS, June 2024* , vol., no., pp.1-153, 26 June 2024.

Abstract

The objective of this document is to specify the assessment methods to evaluate compliance of stationary and dynamic wireless power transfer (WPT) systems with electromagnetic human exposure guidelines (specific absorption rate (SAR), internal electric fields, or current density including contact currents). The frequency range covered by this document is from 1 kHz to 30 MHz. This version of the standard focuses on exposures from inductive WPT systems. Future versions will consider extended guidance for assessments of exposure from capacitive WPT systems.

https://ieeexplore.ieee.org/document/10576682

--

Evaluation of oxidative stress and genetic instability among residents near mobile phone base stations in Germany

Gulati S, Mosgoeller W, Moldan D, Kosik P, Durdik M, Jakl L, Skorvaga M, Markova E, Kochanova D, Vigasova K, Belyaev I. Evaluation of oxidative stress and genetic instability among residents near mobile phone base stations in Germany. Ecotoxicology and Environmental Safety, Volume 279, 2024, doi: 10.1016/j.ecoenv.2024.116486.

Highlights

- Residents exposed to RF-EMR from base stations were tested for genetic instability.
- The years long exposure had no measurable effect on specific cancer related genes.
- We found high level of lipid peroxidation and DNA-lesions but not significant.
- Chromosomal aberrations were significantly more pronounced in highly exposed group.

Abstract

Human exposure to radiofrequency electromagnetic fields (RF-EMF) is restricted to prevent thermal effects in the tissue. However, at very low intensity exposure "non-thermal" biological effects, like oxidative stress, DNA or chromosomal aberrations, etc. collectively termed genomic-instability can occur after few hours. Little is known about chronic (years long) exposure with non-thermal RF-EMF. We identified two neighboring housing estates in a rural region with residents exposed to either relatively low (control-group) or relatively high (exposed-group) RF-EMF emitted from nearby mobile phone base stations (MPBS). 24 healthy adults that lived in their homes at least for 5 years volunteered. The homes were surveyed for common types of EMF, blood samples were tested for oxidative status, transient DNA alterations, permanent chromosomal damage, and specific cancer related genetic markers, like MLL gene rearrangements. We documented possible confounders, like age, sex, nutrition, life-exposure to ionizing radiation (X-rays), occupational exposures, etc. The groups matched well, age, sex, lifestyle and occupational risk factors were similar. The years long exposure had no measurable effect on MLL gene rearrangements and c-Abl-gene transcription modification. Associated with higher exposure, we found higher levels of lipid oxidation and oxidative DNA-lesions, though not statistically significant. DNA double strand breaks, micronuclei, ring chromosomes, and acentric chromosomes were not significantly different between the groups. Chromosomal aberrations like dicentric chromosomes ($p=0.007$), chromatid gaps ($p=0.019$), chromosomal fragments ($p<0.001$) and the total of chromosomal aberrations ($p<0.001$) were significantly higher in the exposed group. No potential confounder interfered with these

findings. Increased rates of chromosomal aberrations as linked to excess exposure with ionizing radiation may also occur with non-ionizing radiation exposure. Biological endpoints can be informative for designing exposure limitation strategies. Further research is warranted to investigate the dose-effect-relationship between both, exposure intensity and exposure time, to account for endpoint accumulations after years of exposure. As established for ionizing radiation, chromosomal aberrations could contribute to the definition of protection thresholds, as their rate reflects exposure intensity and exposure time.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0147651324005621

--

Analysis of the metabolic profile of humans naturally exposed to RF-EM radiation

Rangesh NM, Malaisamy AK, Kumar N, Kumar S. Analysis of the metabolic profile of humans naturally exposed to RF-EM radiation. Metabolomics. 2024 May 18;20(3):55. doi: 10.1007/s11306-024-02121-2.

Abstract

Introduction: The world is experiencing exponential growth in communication, especially wireless communication. Wireless connectivity has recently become a part of everyone's daily life. Recent developments in low-cost, low-power, and miniature devices contribute to a significant rise in radiofrequency-electromagnetic field (RF-EM) radiation exposure in our environment, raising concern over its effect on biological systems. The inconsistent and conflicting research results make it difficult to draw definite conclusions about how RF-EM radiation affects living things.

Objectives: This study identified two micro-environments based on their level of exposure to cellular RF-EM radiation, one with significantly less exposure and another with very high exposure to RF-EM radiation. Emphasis is given to studying the metabolites in the urine samples of humans naturally exposed to these two different microenvironments to understand short-term metabolic dysregulations.

Methods: Untargeted 1H NMR spectroscopy was employed for metabolomics analyses to identify dysregulated metabolites. A total of 60 subjects were recruited with 5 ml urine samples each. These subjects were divided into two groups: one highly exposed to RF-EM (n = 30) and the other consisting of low-exposure populations (n = 30).

Results: The study found that the twenty-nine metabolites were dysregulated. Among them, 19 were downregulated, and 10 were upregulated. In particular, Glyoxylate and dicarboxylate and the TCA cycle metabolism pathway have been perturbed. The dysregulated metabolites were validated using the ROC curve analysis.

Conclusion: Untargeted urine metabolomics was conducted to identify dysregulated metabolites linked to RF-EM radiation exposure. Preliminary findings suggest a connection between oxidative stress and gut microbiota imbalance. However, further research is needed to validate these biomarkers and understand the effects of RF-EM radiation on human health. Further research is needed with a diverse population.

Conclusion

The purpose of this study was to initiate an exploration using urine metabolomics and ^{1}H NMR spectroscopy to identify specific dysregulated metabolites associated with RF-EM radiation exposure. The objective was to pinpoint potential biomarkers for future validation. Based on these preliminary findings, there appears to be a possible connection between RF-EM radiation exposure, oxidative stress, and disruptions in the gut microbiota balance. Nevertheless, it's imperative to stress that additional investigations with a larger and more diverse population are necessary to validate these identified biomarkers and gain a more comprehensive understanding of the effects of RF-EM radiation exposure on human health. Furthermore, individuals with lower levels of radiation exposure exhibited metabolite signatures of a healthier gut microbiota system, indicating a potential association between radiation exposure and microbiota health. However, it is crucial to emphasize that further research is essential for confirmation. Future studies involve conducting a comprehensive study involving individuals with substantial RF-EM radiation exposure alongside a control group of unexposed individuals.

https://pubmed.ncbi.nlm.nih.gov/38762651/

--

Investigation of the Adverse Health Effects of Cell Phone Radiation and Propose Solutions to Minimize Them: A Systematic Review

Elyasi H, Ghanbari M, Nadri F. Investigation of the Adverse Health Effects of Cell Phone Radiation and Propose Solutions to Minimize Them: A Systematic Review. Indian J Occup Environ Med. 2024 Jan-Mar;28(1):18-22. doi: 10.4103/ijoem.ijoem_89_23.

Abstract

Today, mobile phones are one of the most common devices emitting electromagnetic radiation and are available to more than seven billion people in different age groups around the world. The effects of electromagnetic radiation on biological systems have been studied for several years. In this systematic review to find relevant articles, international databases such as Google Scholar, PubMed, Scopus, Science Direct, Web of Science, and Cochrane were used since 2007-2022 by selecting appropriate keywords. The result revealed that exposure to cell phone radiation can lead to disturb in the metabolic activity of the cerebellum by increasing the migration of granulosa cells, decrease the water around the fetus in pregnant women, decrease in the number of blood plates, increasing levels of ALT and AST that they are the key biomarkers of liver damage, decrease of phagocytosis and induced apoptosis of neutrophils, changes at the level of glucose and even at the microscopic level of pancreas this may be a predisposing factor for diabetes, increment in tissue temperature in all depth of the brain tissue, EMF increase the volume, weight, and atresia follicles of the ovaries of the children, also it can cause oxidative stress, DNA fragmentation, etc. Mobile phone radiation is harmful and depends on its intensity, frequency, wave type, and exposure duration. It can cause different biological effects in humans. Due to the uncertainty of the results and mechanism of the effect of these waves, research in this field is still ongoing.

Open access paper: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11111148/

--

Evaluation of neonatal outcomes according to the specific absorption rate values of phones used during pregnancy

Büyükeren M, Karanfil Yaman F. Evaluation of neonatal outcomes according to the specific absorption rate values of phones used during pregnancy. J Turk Ger Gynecol Assoc. 2024 Mar 6;25(1):7-12. doi: 10.4274/jtgga.galenos.2023.2022-10-1.

Abstract

Objective: The aim was to compare neonatal outcomes according to cell phone specific absorption rate (SAR) levels and daily time spent on cell phones by pregnant women.

Material and Methods: Women who gave birth at Konya City Hospital between September 2020 and February 2021 were included in this retrospective study. Gestational ages, birth weight, birth length, head circumference, sex, 5-minute APGAR scores, neonate postpartum resuscitation requirement, delivery type, the model of phone used by the pregnant women, and the average time spent on the phone during a day were recorded. To determine the relation between the SAR values of the phones used and delivering a small for gestational age (SGA) baby, receiver operating characteristic curve analysis was performed.

Results: In total 1495 pregnant women were included. The rate of delivering a SGA fetus was significantly higher in women who used phones with higher SAR values (p=0.001). The cut-off value for the SAR level was 1.23 W/kg with 69.3% sensitivity and 73.0% specificity (area under the curve: 0.685; 95% confidence interval: 0.643-0.726). No correlation was found between time spent on the phone and SGA birth rate. Although both phone SAR values and time spent on the phone were higher in the symmetrical SGA group compared to the asymmetrical SGA group, the difference was not significant (p>0.05). Although the women who had preterm delivery had higher phone SAR values and had spent more time on the phone compared to those who had term deliveries, the difference was again not significant (p>0.05).

Conclusion: As the SAR values of cell phones used during pregnancy increased, there was a trend towards delivering a SGA baby.

Open access paper: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10921072/

--

1.7 GHz LTE radiofrequency EMF with stable power monitoring and efficient thermal control has no effect on the proliferation of various human cell types

My note: The study employed a Western blot analysis which is less likely to detect DNA damage than a comet assay.

Goh J, Suh D, Park G, Jeon S, Lee Y, Kim N, Song K. 1.7 GHz long-term evolution radiofrequency electromagnetic field with stable power monitoring and efficient thermal control has no effect on the proliferation of various human cell types. PLoS One. 2024 May 7;19(5):e0302936. doi: 10.1371/journal.pone.0302936.

Abstract

Long-term evolution (LTE) radiofrequency electromagnetic field (RF-EMF) is widely used in communication technologies. Thus, the influence of RF-EMF on biological systems is a major public concern and its physiological effects remain controversial. In our previous study, we showed that continuous exposure of various human cell types to 1.7 GHz LTE RF-EMF at a specific absorption rate (SAR) of 2 W/Kg for 72 h can induce cellular senescence. To understand the precise cellular effects of LTE RF-EMF, we elaborated the 1.7 GHz RF-EMF cell exposure system used in the previous study by replacing the RF signal generator and developing a software-based feedback system to improve the exposure power stability. This refinement of the 1.7 GHz LTE RF-EMF generator facilitated the automatic regulation of RF-EMF exposure, maintaining target power levels within a 3% range and a constant temperature even during the 72-h-exposure period. With the improved experimental setup, we examined the effect of continuous exposure to 1.7 GHz LTE RF-EMF at up to SAR of 8 W/Kg in human adipose tissue-derived stem cells (ASCs), Huh7, HeLa, and rat B103 cells. Surprisingly, the proliferation of all cell types, which displayed different growth rates, did not change significantly compared with that of the unexposed controls. Also, neither DNA damage nor cell cycle perturbation was observed in the 1.7 GHz LTE RF-EMF-exposed cells. However, when the thermal control system was turned off and the subsequent temperature increase induced by the RF-EMF was not controlled during continuous exposure to SAR of 8 W/Kg LTE RF-EMF, cellular proliferation increased by 35.2% at the maximum. These observations strongly suggest that the cellular effects attributed to 1.7 GHz LTE RF-EMF exposure are primarily due to the induced thermal changes rather than the RF-EMF exposure itself.

Conclusion

To understand the precise cellular effect of 1.7 GHz LTE RF-EMF, we developed an RF-EMF cell exposure system with an improved RF signal generator and control software. With a refined RF-EMF exposure system, we could maintain a consistent target power during the 72-h-exposure period with minimal thermal effects. With this refined experimental setup, exposure to 1.7 GHz LTE RF-EMF at the SAR ranging from 0.4 W/Kg to 8 W/Kg neither affects the proliferation of various human cells with different growth rates nor induces DNA damage and cell cycle perturbation. Before upgrading the exposure system, we observed that the exposure of human cells to 1.7 GHz RF-EMF increased or decreased cell proliferation, depending on the SAR values. In addition, we verified that exposure to 1.7 GHz RF-EMF with this refined system affected cell proliferation when heat was not properly controlled. Altogether, these results suggest that exposure to 1.7 GHz LTE RF-EMF does not directly influence cell proliferation and that the physiological changes induced by RF-EMF might be associated with thermal effects.

Open access paper: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0302936

--

The effects of radiofrequency exposure on cognition: A systematic review and meta-analysis of human observational studies

(My note: Systematic reviews require a multitude of macro- and micro-decisions. When conducted by biased investigative teams, systematic reviews generate erroneous conclusions and policy implications. By design, this review excluded most of the relevant research; hence, its conclusion is very limited in scope.)

Benke G, Abramson MJ, Brzozek C, McDonald S, Kelsall H, Sanagou M, Zeleke BM, Kaufman J, Brennan S, Verbeek J, Karipidis K. The effects of radiofrequency exposure on cognition: A systematic review and meta-analysis of human observational studies. Environment International. 2024. doi: 10.1016/j.envint.2024.108779.

Abstract

Background We aimed to assess evidence of long-term effects of exposure to radiofrequency (RF) electromagnetic fields (EMF) on indicators of cognition, including domains of learning and memory, executive function, complex attention, language, perceptual motor ability and social cognition, and of an exposure–response relationship between RF-EMF and cognition.

Methods We searched PubMed, Embase, PsycInfo and the EMF-Portal on September 30, 2022 without limiting by date or language of publication. We included cohort or case-control studies that evaluated the effects of RF exposure on cognitive function in one or more of the cognitive domains. Studies were rated for risk of bias using the OHAT tool and synthesised using fixed effects meta-analysis. We assessed the certainty of the evidence using the GRADE approach and considered modification by OHAT for assessing evidence of exposures.

Results We included 5 studies that reported analyses of data from 4 cohorts with 4639 participants consisting of 2808 adults and 1831 children across three countries (Australia, Singapore and Switzerland) conducted between 2006 and 2017. The main source of RF-EMF exposure was mobile (cell) phone use measured as calls per week or minutes per day. For mobile phone use in children, two studies (615 participants) that compared an increase in mobile phone use to a decrease or no change were included in meta-analyses. Learning and memory. There was little effect on accuracy (mean difference, MD −0.03; 95% CI −0.07 to 0.02) or response time (MD −0.01; 95% CI −0.04 to 0.02) on the one-back memory task; and accuracy (MD −0.02; 95%CI −0.04 to 0.00) or response time (MD −0.01; 95%CI −0.04 to 0.03) on the one card learning task (low certainty evidence for all outcomes). Executive function. There was little to no effect on the Stroop test for the time ratio ((B-A)/A) response (MD 0.02; 95% CI −0.01 to 0.04, very low certainty) or the time ratio ((D-C)/C) response (MD 0.00; 95% CI −0.06 to 0.05, very low certainty), with both tests measuring susceptibility to interference effects. Complex attention. There was little to no effect on detection task accuracy (MD 0.02; 95% CI −0.04 to 0.08), or response time (MD 0.02;95% CI 0.01 to 0.03), and little to no effect on identification task accuracy (MD 0.00; 95% CI −0.04 to 0.05) or response time (MD 0.00;95% CI −0.01 to 0.02) (low certainty evidence for all outcomes). No other cognitive domains were investigated in children. A single study among elderly people provided very low certainty evidence that more frequent mobile phone use may have little to no effect on the odds of a decline in global cognitive function (odds ratio, OR 0.81; 95% CI 0.42 to 1.58, 649 participants) or a decline in executive function (OR 1.07; 95% CI 0.37 to 3.05, 146 participants), and may lead to a small, probably unimportant, reduction in the odds of a decline in complex attention (OR 0.67;95%CI 0.27 to 1.68, 159 participants) and a decline in learning and memory (OR 0.75; 95% CI 0.29 to 1.99, 159 participants). An exposure–response relationship was not identified for any of the cognitive outcomes.

Discussion This systematic review and meta-analysis found only a few studies that provided very low to low certainty evidence of little to no association between RF-EMF exposure and learning and memory, executive function and complex attention. None of the studies among children reported on global cognitive function or other domains of cognition. Only one study reported a lack of an effect for all domains in elderly persons but this was of very low certainty evidence. Further studies are needed to

address all types of populations, exposures and cognitive outcomes, particularly studies investigating environmental and occupational exposure in adults. Future studies also need to address uncertainties in the assessment of exposure and standardise testing of specific domains of cognitive function to enable synthesis across studies and increase the certainty of the evidence.

Other This review was partially funded by the WHO radioprotection programme and prospectively registered on PROSPERO CRD42021257548.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: [Dr Geza Benke reports financial support was provided by World Health Organization. MA declares that he holds a small parcel of Telstra shares. KK, CB and MS as part of their employment are involved in the provision of advice to the Australian Commonwealth Government, Australian States and Territories and the general public on the risks and health effects of exposure to ionising and non-ionising radiation. KK is also a member of the International Commission on Non-Ionizing Radiation Protection where he contributes in the development and dissemination of science-based advice on limiting exposure to non-ionizing radiation. No member of the review team was involved in screening or extracting data from a study in which he or she was an author].

Acknowledgements

We would like to thank Emilie Van Deventer, WHO, Geneva, Switzerland; Martin Röösli, Swiss TPH, Basel, Switzerland and Anke Huss, Utrecht University, Utrecht, The Netherlands; for their contributions to the protocol.

Supplementary files can be downloaded.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024003659

--

Effects of recall and selection biases on modeling cancer risk from mobile phone use: Results from a case-control simulation study

(My note: The models tested in this simulation study were based on questionable assumptions.)

Bouaoun L, Byrnes G, Lagorio S, Feychting M, Abou-Bakre A, Beranger R, Schüz J. Effects of recall and selection biases on modeling cancer risk from mobile phone use: Results from a case-control simulation study. Epidemiology. 2024 May 20. doi: 10.1097/EDE.0000000000001749.

Abstract

Background: The largest case-control study (Interphone Study) investigating glioma risk related to mobile phone use showed a J-shaped relationship with reduced relative risks for moderate use and a 40% increased relative risk among the 10% heaviest regular mobile phone users, using a categorical risk model based on deciles of lifetime duration of use among ever regular users.

Methods: We conducted Monte-Carlo simulations examining whether the reported estimates are

compatible with an assumption of no effect of mobile phone use on glioma risk when the various forms of biases present in the Interphone study are accounted for. Four scenarios of sources of error in self-reported mobile phone use were considered, along with selection bias. Input parameters used for simulations were those obtained from Interphone validation studies on reporting accuracy and from using a non-response questionnaire.

Results: We found that the scenario simultaneously modeling systematic and random reporting errors produced a J-shaped relationship perfectly compatible with the observed relationship from the main Interphone study with a simulated spurious increased relative risk among heaviest users (OR = 1.91) compared to never regular users. The main determinant for producing this J shape was higher reporting error variance in cases compared to controls, as observed in the validation studies. Selection bias contributed to the reduced risks as well.

Conclusions: Some uncertainty remains, but the evidence from the present simulation study shifts the overall assessment to making it less likely that heavy mobile phone use is causally related to an increased glioma risk.

Open access paper: https://pubmed.ncbi.nlm.nih.gov/38771708/

--

Validation of mobile phone use recall in the multinational MOBI-kids study

van Wel L, Huss A, Kromhout H, Momoli F, Krewski D, Langer CE, Castaño-Vinyals G, Kundi M, Maule M, Miligi L, Sadetzki S, Albert A, Alguacil J, Aragones N, Badia F, Bruchim R, Goedhart G, de Llobet P, Kiyohara K, Kojimahara N, Lacour B, Morales-Suarez-Varela M, Radon K, Remen T, Weinmann T, Vrijheid M, Cardis E, Vermeulen R; MOBI-Kids consortium. Validation of mobile phone use recall in the multinational MOBI-kids study. Bioelectromagnetics. 2024 May 22. doi: 10.1002/bem.22507.

Abstract

Potential differential and non-differential recall error in mobile phone use (MPU) in the multinational MOBI-Kids case-control study were evaluated. We compared self-reported MPU with network operator billing record data up to 3 months, 1 year, and 2 years before the interview date from 702 subjects aged between 10 and 24 years in eight countries. Spearman rank correlations, Kappa coefficients and geometric mean ratios (GMRs) were used. No material differences in MPU recall estimates between cases and controls were observed. The Spearman rank correlation coefficients between self-reported and recorded MPU in the most recent 3 months were 0.57 and 0.59 for call number and for call duration, respectively. The number of calls was on average underestimated by the participants (GMR = 0.69), while the duration of calls was overestimated (GMR = 1.59). Country, years since start of using a mobile phone, age at time of interview, and sex did not appear to influence recall accuracy for either call number or call duration. A trend in recall error was seen with level of self-reported MPU, with underestimation of use at lower levels and overestimation of use at higher levels for both number and duration of calls. Although both systematic and random errors in self-reported MPU among participants were observed, there was no evidence of differential recall error between cases and controls. Nonetheless, these sources of exposure measurement error warrant consideration in interpretation of

the MOBI-Kids case-control study results on the association between children's use of mobile phones and potential brain cancer risk.

Highlights

- Self-reported and operator-registered phone calls were compared among MobiKids participants.
- On average, number of calls was underestimated, and duration overestimated.
- No differential recall error was found between brain tumor cases and controls in MobiKids.

Excerpts

The comparison of self-reported and recorded MPU was conducted separately for the number and duration (in minutes) of calls in the 3 months preceding the date of interview. In addition, a subset of subjects who had data available for 1 year, and a subset for 2 years preceding the date of interview were assessed....

A total of 702 subjects from eight different countries (Canada, France, Germany, Greece, Israel, Italy, Korea, and Spain) had data on both self-reported and recorded MPU in the 3 months preceding the interview date (Table 1)....

The overall GMRs of self-reported versus recorded MPU in the most recent 3 months were 1.59 for call duration and 0.69 for call number, indicating systematic errors in the form of overreporting of call duration of calls and underreporting of the number of calls. Looking at recall over time for subjects with 2 years of data available, there appears to be a lower level of overreporting of duration of calls for both cases (initial GMR 1.62, 1.44 at 1 year) and controls (initial GMR 1.37, 1.08 at 1 year) between recent and 1 year recall. There is however little difference in recall in call duration over time between the 1 and 2 year time points for either cases or controls (1.44 vs. 1.41 and 1.08 vs. 1.07, respectively) (Table 5). Although this same initial decrease can be seen in controls for recall of number of calls, it is less clear for cases (Table 4)....

While not all subjects from the main MOBI-Kids study provided informed consent to obtain their network operator data and not all operators provided data, we managed to include a large proportion of subjects (24.8%) from the MOBI-Kids case–control study in this validation study. The proportion of subjects where longer-term data was available (at 1- and 2-year time periods) was smaller, with no subjects from some of the participating countries....

... it may not always be clear from billing records who the actual user of the phone was when making calls. Therefore, while billing records provide insight into validity and possible calibration of self-reports, they may not represent the gold standard for studies, especially due to the low number of retained study participants, and also because internet-based calling (VoiP) is not included in the records....

Both the COSMOS (Reedijk et al., 2024) as well as the INTERPHONE validation (Vrijheid et al., 2009) studies found significantly differing ratios between countries. In the present study we did see some differences in GMRs among the eight participating countries, but the differences did not achieve statistical significance. In contrast, the MOBI-Expo validation study by Goedhart et al. (2018) did find significant differences among countries with participants from Greece, Israel, and Korea underestimating the duration of calls while in other countries duration was overestimated....

CONCLUSIONS

We compared self-reported MPU with operator data at 3 months, 1 year, and 2 years preceding the interview date. No indication of differential recall error between cases and controls was found. Both non-differential systematic and random errors were observed, with number of calls being underreported and duration of calls being over-reported on average in both cases and controls. If there are true underlying risks, then the observed non-differential random errors may bias risk estimates towards their null values and decrease study power.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.22507

--

Micro-environmental personal radio-frequency electromagnetic field exposures in Melbourne: A longitudinal trend analysis

Bhatt CR, Henderson S, Sanagou M, Brzozek C, Thielens A, Benke G, Loughran S. Micro-environmental personal radio-frequency electromagnetic field exposures in Melbourne: A longitudinal trend analysis. Environ Res. 2024 Mar 13;251(Pt 2):118629. doi: 10.1016/j.envres.2024.118629

Highlights

- RF-EMF exposure may be affected by changes in telecommunication technologies.
- Personal RF-EMF exposure assessments were undertaken in 2015/16 and 2022.
- Quantile regression models were used to compare exposure levels and overall trend.
- Median total exposure level at follow-up did not change over time.
- Overall trend of total exposure at follow-up increased.

Abstract

Background A knowledge gap exists regarding longitudinal assessment of personal radio-frequency electromagnetic field (RF-EMF) exposures globally. It is unclear how the change in telecommunication technology over the years translates to change in RF-EMF exposure. This study aims to evaluate longitudinal trends of micro-environmental personal RF-EMF exposures in Australia.

Methods The study utilised baseline (2015-16) and follow-up (2022) data on personal RF-EMF exposure (88 MHz-6 GHz) measured across 18 micro-environments in Melbourne. Simultaneous quantile regression analysis was conducted to compare exposure data distribution percentiles, particularly median (P50), upper extreme value (P99) and overall exposure trends. RF-EMF exposures were compared across six exposure source types: mobile downlink, mobile uplink, broadcast, 5G-New Radio, Others and Total (of the aforementioned sources). Frequency-specific exposures measured at baseline and follow-up were compared. Total exposure across different groups of micro-environment types were also compared.

Results For all micro-environmental data, total (median and P99) exposure levels did not significantly change at follow-up. Overall exposure trend of total exposure increased at follow-up. Mobile downlink contributed the highest exposure among all sources showing an increase in median exposure and overall

exposure trend. Of seven micro-environment types, five of them showed total exposure levels (median and P99) and overall exposure trend increased at follow-up.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0013935124005334

--

Evaluating radiofrequency electromagnetic field exposure in confined spaces: a systematic review of recent studies and future directions

Ahsan Ashraf M, Celik T. Evaluating radiofrequency electromagnetic field exposure in confined spaces: a systematic review of recent studies and future directions. Radiat Prot Dosimetry. 2024 Apr 20;200(6):598-616. doi: 10.1093/rpd/ncae045.

Abstract

This study reviews recent research on Radiofrequency Electromagnetic Field (RF-EMF) exposure in confined environments, focusing on methodologies and parameters. Studies typically evaluate RF-EMF exposure using an electric field and specific absorption rate but fail to consider temperature rise in the tissues in confined environments. The study highlights the investigation of RF-EMF exposure in subterranean environments such as subways, tunnels and mines. Future research should evaluate the exposure of communication devices in such environments, considering the surrounding environment. Such studies will aid in understanding the risks and developing effective mitigation strategies to protect workers and the general public.

Conclusion

In this study, we conducted an SLR of 63 articles related to RF-EMF exposure in confined environments. The articles were selected based on a QA and divided into measurement and simulation studies. Our literature analysis demonstrates that RF-EMF exposure is higher in working spaces and transport than in dwellings, largely due to multiple exposure frequencies and excessive users. However, evaluating sub-terrain environments remains limited despite the growing use of wearable wireless devices in these settings. We also identified the different exposimeters and numerical dosimetry software tools commonly used for measuring and simulating RF-EMF exposure in confined environments. Furthermore, we found that the electric field is the most commonly investigated parameter for evaluating RF-EMF exposure using dosimetry simulation and measurement in confined environments. Given the increasing use of wearable wireless devices in sub-terrain and other occupational environments, there is a critical need to evaluate RF-EMF exposure in high-temperature environments, particularly with respect to temperature elevation in tissue. Future studies should evaluate the potential health impacts of long-term exposure to RF-EMF, especially in occupational settings where workers are exposed for extended periods.

Open access paper: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11033578/

--

Optimal design of electromagnetic field exposure maps in large areas

López-Espí PL, Sánchez-Montero R, Guillén-Pina J, Chocano-del-Cerro R, Rojas JAM. Optimal design of electromagnetic field exposure maps in large areas. Environmental Impact Assessment Review. Volume 106, 2024, doi: 10.1016/j.eiar.2024.107525.

Abstract

The mapping of electromagnetic field (EMF) exposure over large areas is a very useful tool for the analysis of epidemiological data and risk assessment. Its production requires a costly measurement process. To optimize the effort and ensure the representativeness of the exposure map, criteria for the selection of the sites to be measured must be established. This paper presents a methodology for conducting EMF exposure maps suitable for risk assessment evaluation in large areas. The proposal combines radio wave propagation criteria and GIS methods to optimize the sampling effort. The design criteria are based on the determination of a rectangular grid of 250 m side and the identification of the emitters within the area under study. Both urban and rural sites are analysed in the proposal and line of sight conditions (LOS) are considered to reduce the number of points required and thus optimize the measurement effort. Depending on the extent and regularity of the surface, the density of measurement points has been estimated to be between 8 and 10 points per square kilometre in the urban area. The proposed methodology has been applied to a case study of a 2.8 km2 urban area within a 35.11 km2 municipality, obtaining an average point density of 9.64 points/km2 in the urban area. The differences in exposure depending on the application of the criteria have been analysed by means of the statistical values of the sets and the subtraction of the maps generated using kriging techniques. According to our results, if LOS measurements are not properly incorporated, the mean value of the EMF is underestimated in the area under study.

Conclusions

In this proposal, the methodology for EMF exposure mapping in large areas has been analysed. To optimize the measurement effort over a large area, a division into 250 × 250 m^2 urban grids should be considered, in which the possible sources of radiation should be known. A measurement under LOS conditions must be performed in each of the grids in which any of the emitters is present. The absence of measurements under LOS conditions implies an underestimation of the mean values. To select the rest of the grids to be measured, the viewshed analysis allows to simplify the number of measurements by grouping those grids determined as N-LOS. To achieve optimal interpolation results over the entire surface, sufficient measurements must also be available on its perimeter. Interpolation by ordinary stable kriging obtains adequate results under LOS and N-LOS conditions. The generated maps contribute to a better perception of risk as they provide an objective and simple tool to show the level of EMF. They do not require complex propagation models but are based on interpolations made within a GIS. They can also be combined, using these techniques, for risk assessment and even, if epidemiological data are available, for possible correlations studies.

The proposal considers the differences between urban areas, with the possibility of multipath, and rural (non-urban) areas. For the analysis of urban areas, a grid of 250 × 250 m^2 has been proposed as a guideline for the spatial organisation of the territory in urban areas and 500 × 500 m^2 in rural areas.

Depending on the extent and regularity of the surface, the density of measurement points has been estimated to be between 8 and 10 points per square kilometre in the urban area. Lower point densities

result in IDW representations and higher densities increase the measurement effort excessively. In the case under study, for the urban area, a density of 9.64 points/km^2 has been reached.

The study has been performed following the six-minute averaging criterion and in a bandwidth from 100 kHz to 3 GHz. The latest ICNIRP recommendations modify these values to take into account, among others, the new 5G signals, not yet present in the area. The urban study was carried out in a mainly residential area, where there are no concentrations of the general public, which could lead to areas of greater exposure due to greater use of the network. For future studies, therefore, the measurement and comparison according to the latest ICNIRP recommendations is still pending in areas where the new 5G networks have been incorporated and in urban micro-environments where there is a higher population density or uses significantly different from residential. Another point of continuation of this research is the correlation of the measured values with tumor statistics and other diseases. The opportunity provided by exposure maps created for entire urban environments may allow, where appropriate, to find possible relationships.

https://www.sciencedirect.com/science/article/pii/S0195925524001124

--

Electromagnetic field exposure monitoring of commercial 28-GHz band 5G base stations in Tokyo, Japan

Liu S, Tobita K, Onishi T, Taki M, Watanabe S. Electromagnetic field exposure monitoring of commercial 28-GHz band 5G base stations in Tokyo, Japan. Bioelectromagnetics. 2024 May 22. doi: 10.1002/bem.22505.

Abstract

Fifth generation (5G) wireless communication is being rolled out around the world. In this work, the latest radio frequency electromagnetic field (EMF) exposure measurement results on commercial 28-GHz band 5G base stations (BSs) deployed in the urban area of Tokyo, Japan, are presented. The measurements were conducted under realistic traffic conditions with a 5G smartphone and using both omnidirectional and horn antennas. First and foremost, in all cases, the electric-field (E-field) intensity is much lower (<-38 dB) than the exposure limits. The E-field intensities for traffic-off cases do not show any significant difference between the two antennas with the maximum being 3.6 dB. For traffic-on cases, the omnidirectional antenna can undesirably capture the radio wave from the smartphone in some cases, resulting in a 7-13 dB higher E-field intensity than that using the horn antenna. We also present comparative results between 4G long term evolution BSs and sub-6-GHz band and 28-GHz band 5G BSs and provide recommendations on acquiring meaningful EMF exposure data. This work is a further step toward the standardization of the measurement method regarding quasi-millimeter/millimeter wave 5G BSs.

https://pubmed.ncbi.nlm.nih.gov/38778514/

Conclusions

The latest EMF exposure measurement results on commercial 28-GHz band 5G FR2 BSs in Tokyo, Japan, were presented in this paper. The measurements were evaluated under realistic traffic conditions and in an isotropic state using both omnidirectional and horn antennas. In all cases (sites and spots, antennas,

and traffic conditions), the E-field intensity was within the limits established by ICNIRP and IEEE/ICES. The maximal time-averaged E-field intensity (118 dBμV/m) reached only 0.016% of the limit. It was found that the E-field intensity varied with measurement height with a variance in amplitude reaching 10–25 dB over different measurement spots. The E-field intensities for traffic-off cases did not show any significant difference between the two antennas with the maximum being 3.6 dB. However, for traffic-on cases, the omnidirectional antenna may undesirably capture the radio wave from the smartphone in some cases, thus leading to a time-averaged E-field intensity 7–13 dB higher than that in the cases of using the horn antenna. Despite this, an omnidirectional antenna is always preferred as it can detect all the reflections and scatterings of the environment. A horn antenna is preferred only in situations where the reflections and scatterings are negligible. On the other hand, the distance between the smartphone and the antenna (50 cm in this study) may be increased to avoid the capture of the irradiation from the smartphone. For the horn antenna, there is generally a 23–27 dB rise in the time-averaged E-field intensity of traffic-on cases compared with that of traffic-off cases. The rise for the omnidirectional antenna is 19–37 dB.

5G, especially 5G FR2, is being continuously deployed. The measurements in this work were mainly limited to the urban area of Tokyo, Japan. With the expansion of the 5G service coverage, measurements in suburban and rural areas will be conducted in the future. Comparative studies between 5G FR1 and FR2 will also be of considerable interest and will be a future target. In summary, investigations will be successively conducted following the deployment of 5G.

--

The exposure of nonhuman living organisms to mobile communication emissions: A survey to establish European stakeholders' policy option preferences

Recuero Virto L, Thielens A, Czerwiński M, Froidevaux J. (2024). The exposure of nonhuman living organisms to mobile communication emissions: A survey to establish European stakeholders' policy option preferences. *Risk Analysis*, 1–15. doi: 10.1111/risa.14322.

Abstract

There is an unprecedented exposure of living organisms to mobile communications radiofrequency electromagnetic field (RF-EMF) emissions. Guidelines on exposure thresholds to limit thermal effects from these emissions are restricted to humans. However, tissue heating can occur in all living organisms that are exposed. In addition, exposure at millimetric frequencies used by 5G may impact surface tissues and organs of plants and small-size species. It is also expected that the addition of 5G to existing networks will intensify radiofrequency absorption by living organisms.

A European Parliament report proposed policy options on the effects of RF-EMF exposure of plants, animals, and other living organisms in the context of 5G: funding more research, implementing monitoring networks, accessing more information from operators on antennas and EMF emissions, and developing compliance studies when antennas are installed. However, there is no evidence on the preferences of relevant stakeholders regarding these policy options. This paper reports the findings of a survey of key European stakeholders' policy option preferences based on the European Parliament's report. It reveals a broad consensus on funding more research on the effects of exposure of plants, animals, and other living organisms to EMFs. It also highlights the need for deliberation concerning the

other policy options that could provide solutions for regulatory authorities, central administrations, the private sector, nongovernmental associations and advocates, and academics. Such deliberation would pave the way for effective solutions, focusing on long-term output from funding research, and enabling short-term socially and economically acceptable actions for all parties concerned.

Excerpts

From a regulatory perspective, current guidelines on exposure thresholds to limit thermal energy from mobile communications emissions developed by the International Commission on Non-Ionizing Radiation Protection (ICNIRP) and by the Institute of Electrical and Electronics Engineers (IEEE) only concern human health (ICNIRP, 1998, 2020; IEEE, 1991, 2019).3 It is worth noting that ICNIRP and IEEE thresholds, developed under the assumptions that biological effects are due to excessive tissue heating and that no effects occur below the threshold, remain controversial (ICBE-EMF, 2022). According to the International Commission on the Biological Effects of Electromagnetic Fields (ICBE-EMF), adverse effects have been observed below the threshold, such as nonthermal induction of reactive oxygen species, DNA damage, cardiomyopathy, carcinogenicity, sperm damage, and neurological effects (e.g., electromagnetic hypersensitivity)....

According to the International Commission on Radiological Protection (ICRP), steps taken to protect the environment from ionizing radiation should fall within the concept of optimization, the latter understood as the process of determining the level of protection and safety that makes exposures, and the probability and magnitude of potential exposures, as low as reasonably achievable, economic and societal factors being taken into account. Reference levels were derived by the ICRP to relate the effects of ionizing radiation to doses to nonhuman organisms, but the institution states that these values are not limited and are not intended to be used in that manner. Potential synergies between ICNIRP and ICRP approaches were already subject to discussion in the past for human health, in particular, whether ALARA approaches developed by the ICRP could be meaningfully applied to nonionizing radiation (ICNIRP/WHO/ICRP/IRPA/ILO, 2014).

To conclude, humans have so far chosen not to address the effects of RF-EMF on nonhuman organisms even if there is evidence of thermal effects on all living organisms. The controversial debates on RF-EMF effects on human health have probably hampered early attempts to address the effects on nonhuman organisms. Our results show that, for the consulted stakeholders, this topic calls for some action. It is not a matter of hindering progress, but understanding the optimal conditions for technological proliferation, given the current state of knowledge.

https://onlinelibrary.wiley.com/doi/10.1111/risa.14322

--

RF-EMF Exposure Assessment of Fetus During The First Trimester of Pregnancy

Sandeep S, Vard A, Guxens M, Bloch I, Wiart J. RF-EMF Exposure Assessment of Fetus During The First Trimester of Pregnancy. *IEEE Access*, doi: 10.1109/ACCESS.2024.3404369.

Abstract

This article describes the computational analysis of Radio Frequency - Electromagnetic Field (RF-EMF) exposure of Uterus-Fetus Units (UFUs) embedded inside the body of a 26 year old human female. Realistic UFU models are obtained from ultrasound images acquired for different fetuses and at specific development stages (7 weeks, 9 weeks and 11 weeks old), for which a deep-learning based segmentation method is developed. Each UFU model is then inserted into a computational electromagnetic model of a 26 year old female. The Specific Absorption Rate (SAR) of the fetus at commonly used wireless communication frequencies is estimated using a commercially available numerical electromagnetic solver. The Inverted F antenna (IFA), which is a commonly used mobile phone antenna was used as the excitation source. Fetus SAR values are reported for different combinations of excitation frequencies, phone positions and UFU ages. It was found that the fetus SAR for all the cases is well below the maximum allowable exposure limit of 80 mW/kg, as prescribed by ICNIRP. Furthermore, we replaced the embryo with uterus tissues and calculated the SAR in the uterus tissues (i.e. uterus tissues with same volume and shape, and at the same location as that of UFU). The uterus SAR values were found to be only marginally different from that of fetus SAR.

Open access paper:
https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=10536873&isnumber=6514899

--

Regression calibration of self-reported mobile phone use to optimize quantitative risk estimation in the COSMOS study

Reedijk M, Portengen L, Auvinen A, Kojo K, Heinävaara S, Feychting M, Tettamanti G, Hillert L, Elliott P, Toledano MB, Smith RB, Heller J, Schüz J, Deltour I, Poulsen AH, Johansen C, Verheij R, Peeters P, Rookus M, Traini E, Huss A, Kromhout H, Vermeulen R, Study Group TC. Regression calibration of self-reported mobile phone use to optimize quantitative risk estimation in the COSMOS study. Am J Epidemiol. 2024 May 13:kwae039. doi: 10.1093/aje/kwae039.

Abstract

The Cohort Study of Mobile Phone Use and Health (COSMOS) has repeatedly collected self-reported and operator-recorded data on mobile phone use. Assessing health effects using self-reported information is prone to measurement error, but operator data were available prospectively for only part of the study population and did not cover past mobile phone use. To optimize the available data and reduce bias, we evaluated different statistical approaches for constructing mobile phone exposure histories within COSMOS. We evaluated and compared the performance of four regression calibration (RC) methods (simple, direct, inverse, and generalized additive model for location, shape, and scale), complete-case (CC) analysis and multiple imputation (MI) in a simulation study with a binary health outcome. We used self-reported and operator-recorded mobile phone call data collected at baseline (2007-2012) from participants in Denmark, Finland, the Netherlands, Sweden, and the UK. Parameter estimates obtained using simple, direct, and inverse RC methods were associated with less bias and lower mean squared error than those obtained with CC analysis or MI. We showed that RC methods resulted in more accurate estimation of the relation between mobile phone use and health outcomes, by combining self-reported data with objective operator-recorded data available for a subset of participants.

Excerpts

Self-reported duration of mobile phone use (REPORT) at baseline [21] was based on answers to the question: “Over the last three months, on average, how much time per week did you spend talking on a mobile phone?”. The following response options were provided: “< 5 min/week”, “5-29 min/week”, “30-59 min/week”, “1-3 hours/week”, “4-6 hours/week”, and “>6 hours/week”. In the Netherlands and the UK two further categories of call duration were included, “7-9 hours/week” and “10 or more hours/week”, but these were combined into “>6 hours/week” for the present analyses....

Operator-recorded duration of mobile phone use (RECORD) was collected for all participants who provided consent and had a subscription under their own name. Operator-recorded data were used only when available for all mobile phones that were reported (up to two phones in Denmark, the Netherlands, Sweden, Finland, and up to three in the UK) and for at least three full months at the time the baseline questionnaire was administered. Participants with mobile phones that were also used by others were excluded. Operator-recorded data were available for 21% of participants in Denmark, 76% in Finland, 40% in the Netherlands, 63% in Sweden, and 76% in the UK....

A major issue is how well mobile phone use predicts the exposure of interest, namely radiofrequency electromagnetic fields. While past validation studies have been carried out for the 2nd mobile phone generation, showing fair agreement between amount of use and cumulated emission from the handset [4], fewer data are available on the predictive power of mobile phones of the 3rd, 4th and 5th generations that have and are being used by COSMOS participants....

Table 1 (excerpt)

Complete data by country: Denmark (11%: 2993/25912), Finland (70%: 9162/13062), Netherlands (3%: 3039/88466), Sweden (49%: 24881/50678), United Kingdom (58%: 56862/98685)

Recorded call duration in minutes/week by country (geometric mean): Denmark (60.6), Finland (81.5), Netherlands (23.4), Sweden (78.3), United Kingdom (46.5)

Conclusion

This study addressed an important concern in mobile phone research and more generally in environmental epidemiology: how to leverage self-reported exposure estimates that are often available but error-prone, with more objective measurements that may be obtained in only a subset of participants. Our simulation study indicated RC approaches may improve estimation of exposure-outcome relations between mobile phone use and health outcomes within COSMOS. The prospective design and improved exposure assessment within COSMOS compared to that in previous case-control studies are expected to lead to more robust conclusions about possible health effects from use of mobile phones.

Conflict of Interest: MF was vice chairman (2012-2020) of the International Commission on Non-Ionizing Radiation Protection, an independent body setting guidelines for non-ionizing radiation protection. She has served as advisor to a number of national and international public advisory and research steering groups concerning the potential health effects of exposure to non-ionizing radiation, currently for the World Health Organization (WHO). HK was the chair of the Committee on Electromagnetic Fields of the Health Council of The Netherlands till 2022. He currently is a member of the WHO Task Group for the Environmental Health Criteria Monograph on RF-EMF. AH is a member of the International Commission on Non-Ionizing Radiation Protection since 2020, and of the Committee on Electromagnetic Fields of the

Health Council of The Netherlands, and chairs the Swedish Radiation Safety Authority's (SSM) Scientific Council on Electromagnetic Fields since 2020. AA currently is a member of the WHO Task Group for the Environmental Health Criteria Monograph on RF-EMF. MBT is currently member of the WHO groups tasked with systematic review of evidence on non-ionizing radiation and health, that is feeding into the Environmental Health Criteria Monograph on RF-EMF. All other authors declare they have no competing financial interests.

Open access paper:

https://academic.oup.com/aje/advance-article/doi/10.1093/aje/kwae039/7671112?login=false

--

Monitoring of the exposure to electromagnetic fields with autonomous probes installed outdoors in France

Jawad O, Conil E, Agnani J-B, Wang S, Wiart J. Monitoring of the exposure to electromagnetic fields with autonomous probes installed outdoors in France. Comptes Rendus. Physique, Online first (2024), pp. 1-21. doi : 10.5802/crphys.182.

Abstract

The study is based on a new temporal analysis of exposure based on the deployment of autonomous broadband E-field monitoring probes in many French cities. The combination of the probe's data with frequency-selective in situ measurements performed by ANFR and the knowledge of the nearby base station antennas, allows to draw statistical conclusions on the exposure of the population. Indeed, the data collected by the probes reveal that different periodicities exist (seasonality, day/night). This paper shows that the monitoring probes are able to detect the seasonality of the exposure and provide analysis of correlation between monitoring probes and radio environment.

Excerpt

The analysis indicates that the monitoring probes have varying exposure levels. Probes measuring significant levels show a difference in exposure between day and night, a phenomenon observed for the first time in France. An empirical time interval from 8 AM to 11 PM enables to calculate the ratio of averaged E-field levels between day and night. This ratio is between 1.28 and 1.42 for the three probes with the highest RMS level. Several papers, such as [8–10], have characterized the day and night fluctuation, but this has never been demonstrated using French data.

For the first time, the variability of daily working hours has been quantified for all the probes installed in France. The data shows that most of the probes exhibit a 30% variation percentage based on the data gathered from 8 AM to 5 PM. Based on our knowledge, this is the first time that the assessment of the daily variation contributor based on more than 150 probes installed in different environments is achieved. It confirms the level of daily variation contributor to the in situ measurement uncertainty budget, as communicated in the accredited in situ measurement reports [3, 16, 17]....

For the first time, the seasonality of the level of exposure has been analyzed at the French national level. The PCA on dataset No. 2 emphasizes the observation that the positioning of the probe is crucial to

observe a remarkable variation of the exposure level. The probe must not only be installed in an area with many base stations, but also in close proximity to them and in a line-of-sight position for the exposure. A large number of probes measuring low exposure levels can be explained by the fact that the probes are not in a line-of-sight situation. In some cities, probes were installed in low-density areas rather than in front of base stations due to public concern over electromagnetic waves. Upon analysis, we find that some probes measure a lower level of exposure in summer time. This interpretation was confirmed with the Principal Component Analysis, which showed that the second component of the PCA characterizes the difference between probes with higher exposure in summer or winter. This phenomenon can be explained by the fact that some of the cities are very touristic during the summer, leading to increased use of the telecommunication infrastructure. In general, the exposure levels measured by the autonomous probe are very low compared to the limits. The increase of the exposure level is relatively slow, as it has been shown in several ANFR studies [4, 18–20].

Open access paper:

https://comptes-rendus.academie-sciences.fr/physique/articles/10.5802/crphys.182/

--

Statistical Analysis of RF-EMF Exposure Induced by Cellular Wireless Networks in Public Transportation Facilities of the Paris Regionx

Y. Zhang *et al*. Statistical Analysis of RF-EMF Exposure Induced by Cellular Wireless Networks in Public Transportation Facilities of the Paris Regionx. *IEEE Access*, vol. 12, pp. 79741-79753, 2024, doi: 10.1109/ACCESS.2024.3410090.

Abstract

Wireless communications are increasingly used today. Despite such use, there is a significant perception of risk which makes exposure monitoring a significant concern today. The work described in this article was carried out within the framework of the European SEAWave project and the French Beyond5G project. The exposure assessment was evaluated using a personal exposimeter (MVG EMF Spy) whose compactness and ease of use make it more suitable and portable than a system combining measuring probes and spectrum analyzers. Measurements were carried out on the cellular frequency bands used by 2G, 3G, 4G, and 5G, as well as that of Wi-Fi, in different modes of public transportation (RER, metro, tramway, bus, and train) circulating in the Paris region. The measurements have been analyzed by frequency band, type of public transportation, and type of environment encountered. For each set of measurements (e.g., metro lines, tramways), the mean, standard deviation, skewness, and kurtosis were evaluated and analyzed. For all exposure measurements taken in the 700, 800, 900, 1800, 2100, 2600, and 3500 MHz frequency bands, the overall average values are 0.39, 0.43, 0.30, 0.21, 0.18, 0.24 and 0.18 V/m, respectively. These measurements have, in all cases, a significant dispersion as shown by the ratios of standard deviations to mean values. The well-known K-means clustering technique was applied to these four parameters for different subsets of data. The number of clusters k = 3 has been chosen based on the analysis of the optimal value of k for the current dataset. Our analysis indicates that the first group's members display the highest mean values with moderate variance and the lowest values for the third and fourth moments. The second cluster is distinguished by points with large mean and variance,

accompanied by moderate skewness and kurtosis. Conversely, the third group comprises points with the smallest mean and variance values, yet the largest measurements for the third and fourth moments.

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=10549999&isnumber=10380310

--

Expression levels of tam receptors and ligands in the testes of rats exposed to short and middle-term 2100 MHz radiofrequency radiation

Katirci E, Kirimlioglu E, Oflamaz AO, Hidisoglu E, Cernomorcenco A, Yargıcoğlu P, Ozen S, Demir N. Expression levels of tam receptors and ligands in the testes of rats exposed to short and middle-term 2100 MHz radiofrequency radiation. Bioelectromagnetics. 2024 May 9. doi: 10.1002/bem.22504.

Abstract

With advances in technology, the emission of radiofrequency radiation (RFR) into the environment, particularly from mobile devices, has become a growing concern. Tyro 3, Axl, and Mer (TAM) receptors and their ligands are essential for spermatogenesis and testosterone production. RFR has been shown to induce testicular cell apoptosis by causing inflammation and disrupting homeostasis. This study aimed to investigate the role of TAM receptors and ligands in the maintenance of homeostasis and elimination of apoptotic cells in the testes (weeks), short-term sham exposure (sham/1 week), and middle-term sham exposure (sham/10 weeks). Testicular morphology was assessed using hematoxylin-eosin staining, while immunohistochemical staining was performed to assess expression levels of TAM receptors and ligands in the testes of all groups. The results showed that testicular morphology was normal in the control, sham/1 week, and sham/10 weeks groups. However, abnormal processes of spermatogenesis and seminiferous tubule morphology were observed in RFR exposure groups. Cleaved Caspase 3 immunoreactivity showed statistically significant difference in 1 and 10 weeks exposure groups compared to control group. Moreover, there was no significant difference in the immunoreactivity of Tyro 3, Axl, Mer, Gas 6, and Pros 1 between groups. Moreover, Tyro 3 expression in Sertoli cells was statistically significantly increased in RFR exposure groups compared to the control. Taken together, the results suggest that RFR exposure negatively affects TAM signalling, preventing the clearance of apoptotic cells, and this process may lead to infection and inflammation. As a result, rat testicular morphology and function may be impaired.

https://pubmed.ncbi.nlm.nih.gov/38725116/

Highlights

Exposure to radiofrequency radiation (RFR) disrupts normal seminiferous tubule morphology.
RFR exposure in round spermatids may induce apoptosis independent of Tyro 3, Axl, and Mer (TAM) receptors.
TAM receptors and ligands are expressed in rat testis.

Conclusions

TAM receptors and ligands may play an important role in the spermatogenesis processes and may affect the functions of Leydig cells and Sertoli cells. In the present study, we have reported that RFR exposure

impair the morphology of rat testis. However, the impact on testicular function remains uncertain and requires further investigation. RFR exposure may negatively affect TAM signaling and enhances the accumulation of apoptotic cells.

Humans are exposed to RFR at a variable rate on a daily basis. Cell phones are the main source of RFR exposure, further complicating personal exposure times. In this study, cleaved caspase 3, TAM receptors and ligands were evaluated in rat testicular tissue exposed to 2100 MHz RFR. This study highlights the importance of radiofrequency exposure in male infertility. It provides a model for RFR reproductive toxicity studies. In order to better study the toxicity potential of RFR in the male reproductive system, it is necessary to study the intrauterine development and possibly live/healthy birth rates of the fetuses produced by mating RFR-exposed male mice with healthy female mice. In addition, long-term animal studies are urgently needed to investigate the potential health effects of frequencies used by upcoming 5G on the male reproductive system.

--

Effects of non-ionizing radio frequency electromagnetic radiation on the development and behavior of early embryos of *Danio rerio* (zebrafish)

Khira R, Uggini GK. Effects of non-ionizing radio frequency electromagnetic radiation on the development and behavior of early embryos of *Danio rerio*. Electromagn Biol Med. 2024 May 12:1-8. doi: 10.1080/15368378.2024.2352429.

Abstract

Biological effects of radio frequency electromagnetic radiation (RF-EMR) in the range of 900-1800 MHz emerging from the mobile phone were investigated and were found to influence the locomotor pattern when exposure was initiated from 1 hour post fertilization (hpf) in zebrafish embryos (ZE), *Danio rerio*. Mobile phones and other wireless devices offer tremendous advantages. However, on the flipside they are leading to an increased electromagnetic energy in the environment, an excess of which could be termed as electromagnetic pollution. Herein, we tried to understand the effects of RF-EMR emerging from the mobile phone, on the development and behavior of ZE, exposed to RF-EMR (specific absorption rate of 1.13 W/kg and 1800 MHz frequency) 1 hr daily, for 5 days. To understand if there could be any developmental stage-specific vulnerability to RF-EMR, the exposure was initiated at three different time points: 1hpf, 6hpf and 24hpf of ZE development. Observations revealed no significant changes in the survival rate, morphology, oxidative stress or cortisol levels. However, statistically significant variations were observed in the batch where exposure started at 1 hpf, with respect to locomotion patterns (distance travelled: 659.1 ± 173.1 mm Vs 963.5 ± 200.4 mm), which could be correlated to anxiety-like behavior; along with a corresponding increase in yolk consumption (yolk sac area: 0.251 ± 0.019 mm^2 Vs 0.225 ± 0.018 mm^2). Therefore, we conclude that RF-EMR exposure influences the organism maximally during the earliest stage of development, and we also believe that an increase in the time of exposure (corresponding to the patterns of current usage of mobile phones) might reveal added afflictions.

Plain Language Summary

Mobile phones and other wireless devices are on a rampant usage worldwide. They work by radiating low energy radiofrequency electromagnetic radiations. An excessive usage of wireless devices is leading

to increased presence of these radiations in our surroundings. Since these radiations are not physically sensed by the organisms, its impact stays elusive. Nevertheless, the interaction of these radiations with biological systems may produce some unwarranted effects. When we exposed the ZE to the mobile phone radiation daily 1 hr for 5 days, our observations revealed that the youngest of the experimental group showed susceptibility. The effect was evident through haphazard movements and stressed behavior. So, it is important to be aware of the potential effects and take necessary precautions by following safety guidelines, especially when the organism is in its early life stage.

https://pubmed.ncbi.nlm.nih.gov/38734994

--

Multi-frequency electromagnetic radiation induces anxiety in mice via inflammation in the cerebral cortex

Jing X, Menghua L, Lihui Z, Qian W, Xueli W, Xuelong Z, Zhihui L, Guofu D, Changzhen W. Multi-frequency electromagnetic radiation induces anxiety in mice via inflammation in the cerebral cortex. Environ Sci Pollut Res Int. 2024 May 9. doi: 10.1007/s11356-024-33447-y.

Abstract

Modern life is filled with radiofrequency electromagnetic radiation (RF-EMR) in various frequency bands, while the health risks are not clear. In this study, mice were whole-body exposed to 0.9/1.5/2.65 GHz radiofrequency radiation at 4 W/kg for 2 h per day for 4 weeks to investigate the emotional effects. It was found that the mice showed anxiety but no severe depression. The ELISA results showed a significant decrease in amino acid neurotransmitters (GABA, DA, 5-HT), although acetylcholine (ACH) levels were not significantly altered. Furthermore, Western blot results showed that BDNF, TrkB, and CREB levels were increased in the cerebral cortex, while NF-κB levels were decreased. In addition, pro-inflammatory factors (IL-6, IL-1β, TNF-α) were significantly elevated, and anti-inflammatory factors (IL-4, IL-10) tended to decrease. In conclusion, multi-frequency electromagnetic radiation induces an inflammatory response through the CREB-BDNF-TrkB and NF-κB pathways in the cerebral cortex and causes a decrease in excitatory neurotransmitters, which ultimately causes anxiety in mice.

Excerpt

In conclusion, these findings suggest that multi-frequency RF-EMR may induce anxiety by decreasing brain neurotransmitter levels and triggering inflammatory responses via the CREB-BDNF-TrkB pathway (Fig. 5). Currently, there are few studies on the relationship between multi-frequency electromagnetic radiation and mood changes, and this study provides some data support and a direction for future research. However, further in-depth investigations are needed to elucidate the specific mechanism of whether multi-frequency RF-EMR can induce negative emotions.

https://pubmed.ncbi.nlm.nih.gov/38724846/

--

The endocannabinoid system is involved in the anxiety-like behavior induced by dual-frequency 2.65/0.8 GHz electromagnetic radiation in mice

Xue T, Ma RH, Xu C, Sun B, Yan DF, Liu XM, Gao D, Li ZH, Gao Y, Wang CZ. The endocannabinoid system is involved in the anxiety-like behavior induced by dual-frequency 2.65/0.8 GHz electromagnetic radiation in mice. Front Mol Neurosci. 2024 Apr 15;17:1366855. doi: 10.3389/fnmol.2024.1366855.

Abstract

As wireless communication devices gain popularity, concerns about the potential risks of environmental exposure to complex frequency electromagnetic radiation (EMR) on mental health have become a public health issue. Historically, EMR research has predominantly focused on single- frequency electromagnetic waves, neglecting the study of multi-frequency electromagnetic waves, which more accurately represent everyday life. To address these concerns, our study compared the emotional effects of single-frequency and dual-frequency EMR while exploring potential molecular mechanisms and intervention targets. Our results revealed that single-frequency EMR at 2.65 or 0.8 GHz did not induce anxiety-like behavior in mice. However, exposure to dual-frequency EMR at 2.65/0.8 GHz significantly led to anxiety-like behavior in mice. Further analysis of mouse sera revealed substantial increases in corticosterone and corticotrophin releasing hormone levels following exposure to 2.65/0.8 GHz EMR. Transcriptome sequencing indicated a significant decrease in the expression of *Cnr1*, encoding cannabinoid receptor 1 Type (CB1R), in the cerebral. This finding was consistently verified through western blot analysis, revealing a substantial reduction in CB1R content. Additionally, a significant decrease in the endocannabinoid 2-arachidonoylglycerol was observed in the cerebral cortex. Remarkably, administering the cannabinoid receptor agonist Win55-212-2 significantly alleviated the anxiety-like behavior, and the cannabinoid receptor antagonist AM251 effectively counteracted the anti-anxiety effects of Win55-212-2. In summary, our research confirmed that dual-frequency EMR is more likely to induce anxiety-like behavior in mice than single-frequency EMR, with implications for the hypothalamic–pituitary–adrenal axis and the endocannabinoid system. Furthermore, our findings suggest that Win55-212-2 may represent a novel avenue for researching and developing anti-EMR drugs.

Open access paper: https://www.frontiersin.org/articles/10.3389/fnmol.2024.1366855/full

--

Assessment of the potential threats to brain health posed by the radiation from 5G sub-6 GHz base stations in China using dosimetric methods

Lin J, Ding G, Liu X, Li J. Assessment of the potential threats to brain health posed by the radiation from 5G sub-6 GHz base stations in China using dosimetric methods. Environ Sci Pollut Res Int. 2024 May;31(21):31015-31027. doi: 10.1007/s11356-024-33172-6.

Abstract

The 5G sub-6 GHz radio frequency (RF) electromagnetic fields (EMF) are the most widely used in China's communications. The public has expressed concerns about possible brain health effects of the higher frequency bands in 5G compared to 2G, 3G, and 4G bands. It is imperative to empirically investigate the potential health hazards of these novel frequency bands in 5G communication technology. This study evaluates the assessment of brain tissue dose coupling from sub-6 GHz band EMF emitted by base stations in China. Based on the 3D virtual human body model, the simulation environment was established. Dose including specific absorption rate (SAR) and internal electric field (IEF) between 2G, 3G, and 4G bands and 5G sub-6 GHz was investigated using normalized exposure values and exposure

limits. The results indicate that the sub-6 GHz high-frequency band of 5G has the lowest dose value. It can be concluded that high-frequency electromagnetic radiation in 5G sub-6 GHz reduces the dose and health threats to the brain. This provides strong support for the promotion of 5G commutation in China and other regions.

Conclusion

The research employed dosimetric methodology to comparatively evaluate the potential threat of communication EMF emitted by base stations to the human brain in China. Normalized evaluation showed that the dose value of brain tissue decreased with the increase of frequency. Moreover, the study explored the brain dose and health threats in accordance with both international mainstream ICNIRP guidelines and Chinese national standards. The findings indicate that the sub-6 GHz high-frequency band of 5G has the lowest dose value, thus ensuring a low risk of use. However, it should be noted that the choice of industry standards and physical evaluation quantities can slightly influence the evaluation results and trends. The research supplements the inadequacy of laboratory animal experiments and comparatively elaborates on the potential harm of EMF exposure from 5G sub-6 GHz to the brain's health. From a perspective of normalized assessment and the guideline comparison assessment, it can be concluded that high-frequency electromagnetic radiation in 5G sub-6 GHz reduces the harm of dose and risk values. This provides strong support for the promotion of 5G commutation in China and other regions. However, the limitation of this study is that the actual public exposure value does not take into account the measured value as the massive geographic information and statistical data. In the subsequent stage, the investigation of exposure measurement and dose evaluation can be merged with epidemiological research to probe deeper into the repercussions of novel frequency bands on the brain and its affiliated organ system.

Open access paper: https://link.springer.com/article/10.1007/s11356-024-33172-6

--

Whole-body SAR measurements of millimeter-wave base station in reverberation chambers

Eilers Bischoff J, Joshi P, Colombi D, Xu B, Törnevik C. Whole-body SAR measurements of millimeter-wave base station in reverberation chambers. International Journal of Microwave and Wireless Technologies. Published online 2024:1-6. doi:10.1017/S1759078724000515

Abstract

This paper presents a method for measuring whole-body specific absorption rate (WBSAR) of millimeter-wave base stations (BSs) in a reverberation chamber (RC). The absorbed power in the phantom from the equipment under test (EUT) and hence WBSAR is determined as the difference between the total radiated power with and without the phantom. A chamber transfer function is determined and used to include only the absorption in the phantom due to direct illumination from the EUT, i.e., excluding absorption due to the RC multipath reflections. The measurement method was evaluated at 28 GHz using a horn antenna and a commercial massive multi-input–multi-output BS. The experimental results are in good agreement with simulations. The proposed method allows for measurements of WBSAR within 3 minutes, which is much shorter than traditional approaches. The method is suitable for compliance assessments of BS products with the International Commission on Non-Ionizing Radiation

Protection 2020 electromagnetic field exposure guidelines, which extend the applicability of WBSAR as basic restrictions up to 300 GHz.

https://doi.org/10.1017/S1759078724000515

--

Retraction notice to "Exercise ameliorates hippocampal damage induced by Wi-Fi radiation; A biochemical, histological, and immunohistochemical study"

Mohamed AO, Abdel Hafez SMN, Ibrahim RA, Rifaai RA. Retraction notice to "Exercise ameliorates hippocampal damage induced by Wi-Fi radiation; A biochemical, histological, and immunohistochemical study" [J. Chem. Neuroanat. 129 (2023) 102252]. J Chem Neuroanat. 2024 Jun 8:102433. doi: 10.1016/j.jchemneu.2024.102433. Epub ahead of print. PMID: 38851999.

https://pubmed.ncbi.nlm.nih.gov/38851999/

--

Review on the impact of cell phone radiation effects on green plants

Panda DK, Das DP, Behera SK, Dhal NK. Review on the impact of cell phone radiation effects on green plants. Environ Monit Assess. 2024 May 21;196(6):565. doi: 10.1007/s10661-024-12623-0.

Abstract

The aim of this review is to assess the impact of cell phone radiation effects on green plants. Rapid progress in networking and communication systems has introduced frequency- and amplitude-modulated technologies to the world with higher allowed bands and greater speed by using high-powered radio generators, which facilitate high definition connectivity, rapid transfer of larger data files, and quick multiple accesses. These cause frequent exposure of cellular radiation to the biological world from a number of sources. Key factors like a range of frequencies, time durations, power densities, and electric fields were found to have differential impacts on the growth and development of green plants. As far as the effects on green plants are concerned in this review, alterations in their morphological characteristics like overall growth, canopy density, and pigmentation to physiological variations like chlorophyll fluorescence and change in membrane potential etc. have been found to be affected by cellular radiation. On the other hand, elevated oxidative status of the cell, macromolecular damage, and lipid peroxidation have been found frequently. On the chromosomal level, micronuclei formation, spindle detachments, and increased mitotic indexes etc. have been noticed. Transcription factors were found to be overexpressed in many cases due to the cellular radiation impact, which shows effects at the molecular level.

https://pubmed.ncbi.nlm.nih.gov/38773047/

Conclusion

Cellular radiation affects plant seed germination, physiological, biochemical, cytogenetic, and molecular characters. In some cases, the effect has been shown to be positive. There is definitely some impact of these radiations; still there are controversies. Negative impacts can be described as heating stress or

charged effect of electric fields, but how the changes are positive nobody explains. Where the positive changes happen or on what frequency, and if power and SAR value is required have not been described. Not much research was found on wild plants, despite having more height and highly exposed in field conditions. Molecular changes in level of mRNA transcriptions have been seen with very less numbers of publications. Changes in the level of ion transports and electron transport chains have been done in very less numbers. Physical absorption and behavior of plant tissues have to be done more widely. The importance of these studies is most crucial because of the increased demand of technology with more number of users and requirement of high-end communication. Vast research in this field is required in order to avoid further hazards.

--

Effect of moisture content variation on dielectric properties of various plant leaves at microwave frequencies

Chauhan PD, Gadani DH, Rana VA. Effect of moisture content variation on dielectric properties of various plant leaves at microwave frequencies. Sci Rep. 2024 Jun 8;14(1):13204. doi: 10.1038/s41598-024-64266-3.

Abstract

Complex permittivity of Corn, Jowar, Ashoka and Banana plant leaves was measured using a Vector Network Analyzer (Anritsu Shockline Model-MS46322A) over 500 MHz to 15 GHz frequency range, at different moisture contents. The dielectric constant (ε ') and dielectric loss (ε ”) of the leaves decrease with decrease in moisture content over this frequency range. For the leaves having moisture content less than certain critical moisture level (CML), the values of ε ' and ε " remain almost constant with frequency variation. ε ” does not increase appreciably with the increase in moisture content up to CML. CML is found to vary for different types of leaves. At higher moisture level, ε ' and ε ” exhibit frequency dependence. Above certain moisture level in the leaves, ε ” starts to increase slowly with the increase in frequency above 3 GHz approaching towards the dielectric relaxation of water. The values of ε ' and ε ” were also measured for the extract of leaves and compared with the values calculated using Stogryn equations for the same salinity, and the results agree with the calculated values. The measured values of ε ' and ε ” for the leaves were compared with the values calculated using Debye-Cole dual dispersion dielectric model and are found to match very well.

Open access paper: https://www.nature.com/articles/s41598-024-64266-3

--

How the Impact of Electromagnetic Fields on Plants Can Greatly Increase Severity of and Even Occurrence of “Wildfires”: A Four-Part Structure

Pall ML. How the Impact of Electromagnetic Fields on Plants Can Greatly Increase Severity of and Even Occurrence of “Wildfires”: A Four-Part Structure.Ecology & Conservation Science: Open Access. 2024. doi: 10.19080/ECOA.2023.04.555631

Abstract

Low growing plants near buildings and electric powerlines often burn explosively at extreme temperatures, leaving a light gray powder whereas plants away from such sources of electromagnetic fields (EMFs) burn more normally, leaving slightly burned and/or charred materials. It is argued here that EMFs trigger a four-part mechanism producing this unusual pattern. Electronically generated EMFs impact plants via activation of voltage-controlled calcium channels, causing increases in plant terpenes and three other classes of volatiles (polyamines, lipid peroxidation volatiles and methyl jasmonate) and large increases in peroxynitrite and reactive free radicals. Four processes ensue: 1. The volatile terpenes make plants much more flammable. 2. Each of the four classes of volatiles accumulate in the air under very low wind conditions, and the heavy volatile-containing air spreads over the ground, selectively producing plasma membrane depolarization in low growing plants and parts of plants. Depolarization activates the same calcium channels activated by EMFs, selectively spreading and amplifying EMF-like effects to these low growing plants. 3. The terpenes react with free radicals and O2 to produce three classes of explosive terpene-derived chemicals, hydroperoxides, nitrate esters and nitro compounds. Accordingly, the explosive burning of these three classes of terpene derived chemicals in low growing plants produces very rapidly advancing low level firestorms which when they hit adjacent buildings also causes the buildings to burn at extremely high temperatures, leaving a light gray powder. 4. Terpene hydroperoxides because of their low thermostability may cause spontaneous combustion. Appropriate conditions for spontaneous combustion may be limited to plant materials in depressions in the ground. Sixteen fire observations are inconsistent with climate change being the sole cause of fire severity but are consistent with the proposed mechanism. Wind records from four large explosive US fires were examined and were found to be consistent with prediction. High voltage powerline roles in fires may be caused by powerline dirty electricity produced EMFs rather than poor maintenance as has previously been claimed.

Open access paper: https://juniperpublishers.com/ecoa/pdf/ECOA.MS.ID.555631.pdf

--

Oxidative Stress Response of Honey Bee Colonies during Long-Term Exposure at a Frequency of 900 MHz under Field Conditions

Vilić M, Žura Žaja I, Tkalec M, Tucak P, Malarić K, Popara N, Žura N, Pašić S, Gajger IT. Oxidative Stress Response of Honey Bee Colonies (*Apis mellifera* L.) during Long-Term Exposure at a Frequency of 900 MHz under Field Conditions. Insects. 2024 May 20;15(5):372. doi: 10.3390/insects15050372.

Abstract

In this study, oxidative stress and lipid peroxidation in honey bee larvae, pupae and the midguts of adult bees were investigated during a one-year exposure to radiofrequency electromagnetic fields (RF-EMFs) at a frequency of 900 MHz under field conditions. The experiment was carried out on honey bee colonies at three locations with electric field levels of 30 mV m^{-1}, 70 mV m^{-1} and 1000 mV m^{-1}. Antioxidant enzymes, glutathione-S-transferase (GST), catalase (CAT) and superoxide dismutase (SOD) and thiobarbituric acid reactive substances (TBARS) as indicators of lipid peroxidation were measured spectrophotometrically. The GST activity within the same developmental stage showed no significant differences regardless of electric field level or sampling time. The highest GST activity was found in the pupae, followed by activity in the larvae and midguts. Both CAT activity and TBARS concentration were the highest in the midguts, regardless of field level and sampling time. The larvae showed a significantly

higher TBARS concentration at the location with an electric field level of 1000 mV m^{-1} compared to the locations with lower levels. Our results show that RF-EMFs at a frequency of 900 MHz can cause oxidative stress in honey bees, with the larval stage being more sensitive than the pupal stage, but there was no linear relationship between electric field level and effect in any of the developmental stages.

Open access paper: https://www.mdpi.com/2075-4450/15/5/372

--

The Cellular Response Is Determined by a Combination of Different ELF-EMF Exposure Parameters: A Scope Review

López de Mingo I, Rivera González M-X, Maestú Unturbe C. The Cellular Response Is Determined by a Combination of Different ELF-EMF Exposure Parameters: A Scope Review. *International Journal of Molecular Sciences*. 2024; 25(10):5074. doi: 10.3390/ijms25105074.

Abstract

Since the establishment of regulations for exposure to extremely low-frequency (0–300) Hz electromagnetic fields, scientific opinion has prioritised the hypothesis that the most important parameter determining cellular behaviour has been intensity, ignoring the other exposure parameters (frequency, time, mode, waveform). This has been reflected in the methodologies of the in vitro articles published and the reviews in which they are included. A scope review was carried out, grouping a total of 79 articles that met the proposed inclusion criteria and studying the effects of the different experiments on viability, proliferation, apoptosis, oxidative stress and the cell cycle. These results have been divided and classified by frequency, intensity, exposure time and exposure mode (continuous/intermittent). The results obtained for each of the processes according to the exposure parameter used are shown graphically to highlight the importance of a good methodology in experimental development and the search for mechanisms of action that explain the experimental results, considering not only the criterion of intensity. The consequence of this is a more than necessary revision of current exposure protection regulations for the general population based on the reductionist criterion of intensity.

Conclusions

This review arises from the need to consider each of the parameters of exposure to magnetic fields as an entity capable of producing a determined response in the cell. The importance that has been given to the intensity parameter since the establishment of regulations for the exposure of the general public to low-frequency electromagnetic fields has led to it also being the main parameter in in vitro experiments, considering intensity as the determining parameter for the occurrence of the main cellular effects and placing the “dose effect” as one of the main theories explaining the effects found. We must begin to think of combinations of exposure parameters that act at the cellular level as specific codes that give rise to specific responses. Biology, and more specifically cell biology, entails a set of rules that must be protected as a fundamental right. There must be a commitment on the part of scientific researchers in bioelectromagnetism to develop quality experiments based on rigorous working methodologies and the search for experimental conclusions that contribute to existing knowledge in this field in order to establish models of interaction between magnetic fields and biological systems that allow us to think of

magnetic fields not only as a tool for modulating cell behaviour but also as a possible therapeutic application for various fields of medicine such as oncology, neurodegeneration or the healing of wounds and lesions.

Open access paper: https://www.mdpi.com/1422-0067/25/10/5074

--

Modulation of Zebrafish Heart Rate by Alternating Magnetic Fields with Frequencies Close to Heart Rhythm

Krylov VV, Sizov DA, Machikhin AS, Guryleva AV, Tchougounov V, Burlakov AB. Modulation of Zebrafish Heart Rate by Alternating Magnetic Fields With Frequencies Close to Heart Rhythm. *IEEE Journal of Electromagnetics, RF and Microwaves in Medicine and Biology*. 2024. doi: 10.1109/JERM.2024.3397557.

Abstract

Extremely low-frequency magnetic fields (ELF-MF) up to 100 μT exhibit impacts on physiological processes, including heart function. The mechanisms underlying the influence of these fields on fish heart rates remain insufficiently explored. We assumed that the direct impact of ELF-MF with a frequency close to the heart rate could entrain oscillatory processes responsible for autonomously maintaining heart rhythm in zebrafish embryos. Embryos' heart rates ranged from 1.44 to 3 Hz depending on age, and ELF-MF with frequencies precisely matched, 10% higher, or lower than the heart rate were applied. Additionally, embryos experienced ELF-MF with amplitudes varying by an order of magnitude. Almost all tested ELF-MF induced an increased heart rate effect. This effect was the most pronounced when the exposure occurred earlier during ontogenesis. Fields with frequencies close to the heart rate did not entrain cardiac contractions in zebrafish embryos. A significant negative correlation between heart rate increase and ELF-MF frequency was observed for ELF-MF with amplitudes of 1.98–3.2 μT and 46.8 μT but not 30 μT. Probable molecular mechanisms underlying these effects are discussed in terms of magnetic influence on radical pairs within biochemical oscillating processes.

https://ieeexplore.ieee.org/document/10529610

--

Prey can detect predators via electroreception in air

England SJ, Robert D. Prey can detect predators via electroreception in air. Proc Natl Acad Sci U S A. 024 Jun 4;121(23):e2322674121. doi: 10.1073/pnas.2322674121.

Abstract

Predators and prey benefit from detecting sensory cues of each other's presence. As they move through their environment, terrestrial animals accumulate electrostatic charge. Because electric charges exert forces at a distance, a prey animal could conceivably sense electrical forces to detect an approaching predator. Here, we report such a case of a terrestrial animal detecting its predators by electroreception. We show that predatory wasps are charged, thus emit electric fields, and that caterpillars respond to such fields with defensive behaviors. Furthermore, the mechanosensory setae of caterpillars are deflected by these electrostatic forces and are tuned to the wingbeat frequency of their insect

predators. This ability unveils a dimension of the sensory interactions between prey and predators and is likely widespread among terrestrial animals.

Open access paper: https://www.pnas.org/doi/10.1073/pnas.2322674121

--

Associations between solar and geomagnetic activity and cognitive function in the Normative Aging study

Liddie JM, Vieira CLZ, Coull BA, Sparrow D, Koutrakis P, Weisskopf MG. Associations between solar and geomagnetic activity and cognitive function in the Normative Aging study. Environ Int. 2024 Apr 16;187:108666. doi: 10.1016/j.envint.2024.108666.

Abstract

Background: Studies show that changes in solar and geomagnetic activity (SGA) influence melatonin secretion and the autonomic nervous system. We evaluated associations between solar and geomagnetic activity and cognitive function in the Normative Aging Study from 1992 to 2013.

Methods: We used logistic and linear generalized estimating equations and regressions to evaluate the associations between moving averages of sunspot number (SSN) and Kp index (a measure of geomagnetic activity) and a binary measure for Mini-Mental State Examination (MMSE) scores (≤25 or > 25) and six other cognitive tests as continuous measures, combined into one global composite score and considered separately.

Results: A one-IQR increase in same-day SSN and Kp index were associated with 17% (95% CI: 3%, 34%) and 19% (95% CI: 4%, 36%) increases in the odds of low MMSE score. We observed small increases in the global cognitive score with increasing SSN, although we observed decreases specifically in relation to the backwards digit span test.

Conclusions: Periods of high SGA were associated with cognitive function. SGA may not equally impact all aspects of cognitive function, as evidenced by differences in associations observed for the MMSE, global cognitive score, and individual cognitive tests. Given that much of the pathology of cognitive decline in the elderly remains unexplained, studies specifically targeting decline and with longer follow-up periods are warranted.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0160412024002526?via%3Dihub

--

ICNIRP Statement on Short Wavelength Light Exposure from Indoor Artificial Sources and Human Health

Miller S, Cajochen C, Green A, Hanifin J, Huss A, Karipidis K, Loughran S, Oftedal G, O'Hagan J, Sliney DH, Croft R, van Rongen E, Cridland N, d'Inzeo G, Hirata A, Marino C, Röösli M, Watanabe S; International Commission on Non-Ionizing Radiation Protection (ICNIRP). ICNIRP Statement on Short Wavelength Light Exposure from Indoor Artificial Sources and Human Health. Health Phys. 2024 Apr 1;126(4):241-248. doi: 10.1097/HP.0000000000001790.

Abstract

Concerns have been raised about the possibility of effects from exposure to short wavelength light (SWL), defined here as 380-550 nm, on human health. The spectral sensitivity of the human circadian timing system peaks at around 480 nm, much shorter than the peak sensitivity of daytime vision (i.e., 555 nm). Some experimental studies have demonstrated effects on the circadian timing system and on sleep from SWL exposure, especially when SWL exposure occurs in the evening or at night. The International Commission on Non-Ionizing Radiation Protection (ICNIRP) has identified a lack of consensus among public health officials regarding whether SWL from artificial sources disrupts circadian rhythm, and if so, whether SWL-disrupted circadian rhythm is associated with adverse health outcomes. Systematic reviews of studies designed to examine the effects of SWL on sleep and human health have shown conflicting results. There are many variables that can affect the outcome of these experimental studies. One of the main problems in earlier studies was the use of photometric quantities as a surrogate for SWL exposure. Additionally, the measurement of ambient light may not be an accurate measure of the amount of light impinging on the intrinsically photosensitive retinal ganglion cells, which are now known to play a major role in the human circadian timing system. Furthermore, epidemiological studies of long-term effects of chronic SWL exposure per se on human health are lacking. ICNIRP recommends that an analysis of data gaps be performed to delineate the types of studies needed, the parameters that should be addressed, and the methodology that should be applied in future studies so that a decision about the need for exposure guidelines can be made. In the meantime, ICNIRP supports some recommendations for how the quality of future studies might be improved.

Open access paper: https://journals.lww.com/health-physics/fulltext/2024/04000/icnirp_statement_on_short_wavelength_light.7.aspx

--

Mobile telephony radiation exerts genotoxic action and significantly enhances the effects of gamma radiation in human cells

Panagopoulos DJ. Mobile telephony radiation exerts genotoxic action and significantly enhances the effects of gamma radiation in human cells. Gen Physiol Biophys. 2024 Mar;43(2):103-120. doi: 10.4149/gpb_2023036.

Abstract

I previously reported chromosomal damage in human peripheral blood lymphocytes (HPBLs) induced by: a) Mobile telephony (MT) electromagnetic fields (EMFs)/electromagnetic radiation (EMR), b) a high caffeine dose, and c) the combination of the two stressors. HPBLs from the same subjects exposed to gamma radiation at doses 0.1, 0.3, or 0.5 Gy, displayed more aberrations than those exposed to MT EMFs or the high caffeine dose in a dose- dependent manner. When the cells exposed to these gamma radiation doses were pre-exposed to a single 15-min MT EMF exposure, the number of aberrations

increased significantly more than the sum number of aberrations induced by the individual stressors in all subjects. Thus, MT EMF exposure at a power density ~ 136 times below the latest International Commission on Non-Ionizing Radiation Protection-ICNIRP exposure limit, apart from the fact that it is genotoxic by itself, significantly enhanced the genotoxic action of gamma radiation. Since gamma radiation at similar doses is applied for diagnostic and therapeutic purposes, people should be aware of the increased risk during treatment periods. Comparison of the genotoxic action between MT EMF and gamma radiation shows that the ICNIRP limits are, at least, $\sim 4.5\times10^4$ times less stringent than the limits for gamma radiation.

Excerpts

The studies that found real-life UMTS (3G/4G) exposure to be even more genotoxic than real-life GSM (2G) (D'Silva et al. 2017, 2021) are in line with the fact that newer types of MT/WC EMFs (3G, 4G, 5G) transmit increasingly higher amount/density of variable information (speech, text, images, video, Internet) making the signal increasingly complicated, unpredictably varying each moment, and thus, increasingly more bioactive due to the inability of the living organisms to adapt to a highly variable stressor. Thus, **the effects under deployment 5G MT EMF are expected to be even more intense than those of 2G, 3G, 4G**. This should have been seriously considered by the responsible public health authorities....

The disruption of cell electrochemical balance by manmade (polarized and coherent) EMFs through irregular gating of voltage-gated ion channels (VGICs) in cell membranes is described by the **"ion forced-oscillation and VGIC dysfunction" mechanism (IFO-VGIC mechanism)** (Panagopoulos et al. 2000, 2002, 2015b, 2021; Panagopoulos 2022b). According to this mechanism, the mobile ions in the cells are forced to oscillate in parallel and in phase with the applied man-made oscillating EMFs and this coordinated oscillation of electric charge exerts constructive Coulomb forces on the channel sensors of the VGICs similar to those exerted by membrane voltage changes that physiologically gate the VGICs. This causes irregular gating, and thus, dysfunction of the VGICs, which leads to intracellular release of ROS that finally cause genetic/cellular damage (Panagopoulos et al. 2021, 2022b)....

What has been referred to by Pall (2018) as voltage-gated calcium channel activation mechanism ("VGCC activation mechanism") is no other than the IFO-VGIC mechanism specifically on the calcium voltage-gated ion channels, and should not be reported as a different mechanism. Pall claimed he suggested a different mechanism simply because he hypothesized that the VGICs are gated by "direct" forces on their voltage-sensors by "penetrating" RF EMFs instead of ELF forces exerted by the oscillating ions in close proximity to the sensors. The impossibility of Pall's claims is analyzed in commentary papers (Foster and Balzano 2021; Panagopoulos 2021; Arribas et al. 2022).

Conclusions:

(1) MT EMF exposure, apart from the fact that it is genotoxic by itself, significantly enhanced the genotoxic action of gamma radiation in combined exposure;

(2) People/patients who are subjected to diagnostic or therapeutic treatment with ionizing radiation should be prudently advised to avoid using Wireless Communication (WC) devices (mobile/smart phones, wi-fi, cordless domestic phones, etc.) for a few days before, during, and after such treatments;

(3) Medical/radiology practitioners should be specifically educated on the risks of anthropogenic EMF-exposures in addition to those of ionizing radiations;

(4) Comparison with caffeine and gamma radiation suggests that the ICNIRP (2020) limits for WC EMF exposure should be lowered by 40,000 and 45,000 times respectively;

(5) The limit for short-term (acute) exposure should then become 0.1 µW/cm2 and accordingly for long- term exposure 0.001 µW/cm2;

(6) The combined effects of real-life man-made EMFs with a variety of other environmental stressors should be examined as a priority by next studies.

Open access paper:
http://www.elis.sk/download_file.php?product_id=8263&session_id=ef2ijbhbn0hevd1s33q2nvkrp6

--

Effect of exposure to RF-EMF from lab simulator of 5G NR base station on physiological parameters & cognitive abilities of male rats of different ages

(My note: Methodological limitations of this study reduce generalizability to real-world 5G exposures.)

Krivova NA, Kudabaeva MS, Zaeva OB *et al.* The effect of exposure to RF-EMF from the laboratory simulator of 5G NR base station on physiological parameters and cognitive abilities of male wistar rats of different ages. *Sci Rep* **14**, 10283 (2024). doi: 10.1038/s41598-024-60862-5.

Abstract

In this article, the impact of radiofrequency electromagnetic field (RF-EMF) exposure from a simulated base station for the 5G New Radio (5G NR) telecommunication on rats was studied. The base station affects all age groups of the population, thus, for the first time, the experiment was conducted on male Wistar rats of three different ages (juvenile, adult, and presenile). The base station exposure parameters were chosen according to ICNIRP recommendations for limiting the exposure to radiofrequency electromagnetic field: frequency 2.4 GHz with an average specific absorption rate of 0.0076 W/kg and 0.0059 W/kg over the whole body of experimental animals. Throughout the experiment, body weight was examined weekly, and the dynamics of body weight gain was monitored. Rectal and skin surface temperature on the right hind limb was monitored weekly. Testing in the Morris water maze was performed during the last, Week 5, of RF-EMF exposure. After euthanasia, organ weights were determined in experimental and control animals. None of the investigated parameters did show any statistically significant differences between exposed and control animals of the same age. The data obtained can be used to assess the possible consequences of chronic exposure to RF-EMF from 5G NR base stations.

Open access paper: https://www.nature.com/articles/s41598-024-60862-5

--

Human exposure to EMF from 5G base stations: analysis, evaluation and comparison of different assessment methods

Expósito I, Hakizimali C, Sánchez MG, Cuiñas I, Verhaevert J. Human exposure to EMF from 5G base stations: analysis, evaluation and comparison of different assessment methods. Measurement, Volume 229, 2024. doi: 10.1016/j.measurement.2024.114434.

Highlights

This paper analyzes the feasibility of assessing the 5G base stations compliance using broadband field probes and compares their performance with alternative methodologies and equipment.

Performance of three different methodologies and equipment (broadband probes, spectrum analyzers, and drive test scanners), in the context of human exposure to electromagnetic fields (EMF) from 5G base stations, is compared.

Quantification of the uncertainty that the fluctuation in 5G signal levels induces in the assessment of electromagnetic fields exposure is provided.

The use of broadband field probes for 5G exposure assessment is still possible under certain considerations and correcting the results considering the base station load and beamforming effects.

Abstract

5G networks deployment poses new challenges when evaluating human exposure to electromagnetic fields. Fast variation of the user load and beamforming techniques may cause large fluctuations of 5G base stations field level. They may be underestimated, resulting in compliance of base stations not fitting the requirements. Apparently, broadband field meters would not be adequate for measuring such environments. However, we analyze the feasibility of confidently using broadband field meters and compare their performance with alternative equipment. Measurements based on the synchronization signals power level, using spectrum analyzers or drive test scanners, may be valid, if gain differences between the signaling and data radiation patterns are characterized. These methods lead to good results but require more time and knowledge. Nevertheless, using broadband field meters is still possible if the measurement results are corrected considering the base station load. Under specific conditions, explained here, fast assessment of 5G compliance could be provided.

Conclusions

Assessing human exposure to an electromagnetic field in presence of a 5G base station is not an easy task. The implementation of M−MIMO techniques in 5G base stations results in adaptive beamforming. This makes difficult to guarantee that the field levels are at their maximum at the measurement location during the complete measurement period, which would limit the applicability of broadband instruments as having been done for previous generations. In this research, we have compared different methods for 5G exposure assessment, using a broadband field meter with an isotropic probe, a spectrum analyzer and a drive test scanner.

Along the paper, we first give an overview of the 5G signal structure, describing the frequency domain and time domain specifications. Afterwards, possible assessment methods are described. The SSB level is measured using the Keysight FieldFox N9913A SA and the Rohde & Schwarz TSM6 DTS. The values are extrapolated to the worst-case exposure and compared to the measurements done with the

Wavecontrol WPF8 broadband field probe. Measurements are repeated increasing the base station load by performing a heavy download from a 5G user terminal located near the testers.

The proposed methods were field tested at the University of Vigo, Spain, with a commercial 5G base station located on its campus. The measurements were performed at 7 locations in LOS conditions around the base station, gathering data with the three different equipment at the same locations and at the same time. This data collection allows the comparison of the three methodologies under the same radiating conditions.

All results have been analyzed considering the specific measurement uncertainties, which allows a deeper and more precise comparison among them.

From the measurement results, we can extract that the exposure levels are low at this stage of the 5G deployment. When loading the base station, the results showed that using the broadband field meter can overestimate the field level. Thus, it is still a useful method to check if the field levels comply with the regulation in human exposure; very simple and cost-effective compared with others. In-situ measurements of human exposure to EMF have to be practical and easy to carry, involving only the resources and equipment strictly necessary, but without compromising the validity of the results. When the reference levels are surpassed, more accurate methods based in the assessment and extrapolation of the SSB level could be a solution. The drawback is the required post processing, specially correcting the gain difference between SSB and data signals. If not provided by the network operator, this difference can be determined through measurements, as explained along this document. Measuring with an SA in max-hold mode in the bandwidth of the SSB does not work in 5G as it does in LTE, as we cannot be sure if the measured level corresponds to the SSB or to the user data, no matter if we are forcing the load of the station or not.

The analysis of the results demonstrate that broadband instruments can be used for assessing human exposure to EMF in the vicinity of 5G base stations, which radiating elements provide fields with extreme fluctuations in their intensity as a function of the system load and beamforming configuration. This is accurate when measurements are done by forcing an extra load of the station and the pointing of an antenna beam towards the probe. The validation of this fast method as a first attempt to assess the compliance of 5G stations permits the testing of these base stations in an efficient way. Only when broadband instrument results (including their uncertainties) would overpass the reference levels, a more detailed analysis would be necessary, which procedure and tips are also depicted along this paper.

https://www.sciencedirect.com/science/article/pii/S0263224124003191

--

Absorption of 5G sub-6 GHz electromagnetic radiation from base station to male reproduction system

Lin J, Li J, Ding G. Absorption of 5G sub-6 GHz electromagnetic radiation from base station to male reproduction system. Int J Radiat Biol. 2024 May 6:1-8. doi: 10.1080/09553002.2024.2347354.

Abstract

Background: The impact of electromagnetic radiation from communication on the male reproductive system has emerged as a significant concern in public health. A notable distinction of the 5G sub-6 GHz

band, compared to traditional 2G, 3G, and 4G frequency bands, is the inclusion of higher frequency bands. This has raised public concerns regarding the potential effects of these higher frequencies on organisms, particularly their reproductive systems. While it is imperative to investigate the biological effects and potential risks associated with these new frequency bands in laboratory settings, comparing and evaluating differences between various frequency bands remain challenging due to the absence of standardized parameters such as exposure conditions and duration. In contrast, dose assessment offers a simpler and more reliable approach.

Materials and methods: The dose assessment method was employed in this study to investigate the risks associated with sub-6 GHz electromagnetic radiation from 5G base stations on the male reproductive system. A classical human body model (Duke) was utilized, and an electromagnetic simulation environment was established based on the actual polarization direction of the exposed base stations and various body postures. This research explored the effects of field direction, posture, public population, and frequency on the specific absorption rate of the reproductive system.

Results and conclusions: While maintaining the same level of exposure, a higher frequency results in a reduced dosage on reproductive system. Further analysis reveals that, considering the public exposure threshold, the employment of higher frequency bands in 5G sub-6 GHz does not present a greater dosage on reproductive system compared to lower frequency bands. Consequently, with regard to dosage, there is no need for excessive concern among the general public regarding the impact of electromagnetic radiation emitted by 5G base stations operating below 6 GHz on male reproductive health.
https://pubmed.ncbi.nlm.nih.gov/38709715/

Excerpts

The normalized evaluation revealed a strong negative correlation between the RSSAR results and frequency (figures 6 and 8), indicating the reproductive system's response to varying frequencies of electromagnetic waves. It also indicates that high-frequency skin effects are particularly prominent in the reproductive system. In the exposure limit assessment, there was an evident inflection point at 2.2 GHz where the trend of RSSAR values with frequency changes reached its maximum peak (figure 10). Therefore, when considering the exposure threshold, the higher frequency bands in 5G did not show higher dose values on RSSAR, but on the contrary, higher dose values were obtained in the 4G bands (such as 2.2 GHz in this study). These findings suggest that the potential risk of higher frequency electromagnetic waves in 5G sub-6 GHz is lower, as indicated by both normalized and exposure limit assessments.

Restriction and future focus

This research also has certain limitations. The exposure environment has not been considered in the context of the actual 5G exposure environment. With the introduction of 5G base stations, the grid of base stations may become denser and the energy of a single beam of electromagnetic radiation may be higher (Sofri et al. Citation2022). Although reference (Celaya-Echarri et al. Citation2020) simulated and compared the exposure values caused by base stations in the 2G–5G frequency band, the electric field conditions in real communities should also be measured and compared in detail. In future work,

exposure measurements and dose assessing could be combined with epidemiological investigations to further explore the effects of new frequency bands on reproductive and other systems.

Conclusions

The absorbed dose level of electromagnetic radiation emitted by communication base stations on the male reproductive system was analyzed, with a specific focus on the primary frequency points within the sub-6 GHz frequency band. The study revealed that higher frequencies resulted in reduced dose values while maintaining equivalent exposure levels. Further analysis demonstrated that, considering the exposure threshold, the utilization of higher frequencies in 5G sub-6 GHz did not present a greater risk compared to lower frequencies. Therefore, from a dosage assessment perspective, concerns regarding the impact of sub-6 GHz electromagnetic radiation emitted by 5G base stations on male reproductive health are unwarranted.

--

Central Causation of Autism/ASDs via Excessive [Ca^{2+}]i Impacting Six Mechanisms Controlling Synaptogenesis during the Perinatal Period: The Role of Electromagnetic Fields and Chemicals and the NO/ONOO(-) Cycle, as Well as Specific Mutations

Pall ML. Central Causation of Autism/ASDs via Excessive [Ca2+]i Impacting Six Mechanisms Controlling Synaptogenesis during the Perinatal Period: The Role of Electromagnetic Fields and Chemicals and the NO/ONOO(-) Cycle, as Well as Specific Mutations. Brain Sciences. 2024; 14(5):454. doi: 10.3390/brainsci14050454. (Published: 30 April 2024)

Abstract

The roles of perinatal development, intracellular calcium [Ca2+]i, and synaptogenesis disruption are not novel in the autism/ASD literature. The focus on six mechanisms controlling synaptogenesis, each regulated by [Ca2+]i, and each aberrant in ASDs is novel. The model presented here predicts that autism epidemic causation involves central roles of both electromagnetic fields (EMFs) and chemicals. EMFs act via voltage-gated calcium channel (VGCC) activation and [Ca2+]i elevation. A total of 15 autism-implicated chemical classes each act to produce [Ca2+]i elevation, 12 acting via NMDA receptor activation, and three acting via other mechanisms. The chronic nature of ASDs is explained via NO/ONOO(-) vicious cycle elevation and MeCP2 epigenetic dysfunction. Genetic causation often also involves [Ca2+]i elevation or other impacts on synaptogenesis. The literature examining each of these steps is systematically examined and found to be consistent with predictions. Approaches that may be used for ASD prevention or treatment are discussed in connection with this special issue: The current situation and prospects for children with ASDs. Such approaches include EMF, chemical avoidance, and using nutrients and other agents to raise the levels of Nrf2. An enriched environment, vitamin D, magnesium, and omega-3s in fish oil may also be helpful.

Excerpts

Figure 1 is an outline of the mechanisms proposed here and, in some cases, elsewhere, to act during the perinatal period of brain development [5], to disrupt synaptogenesis and therefore cause autism spectrum disorders (ASDs). Accordingly, key factors in autism/ASD causation are proposed to be the six

mechanisms in the lower left part of Figure 1 (in red), each of which have been shown to have important roles in synaptogenesis in the developing brain. These six mechanisms are neuronal migration, dendritic outgrowth, synapse formation, synapse maturation, synaptic pruning, and lowered MeCP2 function. It is shown below that these six mechanisms are regulated by intracellular calcium ([Ca2+]i).

VGCC activity acts via [Ca2+]i to produce diverse biological effects via the pathways shown in Figure 2

.... Let us shift now to ASD treatment and prevention. These are presented here not to give medical advice, but rather to enlighten the reader in the central issues of this special issue: The current situation and prospects for children with ASDs. Both treatment and prevention require avoidance of electronically-generated EMFs and diverse chemicals, however difficult that may be. Let us start with EMFs. There are many sources of EMFs in our living spaces, workplaces, educational and recreational environments, and even outdoors. Many of these are discussed in Section 2.1.

There are “safety guidelines” initially proposed by ICNIRP, an organization set up by the industry and these ICNIRP safety guidelines have been adopted, sometimes with minor modifications, over much of the world. The ICNIRP and similar “safety guidelines” are all based on thermal effects produced over either a 6 min or 30 min period. However, as discussed above, biological effects are caused by electric forces or time-varying magnetic forces produced by electronically-generated EMFs, and are unrelated to thermal effect “safety guidelines”. It follows that all assurances of safety based on these “safety guidelines” have no scientific merit.

Pregnant women should not use cell phones, cordless phones, should avoid Wi-Fi fields, and should use wired connections to the internet, avoid Bluetooth, should not spend time near smart meters, cell phone towers, avoid high voltage powerlines, etc. Pregnant women should avoid being in buildings with high dirty electricity in their electric powerlines. Such avoidance is very challenging. A second choice is to use shielding to protect the fetus and the young baby after birth.

You can get large shielded T-shirts to protect the fetus, wearing a regular cloth T-shirt underneath to minimize body contact. Such shielding materials have large numbers of metal (often silver) fibers. I believe that shielding mostly acts by disrupting the coherence of the EMFs, therefore greatly lowering any electric or magnetic forces. Shielding has been extensively studied in the scientific literature [271,272,273,274,275,276,277]. For example, ref. [271] states that “By using wireless technologies, we increase the influence of the electromagnetic field on the human body. The basic measures for the protection against electromagnetic radiation are: (1) shielding of the radiation source, (2) shielding of the workplace, (3) personal protective equipment, (4) reduction of radiation in the immediate vicinity of the radiation source”. [271] goes on to discuss the shielding effects of both fabrics containing tiny metal fibers and also brick walls.

You can buy shielding paint of two different types to protect yourself in your house and, if you live in an apartment, this may be the only way to protect yourself from your neighbors’ EMFs. Shielding paint can be either graphite (black) paint which can be painted over with other colors. Other more pleasant colored shielding paint can be purchased containing tiny metallic particles. You can also buy open shielding cloth to put over windows, making them into lace like curtains to lower EMF exposures coming through the windows. Similar approaches can be used after birth to protect the young baby. It is highly desirable to purchase a good EMF meter to provide information and prevent the worst exposures throughout the perinatal period.

Open access paper: https://www.mdpi.com/2076-3425/14/5/454

--

The effects of radiofrequency electromagnetic fields exposure on human self-reported symptoms: A systematic review of human experimental studies (WHO SR8)

(Note: Methodological problems with many studies included in this systematic review undermine the validity of this review paper.)

Bosch-Capblanch X, Esu E, Oringanje CM, Dongus S, Jalilian H, Eyers J, Auer C, Meremikwu M, Röösli M. The effects of radiofrequency electromagnetic fields exposure on human self-reported symptoms: A systematic review of human experimental studies. Environment International. doi: 10.1016/j.envint.2024.108612. Available online 2 April 2024.

Abstract

Background The technological applications of radiofrequency electromagnetic fields (RF-EMF) have been steadily increasing since the 1950s exposing large proportions of the population. The World Health Organization (WHO) is assessing the potential health effects of exposure to RF-EMF.

Objectives To systematically assess the effects of exposure to RF-EMF on self-reported non-specific symptoms in human subjects and to assess the accuracy of perception of presence or absence of RF-EMF exposure.

Methods Eligibility criteria: experimental studies carried out in the general population and in individuals with idiopathic environmental intolerance attributed to EMF (IEI-EMF), in any language.

Information sources: Medline, Web of Science, PsycInfo, Cochrane Library, Epistemonikos and Embase and consulted the EMF portal, searched till April 2022.

Risk of bias: we used the Risk of Bias (RoB) tool developed by OHAT adapted to the topic of this review.

Synthesis of results: we synthesized studies using random effects meta-analysis and sensitivity analyses where appropriate.

Results Included studies: 41 studies were included, mostly cross over trials and from Europe, with a total of 2,874 participants.

Synthesis of results: considering the primary outcomes, we carried out meta-analyses of 10 exposure-outcomes pairs. All evidence suggested no or small non-significant effects of exposure on symptoms with high (three comparisons), moderate (four comparisons), low (one comparison) or very low (two comparisons) certainty of evidence. The effects (standard mean difference, where positive values indicate presence of symptom being exposed) in the general population for head exposure were (95% confidence intervals) 0.08 (−0.07 to 0.22) for headache, −0.01 (−0.22 to 0.20) for sleeping disturbances and 0.13 (−0.51 to 0.76) for composite symptoms; and for whole-body exposure: 0.09 (−0.35 to 0.54), 0.00 (−0.15 to 0.15) for sleeping disturbances and −0.05 (−0.17 to 0.07) for composite symptoms. For IEI-EMF individuals SMD ranged from −0.19 to 0.11, all of them with confidence intervals crossing the value

of zero. Further, the available evidence suggested that study volunteers could not perceive the EMF exposure status better than what is expected by chance and that IEI-EMF individuals could not determine EMF conditions better than the general population.

Discussion Limitations of evidence: experimental conditions are substantially different from real-life situations in the duration, frequency, distance and position of the exposure. Most studies were conducted in young, healthy volunteers, who might be more resilient to RF-EMF than the general population. The outcomes of interest in this systematic review were symptoms, which are self-reported. The available information did not allow to assess the potential effects of exposures beyond acute exposure and in elderly or chronic ill people. It cannot be ruled out that a real EMF effect in IEI-EMF groups is masked by a mix with insensitive subjects. However, studies on symptom reporting and/or field perception did not find any evidence that there are particularly vulnerable individuals in the IEI-EMF group, although in open provocation studies, when volunteers were informed about presence or absence of EMF exposure, such differences have been consistently observed.

Interpretation: available evidence suggests that acute RF-EMF below regulatory limits does not cause symptoms and corresponding claims in everyday life are related to perceived and not real EMF exposure status.

Excerpts

4.4. Implications for practice

We could not find any evidence that would support an association between exposure to RF-EMF below regulatory recommendations and a large variety of short term reported symptoms; neither evidence that could suggest the capacity of individuals to detect the exposure to RF-EMF. This is the case for the general population, for IEI-EMF individuals, in the main meta-analyses as well as in the exposure–response and sensitivity analyses.

While these findings cannot rule out other long term potential harms of RF-EMF, they suggest that the presence or absence of common symptoms may not become good clinical predictors of the occurrence or evolution of IEI-EMF. It may also indicate that notion of acute effects from RF-EMF below regulatory limits in the everyday environment are better explained with the nocebo or the attribution hypothesis (Dieudonné, 2020).

4.5. Implications for research

The body of evidence in this review seems to consistently suggest the lack of association between EMF exposure and symptoms, in a variety of conditions and populations. The most critical aspect challenging the certainty of the evidence was the limited detail in the methodology of included studies, particularly in the issues related to the exposure randomisation and allocation concealment. While we hardly downgrades the evidence on the grounds of ‘indirectness’, most of the studies were focusing on adults, young adults and even adolescents, which may have limited the external validity of findings.

In order to confirm the findings of this systematic review, future research should, thus, consider adhering to best reporting practices and considering a wider scope of participants, including, for example, the elderly.

Future research should also be carried out standardising the design of experiments in terms of exposure ranges, timing and sequence of events, so as to make studies more comparable. It may be worth to increase exposure levels above regulatory limits for the general population to explore potential critical threshold for non-specific symptoms. Self-evident such studies need to follow a prudent approach to ensure that any symptoms are transient and do not result in a long term health impairment.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024001983

--

Is the sustainability of exposure to non-ionizing electromagnetic radiation possible?

Calvente I, Núñez MI. Is the sustainability of exposure to non-ionizing electromagnetic radiation possible? Medicina Clínica (English Edition), 162(8):387-393. 2024. doi: 10.1016/j.medcle.2023.11.016.

Abstract

Technological advances imply an increase in artificially generating sources of electromagnetic fields (EMF), therefore, resulting in a permanent exposure of people and the environment (electromagnetic pollution). Inconsistent results have been published considering the evaluated health effects. The purpose of this study was to review scientific literature on EMF to provide a global and retrospective perspective, on the association between human exposure to non-ionizing radiation (NIR, mainly radiofrequency-EMF) and health and environmental effects. Studies on the health effects of 5G radiation exposure have not yet been performed with sufficient statistical power, as the exposure time is still relatively short and also the latency and intensity of exposure to 5G. The safety standards only consider thermal effects, do not contemplate non-thermal effects. We consider relevant to communicate this knowledge to the general public to improve education in this field, and to healthcare professionals to prevent diseases that may result from RF-EMF exposures.

Excerpts

ICNIRP reviews safety guidelines but does not provide solutions, uses SAR, only considers thermal effects, does not consider non-thermal effects. Scientific evidence shows the importance of considering both thermal and non-thermal effects, short and long-term biological effects, and other factors. On the other hand, more frequent periodic reviews of the regulations are needed. According to ICNIRP the last one was in 2020. Even considering the leap in technology use in 2020, should we wait as many years as now (approx. 10) for a new update?

There are countries with stricter regulations than those established by IARC (sic) that are more concerned about the effects of this type of exposure (Italy, Belgium, Austria (Salzburg), Switzerland, Russia, China)....

In the face of such divergent assessments of wireless RF radiation, the practice of ALARA – as low as reasonably achievable – for RF health and safety should be followed.

On the basis of the evidence presented here, is it possible to speak of sustainability in relation to electromagnetic fields?

https://www.sciencedirect.com/science/article/pii/S2387020624001268

--

Environmental Factors as the Main Hormonal Disruptors of Male Fertility

Wdowiak N, Wójtowicz K, Wdowiak-Filip A, Pucek W, Wróbel A, Wróbel J, Wdowiak A. Environmental Factors as the Main Hormonal Disruptors of Male Fertility. J Clin Med. 2024 Mar 29;13(7):1986. doi: 10.3390/jcm13071986.

Abstract

Introduction and objective: Many scientific reports confirm a systematic decline in male semen parameters over the last decades. This phenomenon has been observed in all parts of the world, and its occurrence is associated, among others, with the hazardous effects of some environmental factors. The environmental factors for which the adverse effect on male fertility has been proven include water, air, and soil pollution, as well as electromagnetic fields and ionizing radiation. The aim of this article was the evaluation of the effect of selected environmental factors on male reproductive capacity based on an analysis of the current scientific reports.

Review methods: A systematic literature review was carried out using three databases: PubMed, EMBASE, and Scopus. The search was limited to the period from 2015 until the end of December 2023. Brief description of the state of knowledge: Environmental factors, such as heavy metals, tobacco smoke, pesticides, dioxins, furans, phthalates, and bisphenols, are well-tested substances that exert an adverse effect on male fertility. A harmful effect of electromagnetic fields and water and air pollution on reproductive functions may be expected; however, this has not been fully proven.

Summary: Results obtained by many researchers published to date should evoke great concern regarding the quality of the environment in which we live, as well as fears about the effect of environmental factors not only on male fertility, but also on all aspects of human health. The majority of environmental pollutants affect the male body by causing oxidative stress and through their effect on the endocrine system.

Excerpt

12. Electromagnetic Fields

Electromagnetic radiation (EMR) originating from both the natural environment and human activity constantly affects living organisms. The effect of EMR on the reproductive system may occur as a result of the thermal effect, generation of oxidative stress, changes in the structure of proteins, and causing changes in ion transport through cell membranes; however, these pathomechanisms have not yet been

fully explained. Based on studies conducted to date it is known that electromagnetic fields generate oxidative stress, disrupting the function of sperm mitochondria and causing the activation of cell membrane NADH oxidase [43].

Balance in the free radical system, which is affected by the electromagnetic fields, as well as the activity of ion channels, are together responsible for the process of sperm hyperactivation. Human spermatozoa, in order to become ready to fertilize an oocyte, undergo many metabolic changes affecting the cell membrane, which leads to capacitation occurring in the woman's reproductive tract. As a result of these transformations, sperm can interact with the structures of the corona radiata, and subsequently with the zona pellucida of the oocyte. This process is necessary for fertilization. Possible premature capacitation taking place in the seminal plasma causes energy depletion of the sperm and reduces the chance for fertilization. Balance in the free radical system is the factor responsible for the proper moment of capacitation. Ion channels are responsible for the proper course of this process, including CatSper, a pH-regulated, calcium-selective ion channel, KSper (Slo3), and the voltage-gated channel Hv1. To date, it has been considered that many other channels regulate these processes; however, this was not supported by research, and only the implementation of the patch–clamp technique shed new light on the possibilities of opportunities to learn about this process. It is known that mutations and deletions of the genes responsible for the function of these channels are responsible for the existence of certain types of male infertility. The voltage-gated channel Hv1, which is an electrical voltage sensor responsible for processes related to fertilization, seems to be the most sensitive to the effect of EMF. The results of studies concerning the effect of EMF on the immune system cells confirmed that the frequency close to 15 Hz has the greatest impact on the transport of Ca^{2+} ions into the cell. Therefore, it may be expected that similar relationships will also concern sperm [43,44].

It has long been known that welding arcs are a very harmful source of EMF for males. Their radiation consists of intensive thermal radiation of high-temperature welding gases, welded or cut elements, electrode, and flux material, on which the lines and bands of radiation characteristic of these materials overlap. The gas burner flame temperature usually does not exceed 2000 K. Thus, this radiation consists mainly of infrared and light, and only hydrogen and acetylene burners are characterized by a higher combustion temperature and emit near-ultraviolet light. Scientific reports concerning the effect of welding on male fertility confirm that in males performing this profession, deterioration of the main sperm parameters occurs, and their partners are at an increased risk of miscarriage. The type of harmful effect on the male reproductive system depends on the type of materials to be welded and applies primarily to mild steel [43,44].

Exposure to strong EMF also concerns employees operating radar stations. Currently, various types of radars operate on frequencies from 3 MHz (HF band) to 110 GHz (W band). In the lower part of this range operate radars which measure the height of the ionosphere, and those using the phenomenon of reflection of electromagnetic waves from the ionosphere for detecting objects hidden beyond the horizon, often over a distance of thousands of kilometers. Most radars operate at frequencies ranging from several hundred megahertz to 100 GHz, including speed cameras known to drivers, which are used to take photos of cars and are based on the 34.3 ± 0.1 GHz band. The reports concerning their effect on male fertility are contradictory. Some researchers describe an increase in the percentage of abnormal sperm and reduction in sperm motility, with a simultaneous absence of effect on its density in

employees of radar stations. In turn, other researchers describe a decreased sperm density in people exposed to radar waves. There are also reports that do not indicate any differences in the quality of sperm between males exposed to a radar, and those not exposed to its effect.

Due to the development of telecommunication technologies, at present, almost every person is exposed to EMF related to GSM. Our own study conducted at the beginning of the 'era of mobile phones' demonstrated an increase in the percentage of abnormal sperm and a decrease in the percentage of sperm in progressive movement, together with an increase in exposure to GSM. Studies concerning the effect of mobile phone technology on sperm have been continued by many researchers. They demonstrated that the waves emitted by mobile phones cause an increased production of the reactive oxygen species (ROS), leading to oxidative stress in sperm, and intensification of sperm DNA fragmentation. Similar relationships were confirmed on animal models. In a study in which mice were exposed to radiation of 900 mW/kg for 12 h daily for 7 days an adverse effect of EMR was observed on mitochondrial genome integrity. It was also proven that long-lasting telephone calls lead to an increase in temperature in the brain, which may affect the activity of the hypothalamic–pituitary–gonadal axis [43,44].

A significant exposure to electromagnetic fields also concerns males working on high-voltage electrical installations. It was found that among employees exposed for many years to contact with voltage of 400 kV, fewer children were born compared to the control group. Among children exposed to high voltage, the male gender prevailed. It was also observed that among offspring of males exposed to high voltage, congenital defects occurred more frequently.

There are also other scientific reports that confirm a beneficial effect of electromagnetic fields with lower frequency ranges on both sperm parameters and the function of tissues responsible for human reproduction. Some scientific studies confirmed that exposure of sperm to magnetic fields within the range from 10–50 Hz contributes to the improvement of the motility of human sperm, whereas others presented opposite results.

In animal models, while investigating the exposure of fish reproductive cells to magnetic fields, an improvement was observed in sperm motility parameters, and an increase in the percentage of fertilizations. Electromagnetic fields have found application in the treatment of prostate disorders in animals. A group of researchers confirmed that in dogs, the exposure of the prostate to frequencies of 4–12 Hz has a healing effect on prostate enlargement, not causing changes in sperm parameters of the animals. The results of studies confirming a beneficial effect of electromagnetic fields provide hope for their use in the treatment of human infertility; however, a long-term impact of this type of energy will require meticulous research, especially regarding possible epigenetic effects diagnosed only in future generations. Own studies of sperm exposed in vitro to weak EMF, close to that used at security checkpoints, showed an effect on DNA methylation [43,44].
At present, we are unable to specify what amount of energy related to EMF is harmful, neutral, or beneficial for human reproductive capacity. It may be expected that there are amounts of energy with neutral or beneficial effect, and only exceeding them may cause harmful effects.

Open access paper: https://www.mdpi.com/2077-0383/13/7/1986

--

Influence of power control in the mobile network on the radiation level

Mitić D, Lebl A, Markov Ž. Influence of power control in the mobile network on the radiation level. Journal of Electrical Engineering, 75(2):161-165. 2024. doi: 10.2478/jee-2024-0019.

Abstract

The paper evaluates how the control of transmitted power affects the intensity of radiation in a mobile network cell. Cell models without power control, with standard power control and a model with power control and channel reallocation are considered. The relative reduction of radiation is evaluated and several examples of calculations are presented. Remarks are given on the dependence of radiation reduction on the number of traffic channels, traffic intensity and signal attenuation coefficient. The assessment procedure and results are based on previously verified traffic process simulation models.

Conclusion

The paper shows that control of emitted power, in addition to saving energy and reducing interference, also contributes to the radiation reduction.

Control of the emission power by matching the power with the user's distance affects the reduction of radiation in proportion to the increase in the number of frequency carriers. The radiation level depends on the distribution of the users' surface density in the cell and on the environmental signal attenuation coefficient.

Additional control of the emission power by channel reallocation can further reduce the radiation level. This reduction is greater when there is a smaller number of frequency carriers (but greater than 1), with less offered traffic and with a higher attenuation coefficient.

Open access paper: https://doi.org/10.2478/jee-2024-0019

--

Frequency selective human-centric sub 6 GHz electromagnetic measurements in shopping mall

Berisha D, Rexhëbeqaj-Hamiti V, Dobruna J, Maloku H, Limani Fazliu Z, Ibrani M. (2024). Frequency selective human-centric sub 6 GHz electromagnetic measurements in shopping mall. *Cogent Engineering*, *11*(1). doi: 10.1080/23311916.2024.2340311.

Abstract

Limiting human exposure to radio frequency electromagnetic fields in crowded indoor public spaces such as shopping malls, is identified as one of the key metrics in the process of green and digital environment transformation. To assess the exposure levels in shopping malls, we conducted a human centric extensive measurement campaign using three-axial frequency selective measurement equipment. Our study involved various indoor locations within the mall, capturing electrical field levels at different times and days of the week. A total of 529,340 samples were collected during working days, and 430,020 samples were collected during weekends. We measured E-field strength, power density,

and calculated total exposures for frequency bands ranging from 88 MHz to 5850 MHz. The presented comparative analysis revealed that mobile communications technologies operating at 900 MHz and 1800 MHz were the main contributors to personal exposure levels in different mall environments.

Conclusions

This paper presents the exposure levels in shopping malls during different days of the week, obtained through a rigorous 3-month measurement campaign utilizing frequency selective measurement equipment. The collected data underwent careful processing, including screening of each measured E-field exposure level to ensure accurate sampling and minimize errors.

The average total E-field exposure value during the weekday was 0.39 V/m, while during the weekend 0.33 V/m. The highest contributors to the total exposure levels in various shopping mall environments are mobile communication technologies operating at 900 MHz and 1800 MHz (2G–3G–4G). The Downlink 2G–3G operating at 900 MHz, has an average electric field exposure value of 0.2 V/m during the weekdays and during the weekend, while the Downlink 2G–4G operating at 1800 MHz has an average E-field value of 0.1 V/m during the weekdays and 0.17 V/m during the weekend.

On weekdays, the main factor contributing to total exposure levels was 2G–3G mobile communications operating at 900 MHz (Downlink), accounting for 52.4% of the exposure, followed by mobile communications at 1800 MHz (Uplink and Downlink. Furthermore, the highest E-field values during weekdays were observed in the Downlink 2G–3G band and Wi-Fi 2 G technology, exceeding the upper limit of 6 V/m.

During the weekend, the primary contributor to total exposure levels remained the 2G–3G communications at 900 MHz (Downlink), accounting for 37% of the exposure. However, there was an increase in e-field values for 2G–4G communications at 1800 MHz (Downlink), with its contribution rising to 34% compared to 15% during weekdays. The highest observed value was for Wi-Fi 5G technology, reaching 4.98 V/m, although its contribution to the total exposure levels was only 0.6%.

Based on the cumulative distribution analysis, it can be inferred that during the weekdays, 90% of the measured data have an E-field exposure value of less than 1 V/m. On the other hand, during the weekend, 90% of the measured data have a value smaller than 0.64 V/m. These values are in line with the E-field values recorded in similar studies as referenced in.

In conclusion, based on our extensive measurement campaign, personal exposure levels are higher in shopping malls during the weekdays compared to the weekend. The highest contributors to the total personal exposure at shopping malls are the Mobile communication technologies (Uplink/Downlink) and WI-FI technologies.

Contrary to our initial expectations regarding the indoor environment of the shopping mall, our measurement data reveals that Wi-Fi technologies are not the primary contributors to the total exposure. Instead, mobile communication technologies operating at 900 MHz and 1800 MHz have emerged as the dominant sources due to their high data rates and extensive coverage.

https://www.tandfonline.com/doi/full/10.1080/23311916.2024.2340311

--

Progress in microwave absorbing materials: A critical review

Sharma S, Parne SR, Panda SSS, Gandi S. Progress in microwave absorbing materials: A critical review. Adv Colloid Interface Sci. 2024 Apr 4;327:103143. doi: 10.1016/j.cis.2024.103143.

Abstract

Microwave-absorbing materials play a significant role in various applications that involve the attenuation of electromagnetic radiation. This critical review article provides an overview of the progress made in the development and understanding of microwave-absorbing materials. The interaction between electromagnetic radiation and absorbing materials is explained, with a focus on phenomena such as multiple reflections, scattering, and polarizations. Additionally, types of losses that affect the performance of microwave absorbers are also discussed, including dielectric loss, conduction loss, relaxation loss, magnetic loss, and morphological loss. Each of these losses has different implications for the effectiveness of microwave absorbers. Further, a detailed review is presented on various types of microwave absorbing materials, including carbonaceous materials, conducting polymers, magnetic materials, metals and their composites, 2D materials (such as MXenes and 2D-transition metal dichalcogenides), biomass-derived materials, carbides, sulphides, phosphides, high entropy (HE) materials and metamaterials. The characteristics, advantages, and limitations of each material are examined. Overall, this review article highlights the progress achieved in the field of microwave-absorbing materials. It underlines the importance of optimizing different types of losses to enhance the performance of microwave absorbers. The review also recognizes the potential of emerging materials, such as 2D materials and high entropy materials, in further advancing microwave-absorbing properties.

https://pubmed.ncbi.nlm.nih.gov/38598925/

--

Rosmarinic Acid Protects the Testes of Rats against Cell Phone and Ultra-high Frequency Waves Induced Toxicity

Fatahi Asl J, Goudarzi M, Mansouri E, Shoghi H. Rosmarinic Acid Protects the Testes of Rats against Cell Phone and Ultra-high Frequency Waves Induced Toxicity. Iran J Med Sci. 2024 Apr 1;49(4):237–46. doi: 10.30476/ijms.2023.97695.2952.

Abstract

Background: Cell phone and Ultra-High Frequency (UHF) waves produce oxidative stress and cause testicular toxicity. This investigation was directed to evaluate the effectiveness of Rosmarinic Acid (RA) against oxidative stress caused by UHF radiation in rats.

Methods: Forty-two male Wistar rats were divided into six groups. The control received 5 mL normal saline (0.9% NaCl) by gavage, the cell phone group received 915 MHz, the UHF waves group just received 2450 MHz, the RA/cell phone group received RA plus 915 MHz, RA/UHF waves group received RA plus 2450 MHz, and RA just received RA (20 mg/kg). After 30 days of consecutive radiation, the biochemical and histopathological parameters of their testes were measured. Statistical comparison was made using one-way ANOVA followed by Tukey's post hoc test.

Results: Cell phone and UHF wave radiation significantly diminished the activity of antioxidant enzymes such as superoxide dismutase, catalase, and glutathione peroxidase, and glutathione content (P<0.001). On the opposite, UHF significantly increased oxidative stress indices including malondialdehyde level, nitric oxide level, and protein carbonyl content (P<0.001). UHF also significantly reduced the number of Sertoli cells, spermatogonia, primary spermatocyte, epithelial height, and seminiferous tubular and luminal diameters (P<0.001). RA, as an effective antioxidant, reverses the above-mentioned harms and moderates the adverse effects of UHF on the testes of rats by significantly diminishing the oxidative stress indices and antioxidant enzyme rise and improving the histological parameters (P<0.001).

Conclusion: RA can protect the testes of rats from UHF-induced toxicity by reducing oxidative stress. RA as a food supplement might be useful for protecting humans exposed to UHF environmental contamination.

Open access paper: https://ijms.sums.ac.ir/article_49676.html

--

Panax ginseng Ameliorates Pituitary-Ovarian Dysfunction Induced by RF EMR from Cell Phones via Upregulation of the CREM Signaling Pathway

Oyewopo OA, Badejogbin OC, Ajadi IO, Enye LA, Ajadi MB, Ebuwa IV, Owolabi OV, Areloegbe SE, Olaniyi KS. Panax ginseng Ameliorates Pituitary-Ovarian Dysfunction Induced by Radiofrequency Electromagnetic Radiation from Cell Phones via Upregulation of the CREM Signaling Pathway. Curr Drug Discov Technol. 2024 Apr 30. doi: 10.2174/0115701638279386240425050818.

Abstract

Background: Panax ginseng (PG) is a plant that contains ginsenosides, which are considered adaptogens that confer cellular protection. However, the impact of PG on pituitary-ovarian dysfunction and subsequent infertility is unknown. This study investigated the hypothesis that PG would attenuate pituitary-ovarian dysfunction associated with mobile phone's Radiofrequency Electromagnetic Radiation (RF-EMR) in experimental rat models and the possible involvement of a cAMP Response Element Modulator (CREM)-dependent pathway.

Methods: Twenty adult female Wistar rats were divided randomly into four groups, each consisting of five rats. The control group was administered a vehicle (distilled water) orally, while the P. ginseng group received 200 mg/kg of P. ginseng extract orally. The RF-EMR group was exposed to 900MHz radiation, and the RF-EMR + PG group was exposed to the same radiation while also being treated with 200 mg/kg of P. ginseng orally. These treatments were administered daily for a period of 56 days.

Results: The RF-EMR group exhibited significant reductions in serum levels of LH, FSH, estradiol, and progesterone compared to the control group. Moreover, levels of superoxide dismutase (SOD) and glutathione peroxidase (GPx) were significantly lower in the RF-EMR group compared to the control. Additionally, there was a notable decrease in the expression of the CREM gene, accompanied by disrupted pituitary/ovarian morphology in the RF-EMR group compared to the control. However, the administration of PG mitigated these changes.

Conclusion: The findings of this study indicate that P. ginseng extract shields against pituitary-ovarian impairment linked to RF-EMR exposure from cell phones by boosting antioxidant capacity and promoting the CREM-dependent pathway.

https://pubmed.ncbi.nlm.nih.gov/38693729/

--

Electric and magnetic field pollution in near substations and investigation of anxiety and depressive effects on adult individuals living in this area

Sert C, Başak N, Koruk İ. Electric and magnetic field pollution in near substations and investigation of anxiety and depressive effects on adult individuals living in this area. Electromagn Biol Med. 2024 May 3:1-11. doi: 10.1080/15368378.2024.2348574.

Abstract

Exposure to electromagnetic fields causes a variety of health problems in living systems. We investigated EMF pollution in Şanlıurfa city center and also investigated anxiety-depression symptoms in individuals (18-40 years old) exposed to this pollution. For this purpose, electric field and magnetic field measurements were taken at Electricity Distribution Center and 44 substations (for each transformer), at 0 points, 1 meter away, 2 meters away and the house/office closest to the transformer. The experimental group was individuals living in electricity distribution center residences and individuals living near transformers (n = 55). The control group was selected from individuals who lived outside the city center of Şanlıurfa, did not have transformers or high transmission lines near their homes, and did not have any chronic diseases that could cause stress (n = 50). Anxiety and depression symptoms of the groups were measured using the Beck Anxiety Inventory Scale (BAI) and Beck Depression Inventory Scale (BDI). The relationship between EMF pollution and anxiety-depression was evaluated statistically. Maximum MF and EF values were recorded as 0.22 mT and 65.9 kV/m, respectively. All measured MF values were below standards, but EF values were above standards at some points. In conclusion, there is no statistically convincing evidence of a relationship between EMF exposure and anxiety-depression (p > 0.05). This result shows that there may be more meaningful results in places with higher EMF levels. We interpreted the fact that exposure to electromagnetic fields does not cause anxiety and depression in individuals, as the measured values are below the limit values.

Plain language summary

Electromagnetic field pollution causes various health problems in living systems. Research on this subject continues. We investigated electromagnetic field pollution in Şanlıurfa city center and investigated the psychological effects on individuals exposed to this pollution. For this purpose, electric and magnetic field measurements were made at 0 points, 1 meter, 2 meters away and at the closest places to the transformer in 44 substations. The experimental group was individuals living near a transformer, the control group was individuals living outside the city center and without a transformer/high transmission line near their home. Anxiety and depression symptoms of the groups were measured using the Beck Anxiety and Depression test. The relationship between EMF pollution and anxiety-depression was evaluated statistically. As a result, no statistical change was observed regarding a relationship between electromagnetic pollution and anxiety-depression.

https://pubmed.ncbi.nlm.nih.gov/38699873/

--

The effect of a rotating magnetic field on the antioxidant system in healthy volunteers - preliminary study

Cecerska-Heryć E, Gliźniewicz M, Grygorcewicz B, Serwin N, Stodolak P, Słodzińska W, Birger R, Goszka M, Polikowska A, Budkowska M, Rakoczy R, Dołęgowska B. The effect of a rotating magnetic field on the antioxidant system in healthy volunteers - preliminary study. Sci Rep. 2024 Apr 15;14(1):8677. doi: 10.1038/s41598-024-59391-y.

Abstract

Oxidative stress is characterized by an excessive concentration of reactive oxygen species (ROS) resulting from a disturbance in the balance between ROS production and their removal by antioxidant systems (SOD, CAT, GPx). Prolonged and intense oxidative stress can cause various forms of damage to cells, which markers are total antioxidant capacity (TAC), reactive oxygen species modulator (ROMO1), and malondialdehyde (MDA). It has been demonstrated that magnetic fields can positively affect human health, for example, by reducing oxidative stress. Determination of the effect of a rotating magnetic field (RMF) on the activity/concentration of selected oxidative stress markers. A group of 30 healthy volunteers (15 women and 15 men) (mean age 24.8 ± 5.1) in the study classified into the following groups: internal control group (CG);1 h 25 Hz (samples placed in the field for one hour at 25 Hz); 3 h 25 Hz (samples placed in the field for 3 h at 25 Hz), the 1 h 50 Hz group (placed in RMF for an hour at 50 Hz), and a group of 3 h 50 Hz (samples placed in the field for 3 h at 50 Hz). Serum samples were collected in K_2EDTA tubes.. The magnetic induction value obtained for RMF is 37.06 mT and 42.64 mT. Activity/concentration of selected oxidative stress markers was analyzed by ELISA. The influence of an RMF on the activity/concentration of SOD, MDA, TAC, and ROMO1 was demonstrated ($p < 0.001$; $p = 0.0013$; $p < 0.001$; $p = 0.003$). The RFM can reduce oxidative stress, as evidenced by higher SOD and CAT activities in the CG than in samples placed in the RFM. Prolonged exposure to the RFM at 50 Hz increased the TAC level, indicating an intensification of oxidative stress in these samples. The optimal conditions for staying in the RFM (reducing oxidative stress) are 1 h 50 Hz for SOD and MDA; 3 h 25 Hz for CAT and TAC. In the case of ROMO1, it is stated that 1 h 25 Hz are the optimal conditions for no increased production of ROS.

Open access paper: https://www.nature.com/articles/s41598-024-59391-y

--

Pulsating Extremely Low-Frequency Electromagnetic Fields Influence Differentiation of Mouse Neural Stem Cells towards Astrocyte-like Phenotypes: In Vitro Pilot Study

Isaković J, Slatković F, Jagečić D, Petrović DJ, Mitrečić D. Pulsating Extremely Low-Frequency Electromagnetic Fields Influence Differentiation of Mouse Neural Stem Cells towards Astrocyte-like Phenotypes: In Vitro Pilot Study. Int J Mol Sci. 2024 Apr 4;25(7):4038. doi: 10.3390/ijms25074038.

Abstract

Even though electromagnetic fields have been reported to assist endogenous neurogenesis, little is known about the exact mechanisms of their action. In this pilot study, we investigated the effects of pulsating extremely low-frequency electromagnetic fields on neural stem cell differentiation towards specific phenotypes, such as neurons and astrocytes. Neural stem cells isolated from the telencephalic wall of B6(Cg)-Tyrc-2J/J mouse embryos (E14.5) were randomly divided into three experimental groups and three controls. Electromagnetic field application setup included a solenoid placed within an incubator. Each of the experimental groups was exposed to 50Hz ELF-EMFs of varied strengths for 1 h. The expression of each marker (NES, GFAP, β-3 tubulin) was then assessed by immunocytochemistry. The application of high-strength ELF-EMF significantly increased and low-strength ELF-EMF decreased the expression of GFAP. A similar pattern was observed for β-3 tubulin, with high-strength ELF-EMFs significantly increasing the immunoreactivity of β-3 tubulin and medium- and low-strength ELF-EMFs decreasing it. Changes in NES expression were observed for medium-strength ELF-EMFs, with a demonstrated significant upregulation. This suggests that, even though ELF-EMFs appear to inhibit or promote the differentiation of neural stem cells into neurons or astrocytes, this effect highly depends on the strength and frequency of the fields as well as the duration of their application. While numerous studies have demonstrated the capacity of EMFs to guide the differentiation of NSCs into neuron-like cells or β-3 tubulin+ neurons, this is the first study to suggest that ELF-EMFs may also steer NSC differentiation towards astrocyte-like phenotypes.

Open access paper: https://www.mdpi.com/1422-0067/25/7/4038

--

Invited Perspective: Why Systematic Reviews, Scoping Reviews, and Evidence-to-Decision Frameworks Are Critical for Transparent, Consistent, Equitable, and Science-Based Decision-Making in Environmental Health

My note: Systematic reviews require a multitude of macro- and micro-decisions. When conducted by biased investigative teams, systematic reviews generate erroneous conclusions and policy implications. This applies to EMF as well as chemical exposures.

Chartres N, Joglekar R. Invited Perspective: Why Systematic Reviews, Scoping Reviews, and Evidence-to-Decision Frameworks Are Critical for Transparent, Consistent, Equitable, and Science-Based Decision-Making in Environmental Health. Environ Health Perspect. 2024 Mar;132(3):31304. doi: 10.1289/EHP14346.

No abstract

Excerpt

"... systematic review is simultaneously being corrupted as agencies, including the Texas Commission on Environmental Quality (TCEQ) and the US Environmental Protection Agency (EPA), as part of implementing the amended the Toxic Substances Control Act, are appropriating the term in regulatory decision-making without meeting many of the standards of an empirically based systematic review. 4,9–12 This could result in rulemakings that underestimate the true risk of harm posed by toxic chemical exposures, which could have devastating implications for vulnerable populations, including low-wealth communities of color. 11"

Open access paper: https://ehp.niehs.nih.gov/doi/10.1289/EHP14346

--

Mobile phone use and brain tumour risk – COSMOS, a prospective cohort study

(My note: Numerous methodological problems undermine the validity of the results and conclusions of this study.)

Feychting M, Schüz J, Toledano MB, Vermeulen R, Auvinen A, Poulsen AH, Deltour I, Smith RB, Heller J, Kromhout H, Huss A, Johansen C, Tettamanti G, Elliott P. Mobile phone use and brain tumour risk – COSMOS, a prospective cohort study. Environment International. 2024, doi: 10.1016/j.envint.2024.108552.

Abstract

Background: Each new generation of mobile phone technology has triggered discussions about potential carcinogenicity from exposure to radiofrequency electromagnetic fields (RF-EMF). Available evidence has been insufficient to conclude about long-term and heavy mobile phone use, limited by differential recall and selection bias, or crude exposure assessment. The Cohort Study on Mobile Phones and Health (COSMOS) was specifically designed to overcome these shortcomings.

Methods: We recruited participants in Denmark, Finland, the Netherlands, Sweden, and the UK 2007-2012. The baseline questionnaire assessed lifetime history of mobile phone use. Participants were followed through population-based cancer registers to identify glioma, meningioma, and acoustic neuroma cases during follow-up. Non-differential exposure misclassification was reduced by adjusting estimates of mobile phone call-time through regression calibration methods based on self-reported data and objective operator-recorded information at baseline. Hazard ratios (HR) and 95% confidence intervals (CI) for glioma, meningioma, and acoustic neuroma in relation to lifetime history of mobile phone use were estimated with Cox regression models with attained age as the underlying time-scale, adjusted for country, sex, educational level, and marital status.

Results: 264,574 participants accrued 1,836,479 person-years. During a median follow-up of 7.12 years, 149 glioma, 89 meningioma, and 29 incident cases of acoustic neuroma were diagnosed. The adjusted HR per 100 regression-calibrated cumulative hours of mobile phone call-time was 1.00 (95 % CI 0.98-1.02) for glioma, 1.01 (95 % CI 0.96-1.06) for meningioma, and 1.02 (95 % CI 0.99-1.06) for acoustic neuroma. For glioma, the HR for ≥ 1908 regression-calibrated cumulative hours (90th percentile cut-point) was 1.07 (95 % CI 0.62-1.86). Over 15 years of mobile phone use was not associated with an increased tumour risk; for glioma the HR was 0.97 (95 % CI 0.62-1.52).

Conclusions: Our findings suggest that the cumulative amount of mobile phone use is not associated with the risk of developing glioma, meningioma, or acoustic neuroma.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024001387
Supplementary material:
https://www.sciencedirect.com/science/article/pii/S0160412024001387#s0065

--

Dominance of Smartphone Exposure in 5G Mobile Networks

Chiaraviglio L, Lodovisi C, Bartoletti S, Elzanaty A, M. Slim-Alouini M. Dominance of Smartphone Exposure in 5G Mobile Networks. IEEE Transactions on Mobile Computing, 23(3):2284-2302, 2024. doi: 10.1109/TMC.2023.3252662.

Abstract

The deployment of 5G networks is sometimes questioned due to the impact of ElectroMagnetic Field (EMF) generated by Radio Base Station (RBS) on users. The goal of this work is to analyze such issue from a novel perspective, by comparing RBS EMF against exposure generated by 5G smartphones in commercial deployments. The measurement of exposure from 5G is hampered by several implementation aspects, such as dual connectivity between 4G and 5G, spectrum fragmentation, and carrier aggregation. To face such issues, we deploy a novel framework, called 5G-EA, tailored to the assessment of smartphone and RBS exposure through an innovative measurement algorithm, able to remotely control a programmable spectrum analyzer. Results, obtained in both outdoor and indoor locations, reveal that smartphone exposure (upon generation of uplink traffic) dominates over the RBS one. Moreover, Line-of-Sight locations experience a reduction of around one order of magnitude on the overall exposure compared to Non-Line-of-Sight ones. In addition, 5G exposure always represents a small share (up to 38%) compared to the total one radiated by the smartphone.

Excerpt

Our results demonstrate that the smartphone exposure dominates over the RBS one upon generation of UL traffic, especially when the UE is in Non-Line-of-Sight (NLOS) with respect to the RBS. On the contrary, both smartphone exposure and total EMF are reduced up to one order of magnitude when the smartphone UL traffic traverses a radio link in Line-of-Sight (LOS) with respect to the serving RBS. Interestingly, the exploitation of dual connectivity feature between 4G and 5G reveals that only a small smartphone exposure share (at most equal to 38%) is due to 5G, while the largest exposure levels are derived from the carrier aggregation over 4G bands. Moreover, both total and smartphone exposure-per-bit metrics are inversely proportional to the maximum amount of UL traffic generated by the smartphone in the measurement location, thus suggesting that innovative exposure estimators, based on the reporting of maximum UL traffic from the smartphone, can be designed.

Open access paper: https://doi.ieeecomputersociety.org/10.1109/TMC.2023.3252662

--

Human exposure to EMF from 5G base stations: analysis, evaluation and comparison of different assessment methods

Expósito I, Hakizimali C, García Sánchez M, Cuiñas I, Verhaevert J. Human exposure to EMF from 5G base stations: analysis, evaluation and comparison of different assessment methods. Measurement. 2024. doi: 10.1016/j.measurement.2024.114434.

Highlights

- This paper analyzes the feasibility of assessing the 5G base stations compliance using broadband field probes and compares their performance with alternative methodologies and equipment.
- Performance of three different methodologies and equipment (broadband probes, spectrum analyzers, and drive test scanners), in the context of human exposure to electromagnetic fields (EMF) from 5G base stations, is compared.
- Quantification of the uncertainty that the fluctuation in 5G signal levels induces in the assessment of electromagnetic fields exposure is provided.
- The use of broadband field probes for 5G exposure assessment is still possible under certain considerations and correcting the results considering the base station load and beamforming effects.

Abstract

5G networks deployment poses new challenges when evaluating human exposure to electromagnetic fields. Fast variation of the user load and beamforming techniques may cause large fluctuations of 5G base stations field level. They may be underestimated, resulting in compliance of base stations not fitting the requirements. Apparently, broadband field meters would not be adequate for measuring such environments. However, we analyze the feasibility of confidently using broadband field meters and compare their performance with alternative equipment. Measurements based on the synchronization signals power level, using spectrum analyzers or drive test scanners, may be valid, if gain differences between the signaling and data radiation patterns are characterized. These methods lead to good results but require more time and knowledge. Nevertheless, using broadband field meters is still possible if the measurement results are corrected considering the base station load. Under specific conditions, explained here, fast assessment of 5G compliance could be provided.

Conclusions

Assessing human exposure to an electromagnetic field in presence of a 5G base station is not an easy task. The implementation of M−MIMO techniques in 5G base stations results in adaptive beamforming. This makes difficult to guarantee that the field levels are at their maximum at the measurement location during the complete measurement period, which would limit the applicability of broadband instruments as having been done for previous generations. In this research, we have compared different methods for 5G exposure assessment, using a broadband field meter with an isotropic probe, a spectrum analyzer and a drive test scanner.

Along the paper, we first give an overview of the 5G signal structure, describing the frequency domain and time domain specifications. Afterwards, possible assessment methods are described. The SSB level is measured using the Keysight FieldFox N9913A SA and the Rohde & Schwarz TSM6 DTS. The values are extrapolated to the worst-case exposure and compared to the measurements done with the Wavecontrol WPF8 broadband field probe. Measurements are repeated increasing the base station load by performing a heavy download from a 5G user terminal located near the testers.

The proposed methods were field tested at the University of Vigo, Spain, with a commercial 5G base station located on its campus. The measurements were performed at 7 locations in LOS conditions around the base station, gathering data with the three different equipment at the same locations and at

the same time. This data collection allows the comparison of the three methodologies under the same radiating conditions.

All results have been analyzed considering the specific measurement uncertainties, which allows a deeper and more precise comparison among them.

From the measurement results, we can extract that the exposure levels are low at this stage of the 5G deployment. When loading the base station, the results showed that using the broadband field meter can overestimate the field level. Thus, it is still a useful method to check if the field levels comply with the regulation in human exposure; very simple and cost-effective compared with others. In-situ measurements of human exposure to EMF have to be practical and easy to carry, involving only the resources and equipment strictly necessary, but without compromising the validity of the results. When the reference levels are surpassed, more accurate methods based in the assessment and extrapolation of the SSB level could be a solution. The drawback is the required post processing, specially correcting the gain difference between SSB and data signals. If not provided by the network operator, this difference can be determined through measurements, as explained along this document. Measuring with an SA in max-hold mode in the bandwidth of the SSB does not work in 5G as it does in LTE, as we cannot be sure if the measured level corresponds to the SSB or to the user data, no matter if we are forcing the load of the station or not.

The analysis of the results demonstrate that broadband instruments can be used for assessing human exposure to EMF in the vicinity of 5G base stations, which radiating elements provide fields with extreme fluctuations in their intensity as a function of the system load and beamforming configuration. This is accurate when measurements are done by forcing an extra load of the station and the pointing of an antenna beam towards the probe. The validation of this fast method as a first attempt to assess the compliance of 5G stations permits the testing of these base stations in an efficient way. Only when broadband instrument results (including their uncertainties) would overpass the reference levels, a more detailed analysis would be necessary, which procedure and tips are also depicted along this paper.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0263224124003191

--

5G NR launching in Greece: Preliminary in situ and monitoring network measurements of electromagnetic fields exposure levels at rooftops

Christopoulou MI, Kyritsi T, Yalofas A, Koutounidis D, Karabetsos E. 5G NR launching in Greece: Preliminary in situ and monitoring network measurements of electromagnetic fields exposure levels at rooftops. Bioelectromagnetics. 2024 Mar 5. doi: 10.1002/bem.22502.

Highlights

- In situ measurements of electromagnetic field (EMF) exposure levels at rooftops, close to 117 base stations operating at 5G FR1 in Greece in order to evaluate the contribution of 5G to the total exposure.
- Statistical analysis of the follow-up for inspections of the same base stations, before and after the 5G FR1 launching in Greece.
- Case study of a 5G FR1 base station during its pilot and regular operation based on in situ and broadband & frequency selective 24/7 monitoring sensors measurements data.

Abstract

In Greece, 5G New Radio (NR) has started launching in the end of 2020, at the 3400-3800 MHz (FR1) frequency band. Focusing on 117 Base Stations (BSs) which were already equipped with 5G NR antennas, in situ broadband and frequency selective measurements have been conducted at minimum three points of interest, at adjacent rooftops (when accessible). The points have been selected according to the sweeping method and the electric field strength (E) value has been stored on the selected worst-case scenario point. Spectrum analysis was conducted in the FR1, for the allocated spectrum that corresponds to each mobile communication provider, in order to get preliminary results concerning the contribution of the 5G NR emissions in the general public exposure levels. The vast majority of the in situ measurements has been conducted in urban environments from the beginning of 2021 until the mid of 2022, since in Greece 5G NR services launching started from the big cities. Additionally, a 5G NR BS, installed in a suburban environment (in the city of Kalamata) is thoroughly investigated during its pilot and regular operation, based on broadband and frequency selective measurements data derived by the National Observatory of Electromagnetic Fields (NOEF) monitoring sensor network. In situ measurement data within the 5G NR frequency range are verified via the NOEF's output. The 5G NR contribution to the total E-field levels is assessed in time, from pilot to regular operation of the BS. In all cases, compliance with the reference levels for general public exposure is affirmed.

https://pubmed.ncbi.nlm.nih.gov/38444067/

--

The determinants of legislation for radiofrequency electromagnetic fields (RF-EMFs) with the onset of 5G: An empirical analysis

Recuero Virto L, Czerwiński M, Froidevaux J. The determinants of legislation for radiofrequency electromagnetic fields (RF-EMFs) with the onset of 5G: An empirical analysis with a worldwide cross-sectional dataset. Risk Anal. 2024 Mar 31. doi: 10.1111/risa.14298.

Abstract

The unprecedented exposure of radiofrequency electromagnetic field (RF-EMF) to humans from mobile communications raises serious public concern about the possibility of unexpected adverse health effects and has stimulated authorities to adopt precautionary exposure limits. These limits are distinctly different across countries, and the causes of these differences are unclear from the literature. This article is the first empirical analysis on the determinants of RF-EMF exposure legislation, using a novel cross-sectional database of 164 countries worldwide. The analysis shows that decentralization and mobile competition in countries with low mobile network deployment tend to promote more stringent RF-EMF exposure limits across the dataset with 164 countries. In more decentralized countries, the regions had a greater influence on national legislation and could accommodate local demands with the advent of mobile technology in the 2000s. In contrast, decentralization and mobile competition in countries with high levels of mobile network deployment tend to relax RF-EMF exposure limits in the sample of 61 countries with fifth-generation (5G) technology. Indeed, restrictive RF-EMF exposure limits are constraining 5G deployment in a context of the widespread adoption of mobile-broadband technologies. These results should be useful for policymakers and mobile operators alike to anticipate the outcome of legislation in countries that have yet to introduce 5G technology. The results should also

be useful when reviewing policies and strategies for the implementation of the upcoming 6G technology in frequency bands that will be increasingly higher (above 6 GHz up to THz for very local usage), and hence where the health effects on humans are less well studied.

https://pubmed.ncbi.nlm.nih.gov/38556257/

--

Editorial: Neurological Illness and National Security: Lessons to Be Learned

Relman DA. Neurological Illness and National Security: Lessons to Be Learned. JAMA. Published online March 18, 2024. doi:10.1001/jama.2023.26818

No abstract

Excerpt

In 2016, a set of troubling neurological symptoms was reported through confidential channels by US government personnel based at the US Embassy in Havana, Cuba. As the number of cases in Havana escalated and then similar cases occurred over the next 5 years in other locations around the globe, efforts to understand this syndrome, now known as anomalous health incidents (AHIs), were hampered by their unusual features, incomplete information, nonstandardized clinical testing, delayed reporting, and the sensitive nature of the circumstances, individuals, and their work. A subset of individuals described the abrupt onset, sometimes in the middle of the night, of a loud, grinding, clicking, buzzing, or high-pitched piercing sound inside the head, occasionally likened to a slide whistle, and a sensation of pressure, sometimes in one ear, on one side of the head, or in the face or chest. Most strikingly, these phenomena often displayed strong location dependence, in that they quickly dissipated when the individuals vacated their initial location, and then returned when the location was revisited. In some cases, this location dependence was reported to occur repeatedly by the same individual or by multiple individuals as they moved away from and then returned within minutes to a specific location, such as part of a room. These abrupt-onset sensory phenomena were followed by a mix of vertigo, dizziness, imbalance, blurry vision, tinnitus, headache, nausea, and cognitive dysfunction, sometimes leading to chronic disability.

In this issue of JAMA, Chan et al1 and Pierpaoli et al2 at the National Institutes of Health (NIH) report on an extensive clinical assessment of 86 participants with AHIs and 30 control participants, and on magnetic resonance imaging (MRI) findings of 81 of these participants with AHIs and 48 control participants, respectively. Overall, the authors found few significant differences between participants with AHIs and control participants, and no consistent evidence of brain injury. These findings differ from previous clinical and imaging studies of smaller numbers of cases from Havana and China that found evidence of vestibular, oculomotor, and pupillary abnormalities3,4 and a variety of MRI findings.5

With few differences between cases and controls in the 2 current studies, one might suspect that nothing or nothing serious happened with these cases. This would be ill-advised. Two detailed investigations of AHIs (in which I played a role) found the cases with abrupt-onset, location-dependent sensory phenomena to be unlike any disorder reported in the neurological or general medical literature, and potentially caused by an external mechanism.6-9 The first of these investigations was undertaken

by the US National Academy of Sciences in 2019-2020 and the second by a panel of experts on behalf of the US Intelligence Community in 2021-2022. The latter is described in more detail below. Both of these investigations involved experienced clinicians, detailed interviews with many affected individuals, careful reviews of the literature, and informal reviews of some medical records, but not physical examinations or directed testing. After considering a wide variety of possible mechanisms, both studies concluded that some of the cases with abrupt-onset, location-dependent sensory phenomena could be plausibly explained by exposure to directed, pulsed radiofrequency energy, despite important uncertainties. Others have also pointed to pulsed radiofrequency energy as a plausible mechanism.10,11 The US Intelligence Community has discounted this possibility and concluded that reported symptoms were probably the result of "preexisting conditions, conventional illnesses, and environmental factors," influenced by their assessment that no foreign adversary played a role in these cases.12 While many cases may be explained in this fashion, the evidence that might favor known conditions, illnesses, and factors in some of the cases with abrupt-onset, location-dependent auditory-vestibular phenomena is weak at best....

Open access paper: https://jamanetwork.com/journals/jama/fullarticle/2816534

--

Clinical, biomarker, and research tests among US government personnel and their family members involved in anomalous health incidents

Chan L, Hallett M, Zalewski CK, et al; NIH AHI Intramural Research Program Team. Clinical, biomarker, and research tests among US government personnel and their family members involved in anomalous health incidents. JAMA. Published March 18, 2024. doi:10.1001/jama.2024.2413

Key Points

Questions Do US government officials and their family members involved in anomalous health incidents (AHIs) differ from control participants with respect to clinical, biomarker, and research assessments?

Findings In this exploratory study that included 86 participants reporting AHIs and 30 vocationally matched control participants, there were no significant differences in most tests of auditory, vestibular, cognitive, visual function, or blood biomarkers between the groups. Participants with AHIs performed significantly worse on self-reported and objective measures of balance, and had significantly increased symptoms of fatigue, posttraumatic stress disorder, and depression compared with the control participants; 24 participants (28%) with AHIs presented with functional neurological disorders.

Meaning In this exploratory study, there were no significant differences between individuals reporting AHIs and matched control participants with respect to most clinical, research, and biomarker measures, except for self-reported and objective measures of imbalance; symptoms of fatigue, posttraumatic stress, and depression; and the development of functional neurological disorders in some.

Abstract

Importance Since 2015, US government and related personnel have reported dizziness, pain, visual problems, and cognitive dysfunction after experiencing intrusive sounds and head pressure. The US government has labeled these anomalous health incidents (AHIs).

Objective To assess whether participants with AHIs differ significantly from US government control participants with respect to clinical, research, and biomarker assessments.

Design, Setting, and Participants Exploratory study conducted between June 2018 and July 2022 at the National Institutes of Health Clinical Center, involving 86 US government staff and family members with AHIs from Cuba, Austria, China, and other locations as well as 30 US government control participants.

Exposures AHIs.

Main Outcomes and Measures Participants were assessed with extensive clinical, auditory, vestibular, balance, visual, neuropsychological, and blood biomarkers (glial fibrillary acidic protein and neurofilament light) testing. The patients were analyzed based on the risk characteristics of the AHI identifying concerning cases as well as geographic location.

Results Eighty-six participants with AHIs (42 women and 44 men; mean [SD] age, 42.1 [9.1] years) and 30 vocationally matched government control participants (11 women and 19 men; mean [SD] age, 43.8 [10.1] years) were included in the analyses. Participants with AHIs were evaluated a median of 76 days (IQR, 30-537) from the most recent incident. In general, there were no significant differences between participants with AHIs and control participants in most tests of auditory, vestibular, cognitive, or visual function as well as levels of the blood biomarkers. Participants with AHIs had significantly increased fatigue, depression, posttraumatic stress, imbalance, and neurobehavioral symptoms compared with the control participants. There were no differences in these findings based on the risk characteristics of the incident or geographic location of the AHIs. Twenty-four patients (28%) with AHI presented with functional neurological disorders.

Conclusions and Relevance In this exploratory study, there were no significant differences between individuals reporting AHIs and matched control participants with respect to most clinical, research, and biomarker measures, except for objective and self-reported measures of imbalance and symptoms of fatigue, posttraumatic stress, and depression. This study did not replicate the findings of previous studies, although differences in the populations included and the timing of assessments limit direct comparisons.

Open access paper: https://jamanetwork.com/journals/jama/fullarticle/2816533

--

Neuroimaging findings in US government personnel and their family members involved in anomalous health incidents

Pierpaoli C, Nayak A, Hafiz R, et al; NIH AHI Intramural Research Program Team. Neuroimaging findings in US government personnel and their family members involved in anomalous health incidents. JAMA. Published March 18, 2024. doi:10.1001/jama.2024.2424

Key Points

Question Can a systematic evaluation using quantitative magnetic resonance imaging (MRI) metrics identify potential brain lesions in patients who have experienced anomalous health incidents (AHIs) compared with a well-matched control group?

Findings In this exploratory study that involved brain imaging of 81 participants who experienced AHIs and 48 matched control participants, there were no significant between-group differences in MRI measures of volume, diffusion MRI–derived metrics, or functional connectivity using functional MRI after adjustments for multiple comparisons. The MRI results were highly reproducible and stable at longitudinal follow-ups. No clear relationships between imaging and clinical variables emerged.

Meaning In this exploratory neuroimaging study, there was no significant MRI-detectable evidence of brain injury among the group of participants who experienced AHIs compared with a group of matched control participants. This finding has implications for future research efforts as well as for interventions aimed at improving clinical care for the participants who experienced AHIs.

Abstract

Importance US government personnel stationed internationally have reported anomalous health incidents (AHIs), with some individuals experiencing persistent debilitating symptoms.

Objective To assess the potential presence of magnetic resonance imaging (MRI)–detectable brain lesions in participants with AHIs, with respect to a well-matched control group.

Design, Setting, and Participants This exploratory study was conducted at the National Institutes of Health (NIH) Clinical Center and the NIH MRI Research Facility between June 2018 and November 2022. Eighty-one participants with AHIs and 48 age- and sex-matched control participants, 29 of whom had similar employment as the AHI group, were assessed with clinical, volumetric, and functional MRI. A high-quality diffusion MRI scan and a second volumetric scan were also acquired during a different session. The structural MRI acquisition protocol was optimized to achieve high reproducibility. Forty-nine participants with AHIs had at least 1 additional imaging session approximately 6 to 12 months from the first visit.

Exposure AHIs.

Main Outcomes and Measures Group-level quantitative metrics obtained from multiple modalities: (1) volumetric measurement, voxel-wise and region of interest (ROI)–wise; (2) diffusion MRI–derived metrics, voxel-wise and ROI-wise; and (3) ROI-wise within-network resting-state functional connectivity using functional MRI. Exploratory data analyses used both standard, nonparametric tests and bayesian multilevel modeling.

Results Among the 81 participants with AHIs, the mean (SD) age was 42 (9) years and 49% were female;

among the 48 control participants, the mean (SD) age was 43 (11) years and 42% were female. Imaging scans were performed as early as 14 days after experiencing AHIs with a median delay period of 80 (IQR, 36-544) days. After adjustment for multiple comparisons, no significant differences between participants with AHIs and control participants were found for any MRI modality. At an unadjusted threshold ($P < .05$), compared with control participants, participants with AHIs had lower intranetwork connectivity in the salience networks, a larger corpus callosum, and diffusion MRI differences in the corpus callosum, superior longitudinal fasciculus, cingulum, inferior cerebellar peduncle, and amygdala. The structural MRI measurements were highly reproducible (median coefficient of variation <1% across all global volumetric ROIs and <1.5% for all white matter ROIs for diffusion metrics). Even individuals with large differences from control participants exhibited stable longitudinal results (typically, <±1% across visits), suggesting the absence of evolving lesions. The relationships between the imaging and clinical variables were weak (median Spearman $\rho = 0.10$). The study did not replicate the results of a previously published investigation of AHIs.

Conclusions and Relevance In this exploratory neuroimaging study, there were no significant differences in imaging measures of brain structure or function between individuals reporting AHIs and matched control participants after adjustment for multiple comparisons.

Open access paper: https://jamanetwork.com/journals/jama/fullarticle/2816532

--

Do blue light filter applications improve sleep outcomes? A study of smartphone users' sleep quality in an observational setting

Rabiei M, Masoumi SJ, Haghani M, Nematolahi S, Rabiei R, Mortazavi SMJ (2024). Do blue light filter applications improve sleep outcomes? A study of smartphone users' sleep quality in an observational setting. Electromagnetic Biology and Medicine, DOI: 10.1080/15368378.2024.2327432.

Abstract

Exposure to blue light at bedtime, suppresses melatonin secretion, postponing the sleep onset and interrupting the sleep process. Some smartphone manufacturers have introduced night-mode functions, which have been claimed to aid in improving sleep quality. In this study, we evaluate the impact of blue light filter application on decreasing blue light emissions and improving sleep quality. Participants in this study recorded the pattern of using their mobile phones through a questionnaire. In order to evaluate sleep quality, we used a PSQI questionnaire. Blue light filters were used by 9.7% of respondents, 9.7% occasionally, and 80% never. The mean score of PSQI was more than 5 in 54.10% of the participants and less than 5 in 45.90%. ANOVA test was performed to assess the relationship between using blue light filter applications and sleep quality (p-value = 0.925). The findings of this study indicate a connection between the use of blue light filter apps and habitual sleep efficiency in the 31–40 age group. However, our results align only to some extent with prior research, as we did not observe sustained positive effects on all parameters of sleep quality from the long-term use of blue light filtering apps. Several studies have found that blue light exposure can suppress melatonin secretion, exacerbating sleep problems. Some studies have reported that physical blue light filters, such as lenses, can affect melatonin secretion and improve sleep quality. However, the impact of blue light filtering applications remains unclear and debatable.

Plain Language Summary

Using smartphones before bedtime and being exposed to its blue light can make it harder to fall asleep and disrupt your sleep. Some smartphone makers have introduced a night mode feature claiming it can help improve your sleep. In this study, we wanted to find out if using these blue light filters on smartphones really makes a difference. We asked people how often they used blue light filters on their phones and also had them fill out a questionnaire about their sleep quality. Only about 10% of people said they used blue light filters regularly, another 10% used them occasionally, and the majority, around 80%, never used them. When we looked at the results, more than half of the participants had sleep scores higher than 5, indicating they might have sleep problems. Less than half had sleep scores lower than 5, suggesting better sleep quality. We used some statistical tests to see if using blue light filters had any link to sleep quality, and the results showed that there was only a connection between the use of blue light filter apps and habitual sleep efficiency in the 31–40 age group. Our findings matched what other studies have found before, that using blue light filters on smartphones may not significantly help improve sleep. So, while it might be a good idea to limit smartphone use before bed, using a blue light filter app may not be the magic solution for better sleep.

https://www.tandfonline.com/doi/full/10.1080/15368378.2024.2327432

--

Mobile phone radiation disturbs cytokinesis and causes cell death in buccal cells: Results of controlled human intervention study

Kundi M, Nersesyan A, Schmid G, Hutter HP, Eibensteiner F, Mišík M, Knasmüller S. Mobile phone specific radiation disturbs cytokinesis and causes cell death but not acute chromosomal damage in buccal cells: Results of a controlled human intervention study. Environ Res. 2024 Mar 5:118634. doi: 10.1016/j.envres.2024.118634.

Abstract

Several human studies indicate that mobile phone specific electromagnetic fields may cause cancer in humans but the underlying molecular mechanisms are currently not known. Studies concerning chromosomal damage (which is causally related to cancer induction) are controversial and are based on the use of questionnaires to assess the exposure. We realized the first human intervention trial in which chromosomal damage and acute toxic effects were studied under controlled conditions. The participants were exposed via headsets at one randomly assigned side of the head to low and high doses of a UMTS signal (n = 20, to 0.1 W/kg and n = 21 to 1.6 W/kg Specific Absorption Rate) for 2h on 5 consecutive days. Before and three weeks after the exposure buccal cells were collected from both cheeks and micronuclei (MN, which are formed as a consequence of structural and numerical chromosomal aberrations) and other nuclear anomalies reflecting mitotic disturbance and acute cytotoxic effects were scored. We found no evidence for induction of MN and of nuclear buds which are caused by gene amplifications, but a significant increase of binucleated cells which are formed as a consequence of disturbed cell divisions, and of karyolitic cells, which are indicative for cell death. No such effects were seen in cells from the less exposed side. Our findings indicate that mobile phone specific high frequency electromagnetic fields do not cause acute chromosomal damage in oral mucosa cells under the present experimental conditions. However, we found clear evidence for disturbance of the cell cycle and cytotoxicity. These effects may play a causal role in the induction of adverse long term health effects in humans.

Final paragraph of paper:

As mentioned in the introduction, evidence is accumulating that exposure to HF-EMF is associated with specific brain tumors (Brabant et al., 2023; Carlberg et al., 2017; Coureau et al., 2014; Hardell and Carlberg, 2015; Hardell et al., 2013; IARC, 2013; INTERPHONE Study Group, 2010). The results of the present investigation indicate that molecular mechanisms other than chromosomal damage may cause neoplastic transformation of the cells as a consequence of exposure to mobile phone specific HF-EMF. As described in the result section, we found in the present study clear evidence for induction of acute toxicity and disturbance of the cell cycle (cytokinesis) as a consequence of exposure to a high radiation dose (1.6 W/kg). It is possible that these effects cause inflammatory responses and/or release of ROS, which were seen in a number of laboratory studies (e.g. Alipour et al., 2022; Benavides et al., 2023; IARC, 2013; Yakymenko et al., 2016). These processes may possibly lead to formation of neoplastic cells.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0013935124005383?via%3Dihub

--

Evaluation of neonatal outcomes according to the specific absorption rate values of phones used during pregnancy

Büyükeren M, Karanfil Yaman F. Evaluation of neonatal outcomes according to the specific absorption rate values of phones used during pregnancy. J Turk Ger Gynecol Assoc. 2024 Mar 6;25(1):7-12. doi: 10.4274/jtgga.galenos.2023.2022-10-1.

Abstract

Objective: The aim was to compare neonatal outcomes according to cell phone specific absorption rate (SAR) levels and daily time spent on cell phones by pregnant women.

Material and methods: Women who gave birth at Konya City Hospital between September 2020 and February 2021 were included in this retrospective study. Gestational ages, birth weight, birth length, head circumference, sex, 5-minute APGAR scores, neonate postpartum resuscitation requirement, delivery type, the model of phone used by the pregnant women, and the average time spent on the phone during a day were recorded. To determine the relation between the SAR values of the phones used and delivering a small for gestational age (SGA) baby, receiver operating characteristic curve analysis was performed.

Results: In total 1495 pregnant women were included. The rate of delivering a SGA fetus was significantly higher in women who used phones with higher SAR values (p=0.001). The cut-off value for the SAR level was 1.23 W/kg with 69.3% sensitivity and 73.0% specificity (area under the curve: 0.685; 95% confidence interval: 0.643-0.726). No correlation was found between time spent on the phone and SGA birth rate. Although both phone SAR values and time spent on the phone were higher in the symmetrical SGA group compared to the asymmetrical SGA group, the difference was not significant (p>0.05). Although the women who had preterm delivery had higher phone SAR values and had spent more time on the phone compared to those who had term deliveries, the difference was again not significant (p>0.05).

Conclusion: As the SAR values of cell phones used during pregnancy increased, there was a trend towards delivering a SGA baby.

Open access paper: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10921072/

--

Do somatic symptom distress and attribution predict symptoms associated with environmental factors?

Köteles F, Nordin S. Do somatic symptom distress and attribution predict symptoms associated with environmental factors? J Psychosom Res. 2024 Mar 1;179:111637. doi: 10.1016/j.jpsychores.2024.111637.

Abstract

Objective: Not much is known on the development of symptoms associated with environmental factors (SAEF), also known as (idiopathic) environmental intolerances. Findings from qualitative studies suggest that appearance of symptoms might be the first step, followed by the acquisition of a specific attribution. The current study investigated cross-sectional and longitudinal (three years) associations between attribution and symptoms with respect to symptoms associated with chemical substances, certain indoor environments (buildings), sounds, and electromagnetic fields (EMFs).

Methods: We used data from the first two waves of the population-based Västerbotten Environmental Health Study (n = 2336). Participants completed the Patient Health Questionnaire Somatic Symptom Scale (PHQ-15), the Environmental Symptom-Attribution Scale, and answered single questions on the four aforementioned SAEFs.

Results: Using binary logistic regression analyses, all four SAEFs showed significant cross-sectional associations with somatic symptom distress and the respective attribution. In the longitudinal analysis, development of SAEF-Sound and SAEF-Chemicals were predicted by both somatic symptom distress and attribution. SAEF-EMFs was predicted only by attribution, whereas neither somatic symptom distress nor attribution forecasted SAEF-Buildings.

Conclusion: Overall, these findings suggest that attribution (i.e., a specific expectation) plays a substantial role in the development and maintenance of many SAEFs.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0022399924000497?via%3Dihub

--

Micro-environmental personal radio-frequency electromagnetic field exposures in Melbourne: A longitudinal trend analysis

Bhatt CR, Henderson S, Sanagou M, Brzozek C, Thielens A, Benke G, Loughran S. Micro-environmental personal radio-frequency electromagnetic field exposures in Melbourne: A longitudinal trend analysis. Environ Res. 2024 Mar 13:118629. doi: 10.1016/j.envres.2024.118629.

Abstract

Background: A knowledge gap exists regarding longitudinal assessment of personal radio-frequency electromagnetic field (RF-EMF) exposures globally. It is unclear how the change in telecommunication technology over the years translates to change in RF-EMF exposure. This study aims to evaluate longitudinal trends of micro-environmental personal RF-EMF exposures in Australia.

Methods: The study utilised baseline (2015-16) and follow-up (2022) data on personal RF-EMF exposure (88 MHz-6 GHz) measured across 18 micro-environments in Melbourne. Simultaneous quantile regression analysis was conducted to compare exposure data distribution percentiles, particularly median (P50), upper extreme value (P99) and overall exposure trends. RF-EMF exposures were compared across six exposure source types: mobile downlink, mobile uplink, broadcast, 5G-New Radio, Others and Total (of the aforementioned sources). Frequency-specific exposures measured at baseline and follow-up were also compared. Total exposure across different groups of micro-environment types were also compared.

Results: For all micro-environmental data, total (median and P99) exposure levels did not significantly change at follow-up. Overall exposure trend of total exposure increased at follow-up. Mobile downlink contributed the highest exposure among all sources showing an increase in median exposure and overall exposure trend. Of seven micro-environment types, five of them showed total exposure levels (median and P99) and overall exposure trend increased at follow-up.

Excerpt

The assessment of change in total personal RF-EMF exposure distribution at follow-up across all micro-environments and those for different micro-environment types showed inconsistent changes in the exposure levels and overall exposure trend. The median and upper extreme total RF-EMF exposure levels across the micro-environments showed no significant change; whilst overall trend of total exposure at follow-up increased during the study period. Mobile downlink, the largest exposure source, median exposure and overall exposure trend increased (26.7%, up to 34.3%, respectively) at follow-up. The observed increase in median and upper extreme exposure levels as well as overall total exposure trend at follow-up remained consistent for the majority of micro-environment types.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0013935124005334?via%3Dihub

--

New-generation electronic appliances and cardiac implantable electronic devices: a systematic literature review of mechanisms and in vivo studies

Kewcharoen J, Shah K, Bhardwaj R, Contractor T, Turagam MK, Mandapati R, Lakkireddy D, Garg J. New-generation electronic appliances and cardiac implantable electronic devices: a systematic literature

review of mechanisms and in vivo studies. J Interv Card Electrophysiol. 2024 Mar 5. doi: 10.1007/s10840-024-01777-z.

Abstract

Introduction: Cardiac implantable electronic device (CIED) functions are susceptible to electromagnetic interference (EMI) from electromagnetic fields (EMF). Data on EMI risks from new-generation electronic appliances (EA) are limited.

Objective: We performed a systematic literature review on the mechanisms of EMI, current evidence, and recently published trials evaluating the effect of EMF on CIEDs from electric vehicles (EV), smartphone, and smartwatch technology and summarize its safety data.

Methods: Electronic databases, including PubMed and EMBASE, were searched for in vivo studies evaluating EMF strength and incidence between CIEDs and commercial EVs, new-generation smartphones, and new-generation smartwatches.

Results: A total of ten studies (three on EVs, five on smartphones, one on smartphones, one on smartphones and smartwatches) were included in our systematic review. There was no report of EMI incidence associated with EVs or smartwatches. Magnet-containing smartphones (iPhone 12) can cause EMI when placed directly over CIEDs - thereby triggering the magnet mode; otherwise, no report of EMI was observed with other positions or smartphone models.

Conclusion: Current evidence suggests CIED recipients are safe from general interaction with EVs/HEVs, smartphones, and smartwatches. Strictly, results may only be applied to commercial brands or models tested in the published studies. There is limited data on EMI risk from EVs wireless charging and smartphones with MagSafe technology.

https://pubmed.ncbi.nlm.nih.gov/38443707/

--

Electric vehicles and health: A scoping review

Pennington AF, Cornwell CR, Sircar KD, Mirabelli MC. Electric vehicles and health: A scoping review. Environ Res. 2024 Mar 16:118697. doi: 10.1016/j.envres.2024.118697.

Abstract

Background: The health impacts of the rapid transition to the use of electric vehicles are largely unexplored. We completed a scoping review to assess the state of the evidence on use of battery electric and hybrid electric vehicles and health.

Methods: We conducted a literature search of MEDLINE, Embase, Global Health, CINAHL, Scopus, and Environmental Science Collection databases for articles published January 1990 to January 2024. We included articles if they presented observed or modeled data on the association between battery electric or hybrid electric cars, trucks, or buses and health-related outcomes. We abstracted data and

summarized results.

Results: Out of 897 reviewed articles, 52 met our inclusion criteria. The majority of included articles examined transitions to the use of electric vehicles (n = 49, 94%), with fewer studies examining hybrid electric vehicles (n = 11, 21%) or plug-in hybrid electric vehicles (n = 8, 15%). The most common outcomes examined were premature death (n = 41, 79%) and monetized health outcomes such as medical expenditures (n = 33, 63%). We identified only one observational study on the impact of electric vehicles on health; all other studies reported modeled data. Almost every study (n = 51, 98%) reported some evidence of a positive health impact of transitioning to electric or hybrid electric vehicles, although magnitudes of association varied. There was a paucity of information on the environmental justice implications of vehicle transitions.

Conclusions: The results of the current literature on electric vehicles and health suggest an overall positive health impact of transitioning to electric vehicles. Additional observational studies would help expand our understanding of the real-world health effects of electric vehicles. Future research focused on the environmental justice implications of vehicle fleet transitions could provide additional information about the extent to which the health benefits occur equitably across populations.

https://pubmed.ncbi.nlm.nih.gov/38499224/

--

Electromagnetic exposure analysis of the subway passenger under the civil communication system radiation

Zhou WY, Zhang XY, Lu M. Electromagnetic exposure analysis of the subway passenger under the civil communication system radiation. PLoS One. 2024 Mar 11;19(3):e0300049. doi: 10.1371/journal.pone.0300049.

Abstract

In order to assess the electromagnetic exposure safety of passengers under the civil communication system of the subway, the radio-frequency (RF) electromagnetic environment of subway carriage is established by using COMSOL Multiphysics software, it includes a 1-1/4 " leaky coaxial cable (LCX1) and a 1-5/8" leaky coaxial cable (LCX2), which are designed to be the exposure sources, and twelve passengers at different position. The electromagnetic environment model has been verified through field measurement. The exposure dose distribution of twelve passengers is compared and analyzed, when LCX1 and LCX2 works respectively. The simulated results show that, to compare with LCX2, the electromagnetic dose absorbed by the passengers is reduced by 9.19% and 22.50% at 2100 MHz and 2600 MHz respectively. The specific absorption rate (SAR) of passengers obtains the maximum value of 1.91×10^{-4} W/Kg and the temperature rise to 0.214 K when the LCX1 works at 3400 MHz. By comparing with the public exposure limitation of the International Commission of Non-Ionizing Radiation Protection (ICNIRP), it demonstrates the electromagnetic exposure safety of the passengers under the civil communication system. More importantly, the proposed LCX1 not only could add the 5G signal cover but also lower the SAR absorbed by the passengers, which indicates that the public electromagnetic exposure dose could be reduced by adjusting the radiation performances of exposure source, which provide a new way for electromagnetic protecting.

Open access paper: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0300049

--

Impact of specific electromagnetic radiation on wakefulness in mice

Deng H, Liu L, Tang X, Lu Y, Wang X, Zhao Y, Shi Y. Impact of specific electromagnetic radiation on wakefulness in mice. Proc Natl Acad Sci U S A. 2024 Apr 9;121(15):e2313903121. doi: 10.1073/pnas.2313903121.

Abstract

Electromagnetic radiation (EMR) in the environment, particularly in the microwave range, may constitute a public health concern. Exposure to 2.4 GHz EMR modulated by 100 Hz square pulses was recently reported to markedly increase wakefulness in mice. Here, we demonstrate that a similar wakefulness increase can be induced by the modulation frequency of 1,000 Hz, but not 10 Hz. In contrast to the carrier frequency of 2.4 GHz, 935 MHz EMR of the same power density has little impact on wakefulness irrespective of modulation frequency. Notably, the replacement of the 100 Hz square-pulsed modulation by sinusoidal-pulsed modulation of 2.4 GHz EMR still allows a marked increase of wakefulness. In contrast, continuous sinusoidal amplitude modulation of 100 Hz with the same time-averaged power output fails to trigger any detectable change of wakefulness. Therefore, alteration of sleep behavior by EMR depends upon not just carrier frequency but also frequency and mode of the modulation. These results implicate biological sensing mechanisms for specific EMR in animals.

Significance

Increased wakefulness in mice was previously found to be a direct result of prolonged exposure to 2.4 GHz electromagnetic radiation (EMR) with 100 Hz square-pulsed modulation at 1/8 duty cycle. Several key issues remain unaddressed. Does the frequency of the square-pulsed modulation matter? Are the sharp edges of the square pulses a major contributor to sleep/wakefulness alteration? Can carrier frequencies other than 2.4 GHz induce sleep/wakefulness alteration? Does the duty cycle matter? In this study, we answer these questions by demonstrating the dependency of sleep/wakefulness alteration on EMR modulation frequency, carrier frequency, and modulation mode.

Excerpts

Modeling and experimental assessment of human exposure to 935 MHz or 2.14 GHz EMR at a dose level of 3.6 W/kg local SAR revealed a maximal skin temperature increase of 0.31 °C and a brain temperature elevation of <0.1 °C (21). In our case, the maximum local SAR value measured with the cSAR3D testing system is 3.6 W/kg and the averaged SAR is 2.81 ± 0.15 W/kg. Hence, the observed EMR impact on sleep in our study is most likely nonthermal....

In conclusion, our study reveals distinct specificity of EMR. Prolonged exposure to the carrier frequency 2.4 GHz EMR with square pulse modulation of different frequencies induces varying changes of wakefulness in mice. In contrast to 2.4 GHz, the carrier frequency 935 MHz has little impact on wakefulness or NREM sleep. Then, 2.4 GHz EMR with sinusoidal pulse modulation, but not continuous sinusoidal modulation of 100 Hz, results in an increase in wakefulness. These findings link specific biological responses to specific parameters of EMR, namely carrier frequency, modulation frequency, and modulation mode. The underlying mechanisms for these observations remain to be unveiled.

https://pubmed.ncbi.nlm.nih.gov/38557178/

--

Effect of 2.45 GHz Microwave Radiation on the Inner Ear: A Histopathological Study on 2.45 GHz Microwave Radiation and Cochlea

Tahir E, Akar Karadayı A, Gülşen Gürgen S, Korunur Engiz B, Turgut A. Effect of 2.45 GHz Microwave Radiation on the Inner Ear: A Histopathological Study on 2.45 GHz Microwave Radiation and Cochlea. J Int Adv Otol. 2024 Jan;20(1):35-43. doi: 10.5152/iao.2024.231142.

Abstract

Background: The present study aims to determine the possible low dose-dependent adverse effects of 2.45 GHz microwave exposure and Wi-Fi frequency on the cochlea.

Methods: Twelve pregnant female rats (n=12) and their male newborns were exposed to Wi-Fi frequencies with varying electric field values of 0.6, 1.9, 5, 10 V/m, and 15 V/m during the 21-day gestation period and 45 days after birth, except for the control group. Auditory brainstem response testing was performed before exposure and sacrification. After removal of the cochlea, histopathological examination was conducted by immunohistochemistry methods using caspase (cysteine-aspartic proteases, cysteine aspartates, or cysteine-dependent aspartate-directed proteases)-3, -9, and terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL). Kruskal-Wallis and Wilcoxon tests and multivariate analysis of variance were used.

Results: Auditory brainstem response thresholds in postexposure tests increased statistically significantly at 5 V/m and above doses. When the number of apoptotic cells was compared in immunohistochemistry examination, significant differences were found at 10 V/m and 15 V/m doses ($F(5,15)=23.203$, $P=.001$; Pillai's trace=1.912, $\eta 2=0.637$). As the magnitude of the electric field increased, all histopathological indicators of apoptosis increased. The most significant effect was noted on caspase-9 staining ($\eta 2$ c9=0.996), followed by caspase-3 ($\eta 2$ c3=0.991), and TUNEL staining ($\eta 2$ t=0.801). Caspase-3, caspase-9, and TUNEL-stained cell densities increased directly by increasing the electric field and power values.

Conclusion: Apoptosis and immune activity in the cochlea depend on the electric field and power value. Even at low doses, the electromagnetic field in Wi-Fi frequency damages the inner ear and causes apoptosis.

Open access paper: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10895889/

--

Ashwagandha Diminishes Hippocampal Apoptosis Induced by Wi-Fi Radiation in Male Quails

Gupta V, Srivastava R. Ashwagandha Diminishes Hippocampal Apoptosis Induced by Microwave Radiation by Acetylcholinesterase Dependent Neuro-Inflammatory Pathway in Male Coturnix coturnix Japonica. Neurochem Res. 2024 Mar 20. doi: 10.1007/s11064-024-04127-7.

Abstract

Microwave radiation (MWR) has been linked to neurodegeneration by inducing oxidative stress in the hippocampus of brain responsible for learning and memory. Ashwagandha (ASW), a medicinal plant is known to prevent neurodegeneration and promote neuronal health. This study investigated the effects of MWR and ASW on oxidative stress and cholinergic imbalance in the hippocampus of adult male Japanese quail. One control group received no treatment, the second group quails were exposed to MWR at 2 h/day for 30 days, third was administered with ASW root extract orally 100 mg/day/kg body weight and the fourth was exposed to MWR and also treated with ASW. The results showed that MWR increased serum corticosterone levels, disrupted cholinergic balance and induced neuro-inflammation. This neuro-inflammation further led to oxidative stress, as evidenced by decreased activity of antioxidant enzymes SOD, CAT and GSH. MWR also caused a significant decline in the nissil substances in the hippocampus region of brain indicating neurodegeneration through oxidative stress mediated hippocampal apoptosis. ASW, on the other hand, was able to effectively enhance the cholinergic balance and subsequently lower inflammation in hippocampus neurons. This suggests that ASW can protect against the neurodegenerative effects of MWR. ASW also reduced excessive ROS production by increasing the activity of ROS-scavenging enzymes. Additionally, ASW prevented neurodegeneration through decreased expression of caspase-3 and caspase-7 in hippocampus, thus promoting neuronal health. In conclusion, this study showed that MWR induces apoptosis and oxidative stress in the brain, while ASW reduces excessive ROS production, prevents neurodegeneration and promotes neuronal health.

Excerpt

Birds were exposed to 2.45 GHz of MW irradiation using Ruckus R310 Wi-fi router with maximum transmission strength of 25dbm at the middle of the cage. Average power density obtained within each partition of cage was found to be 0.1264 mw/cm^2 and overall specific absorption rate (SAR) obtained per bird was determined to be 0.9978 W/Kg.

https://pubmed.ncbi.nlm.nih.gov/38506951/

--

Does Microwave Exposure at Different Doses in the Pre/Postnatal Period Affect Growing Rat Bone Development?

Karadayi A, Sarsmaz H, Çigel A, Engiz B, Ünal N, Ürkmez S, Gürgen S. Does Microwave Exposure at Different Doses in the Pre/Postnatal Period Affect Growing Rat Bone Development? Physiol Res. 2024 Mar 11;73(1):157-172.

Abstract

Effects of pre/postnatal 2.45 GHz continuous wave (CW), Wireless-Fidelity (Wi-Fi) Microwave (MW) irradiation on bone have yet to be well defined. The present study used biochemical and histological methods to investigate effects on bone formation and resorption in the serum and the tibia bone tissues

of growing rats exposed to MW irradiation during the pre/postnatal period. Six groups were created: one control group and five experimental groups subjected to low-level different electromagnetic fields (EMF) of growing male rats born from pregnant rats. During the experiment, the bodies of all five groups were exposed to 2.45 GHz CW-MW for one hour/day. EMF exposure started after fertilization in the experimental group. When the growing male rats were 45 days old in the postnatal period, the control and five experimental groups' growing male and maternal rats were sacrificed, and their tibia tissues were removed. Maternal rats were not included in the study. No differences were observed between the control and five experimental groups in Receptor Activator Nuclear factor-kB (RANK) biochemical results. In contrast, there was a statistically significant increase in soluble Receptor Activator of Nuclear factor-kB Ligand (sRANKL) and Osteoprotegerin (OPG) for 10 V/m and 15 V/m EMF values. Histologically, changes in the same groups supported biochemical results. These results indicate that pre/postnatal exposure to 2.45 GHz EMF at 10 and 15 V/m potentially affects bone development.

Excerpt

In the present study, the effects of 2.45 GHz MW radiation on the bone of healthy rat tibia exposed to different doses of EMF during the prenatal and postnatal period were investigated using biochemical methods such as RANK, RANKL, OPG, and histopathological methods such as Tunel and immunohistochemical straining. Our findings showed that 2.45 GHz low-level MW radiation at 10 V/m (the peak SAR 10g value 14.4 mW/kg) and 15 V/m (the peak SAR 10g value 33.8 mW/kg) could cause changes in the bone. To our knowledge, our study seems to be the first investigation in literature focusing on effects on the bone of 2.45 GHz low-level MW radiation at different EMF values. Additionally, this research is the first article to determine the level of thermal and non-thermal effects on bone.

Open access paper: http://www.biomed.cas.cz/physiolres/pdf/73/73_157.pdf

--

The neuroprotective effects of baobab and black seed on the rat hippocampus exposed to a 900-MHz electromagnetic field

Mohamed H, Deniz OG, Kaplan S. The neuroprotective effects of baobab and black seed on the rat hippocampus exposed to a 900-MHz electromagnetic field. J Chem Neuroanat. 2024 Mar 4;137:102405. doi: 10.1016/j.jchemneu.2024.102405.

Abstract

This study investigated the potential effects on the hippocampus of electromagnetic fields (EMFs) disseminated by mobile phones and the roles of baobab (Adansonia digitata) (AD) and black seed (Nigella sativa) (BS) in mitigating these. Fifty-six male, 12-week-old Wistar albino rats were divided into eight groups of seven animals each. No EMF exposure was applied to the control, AD or BS groups, while the rats in the Sham group were placed in an EMF system with no exposure. A 900-MHz EMF was applied to the EMF+AD, EMF+BS, EMF+AD+BS and EMF groups for 1 hour a day for 28 days. Pyramidal neurons in the hippocampus were subsequently counted using the optical fractionator technique, one of the unbiased stereological methods. Tissue sections were also evaluated histopathologically under light and electron microscopy. The activities of the enzymes catalase (CAT) and superoxide dismutase (SOD) were also determined in blood serum samples. Analysis of the stereological data revealed no statistically significant differences between the EMF and control or sham groups in terms of pyramidal neuron

numbers ($p>0.05$). However, stereological examination revealed a crucial difference in the entire hippocampus between the control group and the AD ($p<0.01$) and BS ($p<0.05$) groups. Moreover, exposure to 900-MHz EMF produced adverse changes in the structures of neurons at histopathological analysis. Qualitative examinations suggest that a combination of herbal products such as AD and BS exerts a protective effect against such EMF side-effects.

https://pubmed.ncbi.nlm.nih.gov/38447905/

Conclusion

We suggest that using appropriate quantities of natural antioxidants in combination with foodstuffs can inhibit or reduce the harmful effects of EMF radiation on the neurons of the brain. The human population, and especially children, should also be protected against exposure to radiation, especially that emitted from mobile phones. To the best of our knowledge, no prior study has demonstrated the effect of AD and BS in the EMF exposed rat hippocampus. Further studies focusing on the effect mechanism of antioxidants, especially AD and BS, that may represent novel protective substances against the side-effects of EMF radiation in the hippocampus, are now needed. Research involving different methods, durations and doses is therefore required.

--

Health and environmental effects to wildlife from radio telemetry and tracking devices—state of the science and best management practices

Manville AM, Levitt BB, Lai HC. Health and environmental effects to wildlife from radio telemetry and tracking devices—state of the science and best management practices. Frontiers in Veterinary Science, 11. 2024. doi: 10.3389/fvets.2024.1283709.

Abstract

This paper discusses the potential health risks and benefits to tagged wildlife from the use of radio tracking, radio telemetry, and related microchip and data-logger technologies used to study, monitor and track mostly wildlife in their native habitats. Domestic pets, especially canids, are briefly discussed as radio-tagging devices are also used on/in them. Radio tracking uses very high frequency (VHF), ultra-high frequency (UHF), and global positioning system (GPS) technologies, including via satellites where platform terminal transmitters (PTTs) are used, as well as geo-locating capabilities using satellites, radio-frequency identification (RFID) chips, and passive integrated responder (PIT) tags, among others. Such tracking technologies have resulted in cutting-edge findings worldwide that have served to protect and better understand the behaviors of myriad wildlife species. As a result, scientists, field researchers, technicians, fish and wildlife biologists and managers, plus wildlife and other veterinarian specialists, frequently opt for its use without fully understanding the ramifications to target species and their behaviors. These include negative physiological effects from electromagnetic fields (EMF) to which many nonhuman species are exquisitely sensitive, as well as direct placement/use-attachment impacts from radio collars, transmitters, and implants themselves. This paper provides pertinent studies, suggests best management practices, and compares technologies currently available to those considering and/or using such technologies. The primary focus is on the health and environmental risk/benefit decisions

that should come into play, including ethical considerations, along with recommendations for more caution in the wildlife and veterinarian communities before such technologies are used in the first place.

Open access paper: https://www.frontiersin.org/articles/10.3389/fvets.2024.1283709

--

Electromagnetic fields regulate iron metabolism in living organisms: A review of effects and mechanism

Zhen C, Zhang G, Wang S, Wang J, Shang P. Electromagnetic fields regulate iron metabolism in living organisms: A review of effects and mechanism. Prog Biophys Mol Biol. 2024 Mar 4:S0079-6107(24)00023-3. doi: 10.1016/j.pbiomolbio.2024.03.001.

Abstract

The emergence, evolution, and spread of life on Earth have all occurred in the geomagnetic field, and its extensive biological effects on living organisms have been documented. The charged characteristics of metal ions in biological fluids determine that they are affected by electromagnetic field forces, thus affecting life activities. Iron metabolism, as one of the important metal metabolic pathways, keeps iron absorption and excretion in a relatively balanced state, and this process is precisely and completely controlled. It is worth paying attention to how the iron metabolism process of living organisms is changed when exposed to electromagnetic fields. In this paper, the processes of iron absorption, storage and excretion in animals (mammals, fish, arthropods), plants and microorganisms exposed to electromagnetic field were summarized in detail as far as possible, in order to discover the regulation of iron metabolism by electromagnetic field. Studies and data on the effects of electromagnetic field exposure on iron metabolism in organisms show that exposure profiles vary widely across species and cell lines. This process involves a variety of factors, and the complexity of the results is not only related to the magnetic flux density/operating frequency/exposure time and the heterogeneity of the observed object. A systematic review of the biological regulation of iron metabolism by electromagnetic field exposure will not only contributes to a more comprehensive understanding of its biological effects and mechanism, but also is necessary to improve human awareness of the health related risks of electromagnetic field exposure.

Open access paper:
https://www.sciencedirect.com/science/article/pii/S0079610724000233?via%3Dihub

--

0.263 terahertz irradiation induced genes expression changes in *Caenorhabditis elegans (roundworm)*

Shang S, Gao F, Zhang Q, Song T, Wang W, Liu D, Gong Y, Lu X. 0.263 terahertz irradiation induced genes expression changes in *Caenorhabditis elegans*. iScience. 2024 Mar 2;27(4):109391. doi: 10.1016/j.isci.2024.109391.

Abstract

The biosafety of terahertz (THz) waves has emerged as a new area of concern with the gradual application of terahertz radiation. Even though many studies have been conducted to investigate the

influence of THz radiation on living organisms, the biological effects of terahertz waves have not yet been fully revealed. In this study, *Caenorhabditis elegans* (*C. elegans*) was used to evaluate the biological consequences of whole-body exposure to 0.263 THz irradiation. The integration of transcriptome sequencing and behavioral tests of *C. elegans* revealed that high-power THz irradiation damaged the epidermal ultrastructures, inhibited the expression of the cuticle collagen genes, and impaired the movement of *C. elegans*. Moreover, the genes involved in the immune system and the neural system were dramatically down-regulated by high-power THz irradiation. Our findings offer fresh perspectives on the biological impacts of high-power THz radiation that could cause epidermal damage and provoke a systemic response.

Open access paper: https://www.cell.com/iscience/fulltext/S2589-0042(24)00612-6

--

Environmental Pollution and Risk of Childhood Cancer: A Scoping Review of Evidence from the Last Decade

Navarrete-Meneses MdP, Salas-Labadía C, Gómez-Chávez F, Pérez-Vera P. Environmental Pollution and Risk of Childhood Cancer: A Scoping Review of Evidence from the Last Decade. *International Journal of Molecular Sciences*. 2024; 25(6):3284. doi: 10.3390/ijms25063284.

Abstract

The long-term effects of environmental pollution have been of concern as several pollutants are carcinogenic, potentially inducing a variety of cancers, including childhood cancer, which is a leading cause of death around the world and, thus, is a public health issue. The present scoping review aimed to update and summarize the available literature to detect specific environmental pollutants and their association with certain types of childhood cancer. Studies published from 2013 to 2023 regarding environmental pollution and childhood cancer were retrieved from the PubMed database. A total of 174 studies were eligible for this review and were analyzed. Our search strategy brought up most of the articles that evaluated air pollution (29%) and pesticides (28%). Indoor exposure to chemicals (11%), alcohol and tobacco use during pregnancy (16%), electromagnetic fields (12%), and radon (4%) were the subjects of less research. We found a particularly high percentage of positive associations between prenatal and postnatal exposure to indoor (84%) and outdoor (79%) air pollution, as well as to pesticides (82%), and childhood cancer. Positive associations were found between leukemia and pesticides and air pollution (33% and 27%); CNS tumors and neuroblastoma and pesticides (53% and 43%); and Wilms tumor and other rare cancers were found in association with air pollution (50%). Indoor air pollution was mostly reported in studies assessing several types of cancer (26%). Further studies are needed to investigate the mechanisms underlying the potential associations between indoor/outdoor air pollution and pesticide exposure with childhood cancer risk as more preventable measures could be taken.

Open access paper: https://www.mdpi.com/1422-0067/25/6/3284

--

A broadband multi-frequency microwave combined biological exposure setup

Zhao X, Li Z, Liu X, Wang Y, Dong G, Liu Q, Wang C. A broadband multi-frequency microwave combined biological exposure setup. *Rev. Sci. Instrum.* 1 April 2024; 95 (4): 044702. doi: 10.1063/5.0196908

Abstract

With the rapid popularization of wireless electronic devices, there has been an increasing concern about the impacts of the electromagnetic environment on health. However, most research reports on the biological effects of microwaves have focused on a single frequency point. In reality, people are exposed to complex electromagnetic environments that consist of multiple frequency microwave signals in their daily lives. It is important to investigate whether multi-frequency combined microwave energies have different biological effects compared with single frequency microwave energy. Unfortunately, there are limited reports on this topic due to the lack of suitable platforms for research on multi-frequency microwave energy combined with biological exposure. To address this issue, this study presents a setup that has a very wide working frequency bandwidth and can be compatible with single frequency and multi-frequency microwave combined exposure. Moreover, it can achieve relatively equal exposure to multiple biological samples at any frequency point in the working frequency range, which is crucial for electromagnetic biology research. The experimental results are in good agreement with the simulation results, confirming its capability to facilitate the study of complex electromagnetic environment effects on organisms.

Open access paper: https://pubs.aip.org/aip/rsi/article/95/4/044702/3280194/A-broadband-multi-frequency-microwave-combined

--

The effects of radiofrequency electromagnetic fields exposure on tinnitus, migraine and non-specific symptoms in the general and working population: A systematic review and meta-analysis on human observational studies

My note: Numerous problems have been identified with this paper and other systematic reviews commissioned by the WHO for an upcoming monograph on radio frequency effects.

Röösli M, Dongus S, Jalilian H, Eyers J, Esu E, Oringanje CM, Meremikwu M, Bosch-Capblanch X. The effects of radiofrequency electromagnetic fields exposure on tinnitus, migraine and non-specific symptoms in the general and working population: A systematic review and meta-analysis on human observational studies. Environ Int. 2024 Jan;183:108338. doi: 10.1016/j.envint.2023.108338.

Abstract

Background: Applications emitting radiofrequency electromagnetic fields (RF-EMF; 100 kHz to 300 GHz) are widely used for communication (e.g. mobile phones), in medicine (diathermy) and in industry (RF heaters).

Objectives: The objective is to systematically review the effects of longer-term or repeated local and whole human body radiofrequency electromagnetic field (RF-EMF) exposure on the occurrence of symptoms. Primary hypotheses were tinnitus, migraine and headaches in relation to RF-EMF exposure of the brain, sleep disturbances and composite symptom scores in relation to whole-body RF-EMF exposure.

Methods: Eligibility criteria: We included case-control and prospective cohort studies in the general population or workers estimating local or whole-body RF-EMF exposure for at least one week.

Information sources: We conducted a systematic literature search in various databases including Web of Science and Medline. Risk of bias: We used the Risk of Bias (RoB) tool developed by OHAT adapted to the topic of this review.

Synthesis of results: We synthesized studies using random effects meta-analysis.

Results: Included studies: We included 13 papers from eight distinct cohort and one case-control studies with a total of 486,558 participants conducted exclusively in Europe. Tinnitus is addressed in three papers, migraine in one, headaches in six, sleep disturbances in five, and composite symptom scores in five papers. Only one study addressed occupational exposure.

Synthesis of results: For all five priority hypotheses, available research suggests that RF-EMF exposure below guideline values does not cause symptoms, but the evidence is very uncertain. The very low certainty evidence is due the low number of studies, possible risk of bias in some studies, inconsistencies, indirectness, and imprecision. In terms of non-priority hypotheses numerous exposure-outcome combinations were addressed in the 13 eligible papers without indication for an association related to a specific symptom or exposure source.

Discussion: Limitations of evidence: This review topic includes various challenges related to confounding control and exposure assessment. Many of these aspects are inherently present and not easy to be solved in future research. Since near-field exposure from wireless communication devices is related to lifestyle, a particular challenge is to differentiate between potential biophysical effects and other potential effects from extensive use of wireless communication devices that may compete with healthy behaviour such as sleeping or physical activity. Future research needs novel and innovative methods to differentiate between these two hypothetical mechanisms.

Interpretation: This is currently the best available evidence to underpin safety of RF-EMF. There is no indication that RF-EMF below guideline values causes symptoms. However, inherent limitations of the research results in substantial uncertainty.

Other: Funding: This review was partially funded by the WHO radioprotection programme.

Registration: The protocol for this review has been registered in Prospero (reg no CRD42021239432) and published in Environment International (Röösli et al., 2021).

Declaration of competing interest The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. Martin Röösli's research is entirely funded by public or not-for-profit foundations. He has served as advisor to a number of national and international public advisory and research steering groups concerning the potential health effects of exposure to nonionizing radiation, including the World Health Organization, the International Agency for Research on Cancer, the International Commission on Non-Ionizing Radiation Protection, the Swiss Government (member of the working group "mobile phone and radiation" and chair of the expert group BERENIS), the German Radiation Protection Commission

(member of the committee Non-ionizing Radiation (A6) and member of the working group 5G (A630)) and the Independent Expert Group of the Swedish Radiation Safety Authority.

https://www.sciencedirect.com/science/article/pii/S0160412023006116?via%3Dihub

--

Effects of radiofrequency electromagnetic field (RF-EMF) exposure on male fertility: A systematic review of experimental studies on non-human mammals and human sperm in vitro (SR 4)

My note: Numerous problems have been identified with this paper and other systematic reviews commissioned by the WHO for an upcoming monograph on radio frequency effects.

Cordelli E, Ardoino L, Benassi B, Consales C, Eleuteri P, Marino C, Sciortino M, Villani P, Brinkworth MH, Chen G, McNamee JP, Wood AW, Belackova L, Verbeek J, Pacchierotti F. Effects of radiofrequency electromagnetic field (RF-EMF) exposure on male fertility: A systematic review of experimental studies on non-human mammals and human sperm in vitro. Environment International. 2024, doi: 10.1016/j.envint.2024.108509. Open access paper:
https://www.sciencedirect.com/science/article/pii/S0160412024000953

Highlights

Risk of bias, inconsistency, publication bias weakened the certainty of results
RF-EMF is unlike to decrease the fecundity of exposed male rodents.
RF-EMF may affect testicular tissue and sperm quality but the evidence is uncertain.
Impact on surrogate markers of fertility may not translate into functional effects.

Abstract

Background The World Health Organization is coordinating an international project aimed at systematically reviewing the evidence regarding the association between radiofrequency electromagnetic field (RF-EMF) exposure and adverse health effects. Reproductive health outcomes have been identified among the priority topics to be addressed.

Objectives To evaluate the effect of RF-EMF exposure on male fertility of experimental mammals and on human sperm exposed in vitro.

Methods Three electronic databases (PubMed, Scopus and EMF Portal) were last searched on September 17, 2022. Two independent reviewers screened the studies, which were considered eligible if met the following criteria: 1) Peer-reviewed publications of sham controlled experimental studies, 2) Non-human male mammals exposed at any stage of development or human sperm exposed in vitro, 3) RF-EMF exposure within the frequency range of 100 kHz-300 GHz, including electromagnetic pulses (EMP), 4) one of the following indicators of reproductive system impairment:•decrease of fertility: rate of infertile males, rate of nonpregnant females, litter size and in vitro fertilization rate;•effects on semen quality: in animal studies sperm count, in both animal and in vitro studies sperm vitality, morphology and DNA/chromatin alterations;•reproductive organ toxicity: testis-epididymis weight, testis or epididymis histology, testis histomorphometry, testicular cell death, estimated testicular cell

production;•hormonal effects: testosterone level.Two reviewers extracted study characteristics and outcome data. We assessed risk of bias (RoB) using the Office of Health Assessment and Translation (OHAT) guidelines. We categorized studies into 3 levels of overall RoB: low, some or high concern. We pooled study results in a random effects meta-analysis comparing average exposure to no-exposure and in a dose–response meta-analysis using all exposure doses. For experimental animal studies, we conducted subgroup analyses for species, Specific Absorption Rate (SAR) and temperature increase. We grouped studies on human sperm exposed in vitro by the fertility status of sample donors and SAR. We assessed the certainty of the evidence using the GRADE approach after excluding studies that were rated as "high concern" for RoB.

Results One-hundred and seventeen papers on animal studies and 10 papers on human sperm exposed in vitro were included in this review. Only few studies were rated as "low concern" because most studies were at RoB for exposure and/or outcome assessment. Subgrouping the experimental animal studies by species, SAR, and temperature increase partly accounted for the heterogeneity of individual studies in about one third of the meta-analyses. In no case was it possible to conduct a subgroup analysis of the few human sperm in vitro studies because there were always 1 or more groups including less than 3 studies. Among all the considered endpoints, the meta-analyses of animal studies provided evidence of adverse effects of RF-EMF exposure in all cases but the rate of infertile males and the size of the sired litters. The assessment of certainty according to the GRADE methodology assigned a moderate certainty to the reduction of pregnancy rate and to the evidence of no-effect on litter size, a low certainty to the reduction of sperm count, and a very low certainty to all the other meta-analysis results. Studies on human sperm exposed in vitro indicated a small detrimental effect of RF-EMF exposure on vitality and no-effect on DNA/chromatin alterations. According to GRADE, a very low certainty was attributed to these results. The few studies that used EMP exposure did not show effects on the outcomes. A low to very low certainty was attributed to these results.

Discussion Many of the studies examined suffered of severe limitations that led to the attribution of uncertainty to the results of the meta-analyses and did not allow to draw firm conclusions on most of the endpoints. Nevertheless, the associations between RF-EMF exposure and decrease of pregnancy rate and sperm count, to which moderate and low certainty were attributed, are not negligible, also in view of the indications that in Western countries human male fertility potential seems to be progressively declining. It was beyond the scope of our systematic review to determine the shape of the dose–response relationship or to identify a minimum effective exposure level. The subgroup and the dose–response fitting analyses did not show a consistent relationship between the exposure levels and the observed effects. Notably, most studies evaluated RF-EMF exposure levels that were higher than the levels to which human populations are typically exposed, and the limits set in international guidelines. For these reasons we cannot provide suggestions to confirm or reconsider current human exposure limits. Considering the outcomes of this systematic review and taking into account the limitations found in several of the studies, we suggest that further investigations with better characterization of exposure and dosimetry including several exposure levels and blinded outcome assessment were conducted. Protocol registration: Protocols for the systematic reviews of animal studies and of human sperm in vitro studies were published in Pacchierotti et al., 2021. The former was also registered in PROSPERO (CRD42021227729 https://www.crd.york.ac.uk/prospero/display_record.php?RecordID = 227729) and the latter in Open Science Framework (OSF Registration DOI https://doi.org/10.17605/OSF.IO/7MUS3).

Excerpts

4.4. Implications for policy and research

In conclusion, our systematic review and *meta*-analyses indicate a possible detrimental effect of RF-EMF exposure on pregnancy rate and sperm count in experimental mammals, whereas the *meta*-analysis of data on litter size was consistent with null.

Although sperm count is not a functional indicator of male fertility, it is a well-standardised analysis routinely applied in clinical andrology. RF-EMF emitting devices are widely applied and epidemiological surveys seem to indicate that, in Western countries, male fertility potential is declining (Auger et al., 2022, Boulicault et al., 2022, Levine et al., 2017). For these reasons the results of our meta-analyses should not be overlooked at a policy level.

It was beyond the scope of our systematic review to determine the shape of the dose–response relationship or to identify a minimum effective exposure level. For these reasons, we cannot provide suggestions to confirm or reconsider current human exposure limits. Nevertheless, it is of note that most studies on male fertility, semen quality and reproductive organ toxicity investigated exposure levels which were rather high with respect to those relevant for human populations: 75–80 % tested exposure levels above 0.4 W/kg (ICNIRP basic restriction for workers) and 46–53 % tested exposure levels above 4 W/kg (ICNIRP health effect level) (ICNIRP, 2020). Thus, it is not known the extent to which the conclusions of the SR meta-analysis can be applied to human exposure levels. Similarly, it is unknown how much our conclusion can be extrapolated to frequencies below 100 MHz and above 10000 MHz, for which only very few studies were retrieved.

During the systematic review, we identified several methodological limitations in the studies that should be overcome to improve the quality of future research. In particular, blinding during experiment performance and outcome assessment should always be applied to minimize bias, an adequate number of cytological or histological preparations should be analysed, automated methods of analysis should be applied whenever possible, a more standardized and complete reporting of technical methods and results should be adopted. Many studies had to be excluded from the systematic review because of insufficient exposure characterization and a large proportion of included studies were rated at either ‘some’ or ‘high concern’ for RoB for similar reasons. We would recommend that future studies bear the reasons for exclusion or RoB concerns in mind in study design and implementation. There are several papers in the research literature with recommendations on how exposure characterisation concerns can be mitigated, for example Kuster and Schonborn (2000). Finally, studies investigating not just a single level but several exposure levels, spanning from low levels comparable to human exposure to higher levels where mild hyperthermic effects could be expected, should be conducted under the same experimental conditions and target tissue temperature monitoring should be employed.

As a final suggestion for future research, we consider it a priority to obtain a scientifically solid database of possible RF-EMF effects on the best predictive surrogate markers of male infertility in experimental rodents. Based on the results of this research, the possibility of testing directly the RF-EMF impact on male reproductive performance could be considered. In view of the limitations of the approach applying *in vitro* exposure of human sperm, we do not recommend further studies of this kind. Conversely, we suggest exploiting semen quality analysis in human biomonitoring investigations of RF-EMF exposed populations....

Other reviews assessing the impact of RF-EMF exposure on male fertility have recently been published, but these only partially assessed the available literature data (Kesari et al., 2018, Sciorio et al., 2022, Sterling et al., 2022, Vornoli et al., 2019). The few recently published systematic reviews on this topic suffered from some methodological limitations such as the lack of a Risk of Bias analysis (Jaffar et al.,

2019, Kim et al., 2021), they limited analysis to only assess effects on semen parameters or were limited to exposure conditions relevant to mobile phone exposures thereby imposing a SAR cut-off (Yu et al., 2021). International committees on human health protection from electromagnetic fields were unable to draw firm conclusions on the possibility of an adverse effect of RF-EMF on male fertility at exposure levels where humans are typically exposed (ICNIRP, 2020, SCENIHR, 2015)....

We considered only original, controlled experimental studies published in peer-reviewed journals. We excluded non-experimental studies (e.g., human epidemiologic or other observational studies), and studies of exposure of both males and females of a mating pair (additional decision and change from protocol, see Section 4.5.2). We excluded papers reporting reviews, opinions, proceedings or meeting abstracts. We did not impose any year-of-publication or language restriction....

For each endpoint, we first conducted a meta-analysis of exposed vs sham control comparisons. When a study had several exposure groups matched to the same comparator, the means and standard deviations of these exposed groups were combined into one exposed group using the formulas provided in the paragraph 6.5.2.10 of the Cochrane Handbook (Higgins and Li, 2022), so that each study was entered only once into the meta-analysis. The exposure level assigned to that combined exposed group was calculated as the average SAR of the exposed groups in that study weighed by the number of animals in each exposed group. In the forest plots this is indicated with an asterisk after the study ID. Studies that compared each exposed group to another separate sham control group were entered as separate studies in the meta-analysis. When multiple studies were reported in the same paper, this is indicated with a number after the study ID in the forest plot....

All data subject to a *meta*-analysis were graphically synthesized by forest plots. A forest plot was drawn in which the studies were divided according to their overall RoB level as “low or some concern” or “high concern”. We decided to exclude from the assessment of the pooled effect sizes the studies rated at “high concern” for RoB in order to draw conclusions based upon the most robust data (see Section 4.5.2)....

After reading the full text, 175 papers on animal studies were excluded. They are listed in Supplementary File 1a with a justification of the exclusion rationale together with those not retrieved or not translated. Over 45 % of the animal studies were excluded because essential information was missing regarding exposure set-up and/or dosimetry, e.g., details on how the exposure system output was established and maintained or exposure frequency. A further 27 % of the studies were excluded because outcome data were deemed out-of-scope or invalid....

Regarding studies on human sperm in vitro, we excluded 33 papers after reading the full text (Supplementary File 1b). Most papers were excluded because they did not report peer-reviewed original results. Other papers could not be included in the systematic review because exposure conditions and/or dosimetry were insufficiently reported or because the exposure conditions did not provide a sufficient exposure contrast between RF-EMF exposed and sham-exposed samples....

4.1. Summary of the evidence and interpretation of the results From experimental animal studies there is moderate certainty of evidence that RF-EMF exposure reduces rate of pregnancy, moderate certainty of evidence that exposure does not reduce litter size, and low certainty of evidence that exposure lowers sperm count. All other results of animal studies and all results on human sperm exposed *in vitro* have very low certainty. We retrieved few independent studies reporting male reproductive effects after experimental animal exposure to EMP. For this source of exposure, results on pregnancy rate, litter size

and sperm count, all consistent with null, have a low certainty. All other results have a very low certainty.

It can be asked whether the results of our *meta*-analyses are consistent with the hypothesis that higher exposure levels, especially those inducing an hyperthermic effect, are more biologically effective than lower exposure levels. The result on the decrease of pregnancy rate is consistent with this hypothesis, as shown by the observation that the pooled effect size is statistically significant only in the subgroup of studies exposed to SAR equal to or higher than 5 W/kg and the statistically significant slope of the linear dose–response relationship. On the other hand, the results on sperm count do not show an increase of the detrimental effect with increasing SAR and all the models of dose–response relationship tested fit the data poorly. Also for other endpoints (the results of which were rated at very low certainty), a direct relationship between the effect and the exposure level is not evident by the subgroup and dose–response analyses and, in some cases, even the possibility of an inverse relationship is suggested by the data. However, this suggestion is not sustained by a solid adverse outcome pathway, and, in some cases, it is based only on few independent studies. We tested if other variables unequally distributed among the subgroups could have a role in increasing the heterogeneity of the observed results and could confound any underlying dose–effect relationship. Indeed, we showed that the absence of blinding during outcome assessment could strongly influence the results for those endpoints that were not measured by automated methods, thus supporting this hypothesis.

4.2. Limitations in the evidence Of all the papers included in the database of animal studies after the title/abstract evaluation, about 60 % had to be excluded for different reasons, with poor exposure characterization accounting for about 45 % of them....

Financial support This project was partially funded by the World Health Organization (contracts 2020/1026306–0, 2022/1275453–1). WHO provided the basis for the protocol and methodological support throughout the review process. Additional in-kind funds were provided by ENEA, Health Canada and Swinburne University of Technology.

Declaration of competing interest The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments We are grateful to Emilie van Deventer, Maria Rosaria Scarfì and Eric van Rongen for advice regarding the protocols draft and for discussions to ensure consistency in approaches across the multiple ongoing WHO systematic reviews. We wish to thank Flavio Di Marzio for his appreciated graphical help.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024000953

--

Mobile phone use and brain tumour risk – COSMOS, a prospective cohort study

My note: This cohort study suffers from numerous methodological problems that undermine the authors' results and conclusions.

Feychting M, Schüz J, Toledano MB, Vermeulen R, Auvinen A, Poulsen AH, Deltour I, Smith RB, Heller J, Kromhout H, Huss A, Johansen C, Tettamanti G, Paul Elliott P. Mobile phone use and brain tumour risk – COSMOS, a prospective cohort study. Environment International. 2024, doi: 10.1016/j.envint.2024.108552.

Abstract

Background Each new generation of mobile phone technology has triggered discussions about potential carcinogenicity from exposure to radiofrequency electromagnetic fields (RF-EMF). Available evidence has been insufficient to conclude about long-term and heavy mobile phone use, limited by differential recall and selection bias, or crude exposure assessment. The Cohort Study on Mobile Phones and Health (COSMOS) was specifically designed to overcome these shortcomings.

Methods We recruited participants in Denmark, Finland, the Netherlands, Sweden, and the UK 2007–2012. The baseline questionnaire assessed lifetime history of mobile phone use. Participants were followed through population-based cancer registers to identify glioma, meningioma, and acoustic neuroma cases during follow-up. Non-differential exposure misclassification was reduced by adjusting estimates of mobile phone call-time through regression calibration methods based on self-reported data and objective operator-recorded information at baseline. Hazard ratios (HR) and 95% confidence intervals (CI) for glioma, meningioma, and acoustic neuroma in relation to lifetime history of mobile phone use were estimated with Cox regression models with attained age as the underlying time-scale, adjusted for country, sex, educational level, and marital status.

Results 264,574 participants accrued 1,836,479 person-years. During a median follow-up of 7.12 years, 149 glioma, 89 meningioma, and 29 incident cases of acoustic neuroma were diagnosed. The adjusted HR per 100 regression-calibrated cumulative hours of mobile phone call-time was 1.00 (95 % CI 0.98–1.02) for glioma, 1.01 (95 % CI 0.96–1.06) for meningioma, and 1.02 (95 % CI 0.99–1.06) for acoustic neuroma. For glioma, the HR for ≥ 1908 regression-calibrated cumulative hours (90th percentile cut-point) was 1.07 (95 % CI 0.62–1.86). Over 15 years of mobile phone use was not associated with an increased tumour risk; for glioma the HR was 0.97 (95 % CI 0.62–1.52).

Conclusions Our findings suggest that the cumulative amount of mobile phone use is not associated with the risk of developing glioma, meningioma, or acoustic neuroma.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0160412024001387

--

Headache in the international cohort study of mobile phone use and health (COSMOS) in the Netherlands and the United Kingdom

My note: This cohort study suffers from numerous methodological problems that undermine the authors' results and conclusions.

Traini E, Smith RB, Vermeulen R, Kromhout H, Schüz J, Feychting M, Auvinen A, Poulsen AH, Deltour I, Muller DC, Heller J, Tettamanti G, Elliott P, Huss A, Toledano MB. Headache in the international cohort study of mobile phone use and health (COSMOS) in the Netherlands and the United Kingdom. Environmental Research. 2024. doi: 10.1016/j.envres.2024.118290.

Highlights

- Prospective study exploring the relationship between mobile phone use and headache.
- Associations with call-time were largely explained by texting.
- Associations likely to reflect behavioural aspects of mobile phone use.

Abstract

Headache is a common condition with a substantial burden of disease worldwide. Concerns have been raised over the potential impact of long-term mobile phone use on headache due to radiofrequency electromagnetic fields (RF-EMFs). We explored prospectively the association between mobile phone use at baseline (2009–2012) and headache at follow-up (2015–2018) by analysing pooled data consisting of the Dutch and UK cohorts of the Cohort Study of Mobile Phone Use and Health (COSMOS) (N = 78,437). Frequency of headache, migraine, and information on mobile phone use, including use of hands-free devices and frequency of texting, were self-reported. We collected objective operator data to obtain regression calibrated estimates of voice call duration. In the model mutually adjusted for call-time and text messaging, participants in the high category of call-time showed an adjusted odds ratio (OR) of 1.04 (95 % CI: 0.94–1.15), with no clear trend of reporting headache with increasing call-time. However, we found an increased risk of weekly headache (OR = 1.40, 95 % CI: 1.25–1.56) in the high category of text messaging, with a clear increase in reporting headache with increasing texting. Due to the negligible exposure to RF-EMFs from texting, our results suggest that mechanisms other than RF-EMFs are responsible for the increased risk of headache that we found among mobile phone users.

Excerpts

A study conducted in Sweden and Finland as part of COSMOS found limited evidence for an association between weekly headache and the highest level of mobile phone use and no clear trend with increasing call-time (Auvinen et al., 2019). The association of headache with call-time appeared stronger for calls via the Universal Mobile Telecommunication System (UMTS) (3G) network than via the older Global System for Mobile (GSM) (2G) telecommunications technology, despite the latter involving higher RF-EMF exposure levels to the head (van Wel et al., 2021) ....

Headache has been linked to excessive mobile phone use, but the mechanism by which mobile phone use may cause symptoms is not properly understood (Wang et al., 2017; Cerutti et al., 2016; Frey, 1998; Hocking, 1998; Oftedal et al., 2000; Schoeni et al., 2015). Previous research in adolescents has suggested that other exposures related to mobile phone use, but not exposure to RF-EMFs, should be considered the causal factor for various symptoms, as the strongest associations were found with activities that cause minimal RF-EMF exposure to the head, such as texting or gaming (Schoeni et al., 2017). Other studies have indicated that stress or unfavourable usage, such as late-night use, may be associated with an increase in reported health symptoms, such as headache (Szyjkowska et al., 2014; Röösli, 2008; Thomée et al., 2011). It is therefore crucial to distinguish between using a mobile phone for calling and other activities that expose the brain to RF-EMFs at lower levels, such as Internet browsing (Cabré-Riera et al., 2022a; SSM's Scientific Council on Electromagnetic Fields, 2020).

Our study attempted to disentangle the exposure-outcome gradient by considering call-time as a proxy for RF-EMF exposure and texting as a proxy for usage with negligible RF-EMF exposure to the brain (Wall

et al., 2019). This study's mobile phone usage data was gathered between 2009 and 2012. During those years, texting was the most popular activity unrelated to RF-EMF exposure.....

The Headache Impact Test (HIT-6) score with a cut-off of 56 points defined severe weekly headache. The HIT-6 is a tool used to measure the impact headaches have on one's ability to function in various aspects of daily life, including work, school, home, and social contexts. The score, ranging from 36 to 78 points, provides a measure of the degree to which headaches affect daily life and functioning, with higher scores indicating a more significant impact on the participant's overall life (Kosinski et al., 2003)....

Results from two-exposure models mutually adjusting for both call-time and texting at baseline, showed substantially lower risk estimates for weekly headache in the high call-time (RC-hfa) category (OR = 1.04, 95 % CI: 0.94–1.15), and no evidence of a trend (P trend = 0.292) (Table 2). Associations with texting were robust to adjustment for call-time: we observed an increased risk of weekly headache in the high category of texting (OR = 1.40, 95 % CI: 1.25–1.56) and a trend of increasing risk with increasing texting frequency (P trend<0.001), in line with results from the single-exposure model (Table 3)....

In this large international prospective cohort of mobile phone users in the Netherlands and the UK, mobile phone use for calling and texting at baseline was associated with headaches at follow-up. Mutually adjusting for both call-time and texting considerably attenuated risk estimates for call-time, while associations with texting were still strong and robust to adjustment, with a clear exposure-outcome gradient.

Headache has been linked to excessive mobile phone use, but the mechanism by which mobile phone use may cause symptoms is not properly understood (Wang et al., 2017; Cerutti et al., 2016; Frey, 1998; Hocking, 1998; Oftedal et al., 2000; Schoeni et al., 2015). Previous research in adolescents has suggested that other exposures related to mobile phone use, but not exposure to RF-EMFs, should be considered the causal factor for various symptoms, as the strongest associations were found with activities that cause minimal RF-EMF exposure to the head, such as texting or gaming (Schoeni et al., 2017)....

Our study attempted to disentangle the exposure-outcome gradient by considering call-time as a proxy for RF-EMF exposure and texting as a proxy for usage with negligible RF-EMF exposure to the brain (Wall et al., 2019). This study's mobile phone usage data was gathered between 2009 and 2012. During those years, texting was the most popular activity unrelated to RF-EMF exposure.

In both scenarios, we found an increased risk of headache in the high exposure category of mobile phone use with a positive exposure-outcome gradient confirmed by the test for trend. The attenuated risk estimates for call-time in the mutually adjusted model argue against an effect of exposure to RF-EMFs due to the negligible exposure attributed to texting. This conclusion is also supported by comparing call-time analyses with and without hands-free adjustment, where no risk reduction was found among users for the adjusted exposure metrics....

Regarding secondary health outcomes, we found consistent patterns of results for severe weekly headache and migraine at follow-up in terms of increased risk estimates and significant trends. Increasing risk of daily headache was associated with increasing texting (P trend<0.001) but not with increasing call-time (P trend = 0.448) (Table 2, Table 3).

In both scenarios, we found an increased risk of headache in the high exposure category of mobile phone use with a positive exposure-outcome gradient confirmed by the test for trend. The attenuated risk estimates for call-time in the mutually adjusted model argue against an effect of exposure to RF-EMFs due to the negligible exposure attributed to texting. This conclusion is also supported by comparing call-time analyses with and without hands-free adjustment, where no risk reduction was found among users for the adjusted exposure metrics.

In this study, the distribution of the exposure, specifically regarding call-time, differed between Dutch and UK participants. Mobile phone usage behaviour across countries cannot be assumed to be identical due to various factors such as cultural, economic, technological, and market dynamics (Böhm, 2015). To assess the consistency of our findings, we showed that defining the top exposure category for call-time based on the 80th percentile cut-off, thereby ensuring the inclusion of Dutch participants in the "high" exposure category, yielded results consistent with those obtained using the 90th percentile as a cut-off. These findings suggested that the association we found between call-time and headache was driven not only by UK but also Dutch participants. Of note, all analyses were adjusted for country of residence....

Our study also has limitations. First, we did not have information about "true" RF-EMF exposure. Exposure to RF-EMFs emitted by wireless devices is difficult to quantify, particularly in large populations and over extended periods, as it depends on different factors, such as reception quality or other factors influencing signal strength. In our study, we calculated several exposure metrics as proxies for RF-EMF exposure, which allowed us to estimate the average individual RF-EMF exposure in the population. Additionally, information on other aspects of usage, such as screen time, blue light exposure or unfavourable use at night, may be helpful to include in future studies.

For highly transient and acute symptoms such as headache, using the peak of RF-EMF exposure might be theoretically preferable over the weekly exposure assessed in our study. However, adopting this approach would require substantially different exposure assessment methods that are impractical for large cohort studies, such as asking participants to regularly fill in a detailed usage diary. Given the study design and methodology used to assess RF-EMF exposure in COSMOS, the analysis of the association between RF-EMF peak exposure and reporting of headache symptoms was precluded. In light of the transient nature of headaches, future research may explore the potential effect of peak RF-EMF exposure on symptom onset more thoroughly.

The composition of the Dutch cohort is not representative of the adult population of the Netherlands with respect to sex and age....

Conclusions

In summary, we found that the use of mobile phones, particularly texting, is associated with headaches and migraines, and the associations with call-time were largely explained by texting. As the associations are driven more by text messaging than call-time, they do not appear to be explained by RF-EMF exposure from the mobile device but are likely to reflect lifestyle, other exposures, or behavioural factors associated with mobile phone use.

Declaration of competing interest [Note that text is missing from this paragraph.]

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Maria Feychting reports financial support was provided by Swedish Research Council. Maria Feychting reports financial support was provided by AFA Insurance. Maria Feychting reports financial support was provided by Swedish Research Council for Health, Working Life and Welfare. Maria Feychting reports financial support was provided by Swedish Radiation Safety Authority. Maria Feychting reports financial support was provided by Sweden's Innovation Agency. Paul Elliott reports financial support was provided by Mobile Telecommunications and Health Research. Paul Elliott reports financial support was provided by National Institute for Health and Care Research Health Protection Research Unit. Anssi Auvinen reports financial support was provided by National Technology Agency (TEKES). Anssi Auvinen reports financial support was provided by Pirkanmaa Hospital District. Anssi Auvinen reports financial support was provided by Yrjö Jahnsson Foundation. Anssi Auvinen reports financial support was provided by Mobile Manufacturers Forum. Roel Vermeulen reports financial support was provided by Netherlands Organisation for Health Research and Development. Aslak Harbo Poulsen reports financial support was provided by the Danish Strategic Research Council. Joachim Schuz reports financial support was provided by French Agency for Food, Environmental and Occupational Health & Safety (ANSES). Maria Feychting reports a relationship with World Health Organization that includes:. Maria Feychting reports a relationship with Public Health England Advisory Group on Non-ionizing Radiation that includes:. Maria Feychting reports a relationship with Norwegian Institute of Public Health that includes:. Maria Feychting reports a relationship with Forskningsrådet för hälsa arbetsliv och välfärd that includes:. Maria Feychting reports a relationship with Swedish Radiation Safety Authority that includes:. Maria Feychting reports a relationship with International Commission on Non-Ionizing Radiation Protection that includes:. Aslak Harbo Poulsen reports a relationship with Swedish Radiation Safety Authority that includes:. Anssi Auvinen reports a relationship with International Commission on Non-Ionizing Radiation Protection that includes:. Anssi Auvinen reports a relationship with Scientific Committee on Emerging and Newly Identified Health Risks that includes:. Anssi Auvinen reports a relationship with European Commission, Swedish Radiation Safety Authority that includes:. Anssi Auvinen reports a relationship with World Health Organization that includes:. Anke Huss reports a relationship with Swedish Radiation Safety Authority that includes:. Anke Huss reports a relationship with International Commission on Non-Ionizing Radiation Protection that includes:. Anke Huss reports a relationship with BioEM society that includes:. Anke Huss reports a relationship with Gezondheidsraad that includes:. Anke Huss reports a relationship with Swiss Research Foundation for Electricity and Mobile Communication that includes:. Hans Kromhout reports a relationship with Gezondheidsraad that includes:. Hans Kromhout reports a relationship with World Health Organization that includes:. Mireille B Toledano reports a relationship with UK Committee on Medical Aspects of Radiation in the Environment (COMARE) that includes:.

Open access paper: https://www.sciencedirect.com/science/article/pii/S0013935124001944

--

Personal exposure to radiofrequency electromagnetic fields: A comparative analysis of international, national, and regional guidelines

My note: This paper suffers from problems that undermine the authors' results and conclusions.

Ramirez-Vazquez R, Escobar I, Vandenbosch GAE, Arribas E. Personal exposure to radiofrequency electromagnetic fields: A comparative analysis of international, national, and regional guidelines. Environmental Research. 2024. doi: 10.1016/j.envres.2024.118124.

Highlights

- International, National and Regional exposure limits for RF-EMF.
- RF-EMF limits are based on the ALARA (As Low As Reasonably Achievable) Principle.
- Maximum exposure limit must be scientific related, not political or in any other way.
- New technologies and new smartphones require less intensity than previous model

Abstract

A worldwide overview and analysis for the existing limits of human exposure to Radiofrequency Electromagnetic Fields (RF-EMF) is given in this paper. These reference levels have been established by different national and even regional governments, which can be based on the guidelines provided by the recommendations of the International Commission on Non-Ionizing Radiation Protection (ICNIRP), the International Committee on Electromagnetic Safety of the Institute of Electrical and Electronics Engineers (IEEE), and even in the United States of the Federal Communications Commission (FCC), as well as, are based on the so-called precautionary principle. Explicit reference is made to the exposure limits adopted in countries or regions, such as Canada, Italy, Poland, Switzerland, China, Russia, France, and regions of Belgium (Brussels, Flanders, Wallonia), where the limits are much lower than the international standards. The limits are compared to a selected set of in-situ measurements. This clearly shows that the measured values are typically very small compared to the international standards but could be somewhat higher compared to the reduced limits. Based on this observation and the reasonable assumption that the sensitivity of people to Electromagnetic Fields (EMF) is the same everywhere (whole-body), we propose the idea to establish a worldwide reference limit for the general public, thus applicable in all countries, if the ICNIRP considers it appropriate. Research must continue to generate measurement data that demonstrate the levels of exposure to which we are really exposed, and with this, provide arguments to the organizations that established the guidelines, especially the ICNIRP, to evaluate whether the current limits are too much. High and can be modified when considered pertinent. To the best of our knowledge, at no time has the reference level for the general public been exceeded.

Conclusion

The duality of limits in the same country does not seem appropriate, because people who live in one area or another have more or less the same interaction with RF-EMF waves. Therefore, the limits should be the same, following the ALARA principle – As Low As Reasonably Achievable. The reasons for setting the maximum exposure limit should only be scientific related, not political or in any other way.

In this paper, we provided an overview, analyze, and discuss existing limits to RF-EMF. The international reference levels established by ICNIRP are also recommended by WHO, IEEE and FCC, and are adopted by most countries. However, some countries such as Canada, Italy, Poland, Switzerland, China, Russia, France, and regions of Belgium establish more restrictive limits than the international ones. The case of Belgium is rather specific because it establishes three very strict and different limits in Brussels, Flanders, and Wallonia.

In addition to the international limits, we have commented on the BioInitiative Limit, based on the Precautionary Principle as a preventive action, an excessively restrictive value that leaves aside more than 60 % of the personal exposure studies to RF-EMF.

There should be a trend towards unifying the limits of exposure to RF-EMF in all countries. This would allow decisions to be made in a scientific and consensual manner. The solutions could follow the ones from the International Union for Pure and Applied Physics (IUPAP), which is an international organization that has sixty member countries and is in charge, among other things, of the correct use of the International System of Units and of reviewing the measurement units of the physical magnitudes.

Based on the observation and analysis of measured levels that in most cases, the measured levels are very small, we propose that a reference limit be established for the general public, reasonable and applied to all countries, if the ICNIRP considers it appropriate. Therefore, research must continue to generate measurement data that demonstrate the levels of exposure to which we are really exposed, and with this, provide arguments to the organizations that established the guidelines, especially the ICNIRP, so that they evaluate whether the current limits are too high and can be modified, when considered pertinent. To the best of our knowledge, at no time has the reference level been exceeded for the general public, and we believe that it will not be exceeded, since the new smartphones require much less intensity than previous models, and the same goes for new technologies, 5G, 6G.

https://www.sciencedirect.com/science/article/pii/S0013935124000288

--

How to improve IARC's RF-EMF cancer hazard communication

My note: This paper suffers from problems that undermine the authors' analysis.

Wiedemann PM, Croft RJ. How to improve IARC's RF-EMF cancer hazard communication. Bioelectromagnetics. 2024 Feb 13. doi: 10.1002/bem.22499.

Abstract

A crucial aspect of IARC's evaluation of the relative carcinogenicity of agents is the communication of its conclusions. The present paper addressed the experimental risk perception literature pertaining to IARC's radiofrequency electromagnetic field evaluation communication, and derived specific recommendations for improving it.

Open access paper: https://onlinelibrary.wiley.com/doi/10.1002/bem.22499

--

Molecular biomarkers in Electrohypersensitivity and Multiple Chemical Sensitivity: How They Can Help Diagnosis, Follow-Up, and in Etiopathologic Understanding

Irigaray P, Awaida N, Belpomme D. Molecular biomarkers in Electrohypersensitivity and Multiple Chemical Sensitivity: How They Can Help Diagnosis, Follow-Up, and in Etiopathologic Understanding. Medical Research Archives, [S.l.], 12(1), 2024. doi: 10.18103/mra.v12i1.4771.

Abstract

Electrohypersensitivty (EHS) and multiple chemical sensitivity (MCS) are new worldwide emerging neurologic disorders in the framework of sensitivity-related environmental pathology. We have recently extended and confirmed our previous observation showing that EHS and MCS share clinically identical symptoms and may co-exist as a unique, common, sensitivity-related neurologic syndrome in 25% of the cases. There is presently no published biological study of these disorders, except the one we have previously published as preliminary. In the present study, we show that EHS and MCS and the combined syndrome share identical biochemical changes. More precisely, by measuring levels of peripheral blood and urine molecular biomarkers in a cohort of 2,018 consecutive cases, we show that both disorders and the combined syndrome can be objectively characterized, in about 90% of the cases, by a decrease in the production of 6-hydroxymelatonin sulfate in urine, while in 30-50% they are characterized by increased levels of histamine and of heat shock proteins (HSP) 27 and/or 70, and of protein S100B and nitrotyrosine in the peripheral blood. Increased levels of histamine and HSP are indicators of low grade inflammation while increased levels of protein S100B and nitrotyrosine are indicators of blood-brain barrier disruption/opening. In addition, we show that in about 15% of the cases anti-myelin autoantibodies can be detected in the peripheral blood, accounting for the occurrence of an autoimmune response. Sensitivity, specificity and reproducibility of the biochemical tests are discussed, as well as the role of these indicators used as biomarkers for the diagnosis and follow-up of patients. We also discuss cases with undetectable biological change for which they can be nevertheless diagnosed by cerebral neurotransmitters analysis in urine and brain imaging. On the basis of these biological data it is suggested that EHS and/or MCS are new brain disorders, generated via a common etiopathogenic mechanism.

https://esmed.org/MRA/mra/article/view/4771

--

Environmental Health Ecosystem Sustainability in The Era of Electromagnetic Radiation Contamination

Matthew UO, Bakare KM, Oyekunle D, Nkeiruka AM, Ebong GN (2024) Environmental Health Ecosystem Sustainability in The Era of Electromagnetic Radiation Contamination. J Comm Med and Pub Health Rep 5(02): doi: 10.38207/JCMPHR/2024/JAN05020418.

Abstract

The primary environmental health sub-disciplines that address the best approaches to environmental health, ecosystem sustainability, and natural habitats are environmental science, toxicology, environmental epidemiology, and occupational and environmental medicine. The intimate connection between environmental toxins and human health gained increased public attention at the beginning of the new millennium. Radiation, chemical, and biological agents are the three primary ecological contaminants. There is contamination in the soil, water, food, and air. The increasing adoption of 5G wireless networks has recently raised interest in its potential to support several digital and critical infrastructures. However, concerns have been raised over the potential health risks associated with fifth-generation wireless networks due to the claims that the electromagnetic radiation in the 5G carrier signal is more potent than in any prior networks. As a subject of public health, environmental health studies the whole range of effects that artificial technology and natural environments have on the general well-being of society-wide space. It establishes the prerequisites for a wholesome atmosphere

as essential in managing factors that may negatively impact the sustainability of ecosystems and the environment's well-being. This study observed that the multistage carcinogenic process, which includes carcinogen activation, oxidative DNA damage, and tumor growth, is aided by hydrogen peroxide formation during the breakdown of water molecules due to 5G electromagnetic frequency radiation as an environmental contaminant.

https://www.acquaintpublications.com/article/environmental_health_ecosystem_sustainability_in_the_era_of_electromagnetic_radiation_contamination

--

Assessment of Radio Frequency Electromagnetic Field Exposure Induced by Base Stations in Several Micro-Environments in France

W. B. Chikha et al. Assessment of Radio Frequency Electromagnetic Field Exposure Induced by Base Stations in Several Micro-Environments in France. IEEE Access, vol. 12, pp. 21610-21620, 2024, doi: 10.1109/ACCESS.2024.3363914.

Abstract

Recently, the monitoring of the radiofrequency electromagnetic field (RF-EMF) exposure induced by cellular networks has received a great deal of attention. In this work, a set of 70 microenvironments (MEs) located in urban and rural areas are selected in France under, on the one hand, the French Beyond5G project, and on the other hand, the 5G expOsure, causaL effects and rIsk perception through citizen engagement (GOLIAT) EU project. The purpose of this study is to assess the RF-EMF DL exposure in residential areas, downtowns, business areas, train stations, and public transport rides. For that, we employ the personal ExpoM-RF4 dosimeter placed inside a backpack to perform the measurements in different MEs. To take into consideration the effect of the presence of the human body near the dosimeter, we propose a correction approach that is mainly based on comparing the measurements given by ExpoM-RF4 to the ones provided by a reference system using the Tektronix real-time spectrum analyzer (RTSA) far from the body. Then, we use metrics, such as the quadratic mean, standard deviation, and median of the electric (E) field to carry out a comparative study between different MEs with different RF bands. It was found that the RF-EMF exposure levels for all MEs are well below the maximum allowable exposure limit prescribed by the International Commission on Non-Ionizing Radiation Protection (ICNIRP). In addition, we perform clustering analyses using the K-Means technique to group the MEs with comparable exposure levels. The results show that the exposure level is low, but generally higher in MEs located in Paris than in the other considered areas (i.e., Massy and three villages, namely Igny, Bures-sur-Yvette and Gif-Sur-Yvette). For example, we observe that outdoor MEs can be grouped into three clusters, where the average total E fields (ATEFs) are 0.77 V/m, 0.35 V/m, and 0.08 V/m for the MEs belonging to the first, second and third clusters, respectively. Note that the first cluster here mainly contains the MEs located in Paris. This can be explained by the important number of antennas deployed in that area to serve the huge amount of users. We also observe few locations with exceptions confirming the presence of heterogeneous environments in the vicinity of some areas. For instance, three MEs in Paris among fifteen have an exposure level similar to Massy MEs in outdoor

https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=10424982&isnumber=10380310

--

Electromagnetic Exposure Level of a Pure Electric Vehicle Inverter Based on a Real Human Body

Dong X, Gao Y, Lu M. The Electromagnetic Exposure Level of a Pure Electric Vehicle Inverter Based on a Real Human Body. *Applied Sciences*. 2024; 14(1):32. https://doi.org/10.3390/app14010032.

Abstract

In order to quantitatively analyze the electromagnetic exposure dose of an inverter in a pure electric vehicle to the driver's body and assess the safety of the electromagnetic exposure, based on a real human anatomy model in the virtual home project, a real human model with several organs and tissues, including muscles, bones, a heart, lungs, a liver, kidneys, a bladder, a skull, a scalp, white matter, and a cerebellum, was constructed. The inverter of a pure electric vehicle is considered to be the electromagnetic exposure source; for this study, an equivalent electromagnetic environment model composed of a real human body, an inverter, and a vehicle body was built. The distribution of induced fields in the driver's tissues and organs was calculated and analyzed using the finite element method. The results show that the distribution of the magnetic flux density, induced electric field, and induced current density in the driver's body was affected by the spatial distance of the inverter. The farther the distance was, the weaker the value was. Specifically, due to the different dielectric properties of the different tissues, the induced field in the different tissues was significantly different. However, the maximum magnetic flux density over the space occupied by the driver's body and induced electric field in the driver's trunk and central nervous system satisfied the exposure limits of the International Commission on Non-Ionization Radiation Protection, indicating that the electromagnetic environments generated by the inverter proposed in this paper are safe for the vehicle driver's health. The numerical results of this study could also effectively supplement the study of the electromagnetic environments of pure electric vehicles and provide some references for protecting the drivers of pure electric vehicles from electromagnetic radiation.

Conclusions

Based on the numerical results, we found that although the B-field in the central region of the inverter exceeds the reference level of the ICNIRP for general public exposure, when the distance from the inverter is increased, the B-field values sharply decrease. The maximum B-field over the space occupied by the driver's body and the E-field in the driver's trunk and CNS are all below the exposure limits defined by the ICNIRP for general public exposure. The electromagnetic environment generated by the inverter adopted in this study (12 kW, Semikron, Germany), at its rated state in the paper, is therefore safe and would not affect the PEV driver's health. Additionally, the results of this study could effectively supplement research regarding the electromagnetic environment of PEVs and provide references for the formulation of industry standards for electromagnetic exposure and the design of vehicles that are safe in terms of electromagnetic exposure.

Open access paper: https://www.mdpi.com/2076-3417/14/1/32

--

SAR Estimations in a Child Due to RF Exposures from Several Laptops in a Classroom Environment

Soares NE, Bulla G, Fernandez-Rodriguez CE, de Salles AAA. SAR Estimations in a Child Due to RF Exposures from Several Laptops in a Classroom Environment. *2023 IEEE MTT-S Latin America Microwave Conference (LAMC)*, San José, Costa Rica, 2023, pp. 58-60, doi: 10.1109/LAMC59011.2023.10375589.

Abstract

This research provides a detailed examination of the exposure to electromagnetic fields (EMF) in a classroom environment. The study is driven by the escalating integration of electronic devices in educational settings and the subsequent necessity to estimate the implications of this on the overall exposure. The employed methodology simulates a classroom filled with several 7-year-old girls realistic models, each using a laptop, which are the primary sources of electromagnetic radiation. The 1g and 10g peak spatial Specific Absorption Rate (psSAR) are calculated for various parts of the child model, including the head, back, and hands. It is observed that, when several laptops are used, the psSAR can be substantially increased (e.g. up to 40-fold). One objective of this study is to enhance the understanding of potential EMF exposure effects in relation to age and to provide guidelines for the design of safer classrooms in the context of widespread laptop usage.

https://ieeexplore.ieee.org/document/10375589

--

Effects of 3.5-GHz radiofrequency radiation on energy-regulatory hormone levels in the blood and adipose tissue

Bektas H, Dasdag S, Altindag F, Akdag MZ, Yegin K, Algul S. Effects of 3.5-GHz radiofrequency radiation on energy-regulatory hormone levels in the blood and adipose tissue. *Bioelectromagnetics* 1–9 (2024). doi: 10.1002/bem.22498.

Highlights

- 3.5 GHz radiofrequency radiation (RFR) may induce alterations in hormones regulating energy metabolism.
- 3.5 GHz RFR may lead to alterations in total antioxidant, total oxidant, and hydrogen peroxide levels.
- Particularly in conjunction with diabetes, 3.5 GHz RFR may result in adverse effects on energy metabolism.
- Although there were changes of the hormone levels in the exposed group, the actual values remained for both sham and exposed groups within the normal range.

Abstract

In recent years exposure of living beings to radiofrequency radiation (RFR) emitted from wireless equipment has increased. In this study, we investigated the effects of 3.5-GHz RFR on hormones that regulate energy metabolism in the body. Twenty-eight rats were divided into four groups: healthy sham (n = 7), healthy RFR (n = 7), diabetic sham (n = 7), and diabetic RFR (n = 7). Over a month, each group spent 2 h/day in a Plexiglas carousel. The rats in the experimental group were exposed to RFR, but the sham groups were not. At the end of the experiment, blood and adipose tissues were collected from euthanized rats. Total antioxidant, total oxidant, hydrogen peroxide, ghrelin, nesfatin-1, and irisin were

determined. Insulin expression in pancreatic tissues was examined by immunohistochemical analysis. Whole body specific absorption rate was 37 mW/kg. For the parameters analyzed in blood and fat, the estimated effect size varied within the ranges of 0.215–0.929 and 0.503–0.839, respectively. The blood and adipose nesfatin-1 ($p = 0.002$), blood and pancreatic insulin are decreased, ($p = 0.001$), gherelin ($p = 0.020$), irisin ($p = 0.020$), and blood glucose ($p = 0.040$) are increased in healthy and diabetic rats exposed to RFR. While nesfatin-1 are negatively correlated with oxidative stress, hyperglycemia and insulin, ghrelin and irisin are positively correlated with oxidative stress and hyperglycemia. Thus, RFR may have deleterious effects on energy metabolism, particularly in the presence of diabetes.

https://onlinelibrary.wiley.com/doi/10.1002/bem.22498

--

Genetic profiling of rat gliomas and cardiac schwannomas from life-time radiofrequency radiation exposure study using a targeted next-generation sequencing gene panel

Brooks AM, Vornoli A, Kovi RC, Ton TVT, Xu M, Mashal A, Tibaldi E, Gnudi F, Li JL, Sills RC, Bucher JR, Mandrioli D, Belpoggi F, Pandiri AR. Genetic profiling of rat gliomas and cardiac schwannomas from life-time radiofrequency radiation exposure study using a targeted next-generation sequencing gene panel. PLoS One. 2024 Jan 17;19(1):e0296699. doi: 10.1371/journal.pone.0296699.

Abstract

The cancer hazard associated with lifetime exposure to radiofrequency radiation (RFR) was examined in Sprague Dawley (SD) rats at the Ramazzini Institute (RI), Italy. There were increased incidences of gliomas and cardiac schwannomas. The translational relevance of these rare rat tumors for human disease is poorly understood. We examined the genetic alterations in RFR-derived rat tumors through molecular characterization of important cancer genes relevant for human gliomagenesis. A targeted next-generation sequencing (NGS) panel was designed for rats based on the top 23 orthologous human glioma-related genes. Single-nucleotide variants (SNVs) and small insertion and deletions (indels) were characterized in the rat gliomas and cardiac schwannomas. Translational relevance of these genetic alterations in rat tumors to human disease was determined through comparison with the Catalogue of Somatic Mutations in Cancer (COSMIC) database. These data suggest that rat gliomas resulting from life-time exposure to RFR histologically resemble low grade human gliomas but surprisingly no mutations were detected in rat gliomas that had homology to the human IDH1 p.R132 or IDH2 p.R172 suggesting that rat gliomas are primarily wild-type for IDH hotspot mutations implicated in human gliomas. The rat gliomas appear to share some genetic alterations with IDH1 wildtype human gliomas and rat cardiac schwannomas also harbor mutations in some of the queried cancer genes. These data demonstrate that targeted NGS panels based on tumor specific orthologous human cancer driver genes are an important tool to examine the translational relevance of rodent tumors resulting from chronic/life-time rodent bioassays.

Excerpts

In this study we have demonstrated that the gliomas and cardiac schwannomas in rats resulting from lifetime exposure to low dose far field RFR that are used for cellular telephone communications, are morphologically similar to low grade human gliomas and that about 25% of the mutations seen in these tumors have corresponding alterations in homologs of human cancer genes. Surprisingly, none of the rat

gliomas examined in this study harbored mutations in *Idh1/2* genes that are common in human gliomas....

In summary, our results demonstrate that regardless of their etiology (due to lifetime RFR exposure or arising spontaneously), rat gliomas are primarily *Idh*1/2 wild type unlike most human gliomas. Histologically, most of the rat gliomas resemble diffuse low-grade gliomas in humans and such gliomas that do not harbor *IDH1/2* mutations in humans are known to have poor prognosis. The genetic alterations in other cancer genes evaluated in this panel provide novel insights into tumor progression in rat gliomas and cardiac schwannomas. The relevance of specific mutations to human cancers is variable, with some genes (*Tp53*, *Cdkn2a*, *Erbb2*, *Chek2*, *Kras* and *Pik3r1*) harboring many alterations with COSMIC relevance while the opposite is true for other target genes (*Idh1/2*, *Atrx*, *Notch1*, *Pten*, *Rb1* and *Setd2*). Several of these conserved mutations in rat tumors do not have comparable alterations in the COSMIC database, suggesting that the orthologous mutations could have different functional consequences in rat carcinogenesis and deserve further study. An important consideration is that molecular differences underlying mutational processes contribute to distinct mutational patterns which could be the result of similar etiology, albeit by different mechanisms.

Several of the variants that were detected in gliomas were also observed in non-tumor brain tissues from interim time point providing an insight into the molecular pathogenesis in rodent carcinogenicity studies and these strategies may be utilized to potentially estimate the cancer hazard risk in shorter term animal studies. Finally, this targeted mutation panel may be refined using data from whole genome or exome sequencing of rat tumors and performing error corrected duplex sequencing to increase the sensitivity to detect rare mutations in exposed non-tumor tissues from early time points.

Open access paper: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0296699

--

Understanding the public voices and researchers speaking into the 5G narrative

Weller S, McCredden J. Understanding the public voices and researchers speaking into the 5G narrative. Front. Public Health. Vol. 11. 2023 doi: 10.3389/fpubh.2023.1339513.

Abstract

The many different voices speaking into the current narrative surrounding the health effects of 5G technologies necessitate an exploration of the background of the various published author spokespersons and their potential motives. This has been attempted recently by de Vocht and Albers. However, that opinion piece used a narrow investigative lens, resulting in an undermining of both the rationality of the concerned general public and the motives of specific researchers. At the same time, biases, conflicts of interest, and flaws found in "independent" reviews were not considered. To address these oversights, an evidence-based appraisal of public opinion and the scientific caliber of authors involved in the 5G health discussion is warranted. Subsequently, this review article presents an analysis of the available Australian data representing public voices, while also conducting a broader investigation of the level of expertise of the recent author-spokespersons based on their experience as scientists, particularly in the area of health effects of radiofrequency electromagnetic fields. This review thus attempts to more clearly illustrate for the reader the caliber and motives of the voices speaking into the

5G narrative. The article concludes with a set of questions that need to be answered to enable scientists to advise policy makers more effectively on matters of 5G and public health.

Open access paper: https://www.frontiersin.org/articles/10.3389/fpubh.2023.1339513/abstract

--

Effects of 700 and 3500 MHz 5G radiofrequency exposure on developing zebrafish embryos

Torres-Ruiz M, Suárez OJ, López V, Marina P, Sanchis A, Liste I, de Alba M, Ramos V. Effects of 700 and 3500 MHz 5G radiofrequency exposure on developing zebrafish embryos. Sci Total Environ. 2024 Jan 8:169475. doi: 10.1016/j.scitotenv.2023.169475.

Abstract

Telecommunications industries are rapidly deploying the fifth generation (5G) spectrum and there is public concern about the safety and health impacts of this type of Radio Frequency Radiation (RFR), in part because of the lack of comparable scientific evidence. In this study we have used a validated commercially available setting producing a uniform field to expose zebrafish embryos (ZFe) to unmodulated 700 and 3500 MHz frequencies. We have combined a battery of toxicity, developmental and behavioral assays to further explore potential RFR effects. Our neurobehavioral profiles include a tail coiling assay, a light/dark activity assay, two thigmotaxis anxiety assays (auditory and visual stimuli), and a startle response - habituation assay in response to auditory stimuli. ZFe were exposed for 1 and 4 h during the blastula period of development and endpoints evaluated up to 120 h post fertilization (hpf). Our results show no effects on mortality, hatching or body length. However, we have demonstrated specific organ morphological effects, and behavioral effects in activity, anxiety-like behavior, and habituation that lasted in larvae exposed during the early embryonic period. A decrease in acetylcholinesterase activity was also observed and could explain some of the observed behavioral alterations. Interestingly, effects were more pronounced in ZFe exposed to the 700 MHz frequency, and especially for the 4 h exposure period. In addition, we have demonstrated that our exposure setup is robust, flexible with regard to frequency and power testing, and highly comparable. Future work will include exposure of ZFe to 5G modulated signals for different time periods to better understand the potential health effects of novel 5G RFR.

https://pubmed.ncbi.nlm.nih.gov/38199355/

--

Effects of radiofrequency field from 5G communication on fecal microbiome and metabolome profiles in mice

Wang, X., Zhou, G., Lin, J. *et al.* Effects of radiofrequency field from 5G communication on fecal microbiome and metabolome profiles in mice. *Sci Rep* **14**, 3571 (2024). https://doi.org/10.1038/s41598-024-53842-2

Abstract

With the rapid development of 5G networks, the influence of the radiofrequency field (RF) generated from 5G communication equipment on human health is drawing increasing attention in public. The study aimed at assessing the effects of long-term exposure to 4.9 GHz (one of the working frequencies of 5G communication) RF field on fecal microbiome and metabolome profiles in adult male C57BL/6 mice. The animals were divided into Sham group and radiofrequency group (RF group). For RF group, the mice were whole body exposed to 4.9 GHz RF field for three weeks, 1 h/d, at average power density (PD) of 50 W/m^2. After RF exposure, the mice fecal samples were collected to detect gut microorganisms and metabolites by 16S rRNA gene sequencing and LC–MS method, respectively. The results showed that intestinal microbial compositions were altered in RF group, as evidenced by reduced microbial diversity and changed microbial community distribution. Metabolomics profiling identified 258 significantly differentially abundant metabolites in RF group, 57 of which can be classified to Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways. Besides, functional correlation analysis showed that changes in gut microbiota genera were significantly correlated with changes in fecal metabolites. In summary, the results suggested that altered gut microbiota and metabolic profile are associated with 4.9 GHz radiofrequency exposure.

Open access paper: https://www.nature.com/articles/s41598-024-53842-2

--

Estimation of SAR Average in Rats during 5G NR Chronic Exposure

Makhmanazarov R, Tseplyaev I, Shipilov S, Krivova N. Estimation of SAR Average in Rats during 5G NR Chronic Exposure. *Applied Sciences*. 2024; 14(1):208. doi: 10.3390/app14010208.

Abstract

To study physiological reactions in the brain and skin of higher mammals exposed to chronic radiofrequency radiation, specific absorption ratio (SAR) determination is required and time-consuming numerical methods are used. The paper deals with the estimation of the whole-body specific absorption rate (SAR) in rats chronically exposed to external electromagnetic fields, as well as the development of a laboratory setup simulating the operation of a fifth-generation 5G New Radio base station (with a signal bandwidth of 15 MHz and a carrier frequency of 2.4 GHz). The paper presents a modified method for theoretical SAR estimation for one-sided irradiation and distributed absorption. Mean whole-body SAR values were estimated by the proposed method and numerically modeled with the CST Microwave Studio simulation software 2020package using primitive rat models. Dielectric parameters in the numerical simulation were used from the software library. The IEEE/IEC 62704-1 algorithm was used to investigate SAR in numerical simulations. The theoretical estimates and numerical simulations were compared for different SAR distributions and were found to be qualitatively comparable. The differences between approximate theoretical estimates and numerical simulations are 7% and 10% for distributed and non-distributed absorptions, respectively. The proposed method, which takes into account the decreasing power flux density, can be used to estimate the approximate whole-body SAR during chronic electromagnetic field exposure in rats.

Open access paper: https://www.mdpi.com/2076-3417/14/1/208

--

The thermal sensation threshold and its reliability induced by the exposure to 28 GHz millimeter-wave

Akiko Y, Shintaro U, Kazuki U, Sachiko K, Norika A, Akimasa H, Yohei O. The thermal sensation threshold and its reliability induced by the exposure to 28 GHz millimeter-wave. Frontiers in Neuroscience. 18. 2024. doi: 10.3389/fnins.2024.1331416.

Abstract

The application of 28 GHz millimeter-wave is prevalent owing to the global spread of fifth-generation wireless communication systems. Its thermal effect is a dominant factor which potentially causes pain and tissue damage to the body parts exposed to the millimeter waves. However, the threshold of this thermal sensation, that is, the degree of change in skin temperature from the baseline at which the first subjective response to the thermal effects of the millimeter waves occurs, remains unclear. Here, we investigated the thermal sensation threshold and assessed its reliability when exposed to millimeter waves. Twenty healthy adults were exposed to 28 GHz millimeter-wave on their left middle fingertip at five levels of antenna input power: 0.2, 1.1, 1.6, 2.1, and 3.4 W (incident power density: 27–399 mW/cm²). This measurement session was repeated twice on the same day to evaluate the threshold reliability. The intraclass correlation coefficient (ICC) and Bland–Altman analysis were used as proxies for the relative and absolute reliability, respectively. The number of participants who perceived a sensation during the two sessions at each exposure level was also counted as the perception rate. Mean thermal sensation thresholds were within 0.9°C–1.0°C for the 126–399 mW/cm² conditions, while that was 0.2°C for the 27 mW/cm² condition. The ICCs for the threshold at 27 and 126 mW/cm² were interpreted as poor and fair, respectively, while those at higher exposure levels were moderate to substantial. Apart from a proportional bias in the 191 mW/cm² condition, there was no fixed bias. All participants perceived a thermal sensation at 399 mW/cm² in both sessions, and the perception rate gradually decreased with lower exposure levels. Importantly, two-thirds of the participants answered that they felt a thermal sensation in both or one of the sessions at 27 mW/cm², despite the low-temperature increase. These results suggest that the thermal sensation threshold is around 1.0°C, consistent across exposure levels, while its reliability increases with higher exposure levels. Furthermore, the perception of thermal sensation may be inherently ambiguous owing to the nature of human perception.

Open access paper:
https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2024.1331416

--

The impact of radiofrequency exposure on Aedes aegypti (Diptera: Culicidae) development [mosquitos]

Nik Abdull Halim NMH, Mohd Jamili AF, Che Dom N, Abd Rahman NH, Jamal Kareem Z, Dapari R. The impact of radiofrequency exposure on Aedes aegypti (Diptera: Culicidae) development. PLoS One. 2024 Feb 27;19(2):e0298738. doi: 10.1371/journal.pone.0298738.

Abstract

Introduction Wireless communication connects billions of people worldwide, relying on radiofrequency electromagnetic fields (RF-EMF). Generally, fifth-generation (5G) networks shift RF carriers to higher frequencies. Although radio, cell phones, and television have benefited humans for decades, higher carrier frequencies can present potential health risks. Insects closely associated with humans (such as mosquitoes) can undergo increased RF absorption and dielectric heating. This process inadvertently impacts the insects' behaviour, morphology, and physiology, which can influence their spread. Therefore, this study examined the impact of RF exposure on Ae. aegypti mosquitoes, which are prevalent in indoor environments with higher RF exposure risk. The morphologies of Ae. aegypti eggs and their developments into Ae. aegypti mosquitoes were investigated.

Methods A total of 30 eggs were exposed to RF radiation at three frequencies: baseline, 900 MHz, and 18 GHz. Each frequency was tested in triplicate. Several parameters were assessed through daily observations in an insectarium, including hatching responses, development times, larval numbers, and pupation periods until the emergence of adult insects.

Results This study revealed that the hatching rate for the 900 MHz group was the highest (79 ± 10.54%) compared to other exposures ($p = 0.87$). The adult emergence rate for the 900 MHz group was also the lowest at 33 ± 2.77%. A significant difference between the groups was demonstrated in the statistical analysis ($p = 0.03$).

Conclusion This work highlighted the morphology sensitivity of Ae. aegypti eggs and their developments in the aquatic phase to RF radiation, potentially altering their life cycle.

This study successfully comprehended the influence of RF exposure on the development of *Aedes* mosquitoes. The 900 MHz RF exposure accelerated the hatching process of *Ae*. *aegypti* mosquitoes and increased the percentage of adult emergence. These findings represented an essential initial stage in understanding the impact of RF radiation on *Aedes* mosquito populations, providing vital insights into the population dynamics. Nevertheless, the potential variability results under distinct study conditions involving RF exposure variation types and the mosquito species studied were necessary. Thus, additional investigation was desirable and crucial to understand the consequences of RF exposure comprehensively on *Ae*. *albopictus* and determine the most efficient approaches for identifying the most effective strategies for dengue vector control. Studies are actively investigating the influence of RF exposure on insects (particularly mosquitoes) due to data suggesting that it can impact hatching and developmental processes. Hence, further studies should be conducted to fully comprehend the scope of these effects and clarify their practical relevance in preventing dengue spread. These ongoing studies are pivotal in learning the potential utility of RF exposure to mitigate the spread of dengue disease and in establishing the most efficacious approaches for translating this knowledge into practical control measures.

Open access paper: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0298738

--

Detrimental impact of cell phone radiation on sperm DNA integrity (Review)

Koohestanidehaghi Y, Khalili MA, Dehghanpour F, Seify M. Detrimental impact of cell phone radiation on sperm DNA integrity. Clin Exp Reprod Med. 2024 Jan 24. doi: 10.5653/cerm.2023.06121.

Abstract

Radiofrequency electromagnetic radiation (RF-EMR) from various sources may impact health due to the generation of frequency bands. Broad pulses emitted within frequency bands can be absorbed by cells, influencing their function. Numerous laboratory studies have demonstrated that mobile phones-generally the most widely used devices-can have harmful effects on sex cells, such as sperm and oocytes, by producing RF-EMR. Moreover, some research has indicated that RF-EMR generated by mobile phones can influence sperm parameters, including motility, morphology, viability, and (most critically) DNA structure. Consequently, RF-EMR can disrupt both sperm function and fertilization. However, other studies have reported that exposure of spermatozoa to RF-EMR does not affect the functional parameters or genetic structure of sperm. These conflicting results likely stem from differences among studies in the duration and exposure distance, as well as the species of animal used. This report was undertaken to review the existing research discussing the effects of RF-EMR on the DNA integrity of mammalian spermatozoa.

EMW can induce oxidative stress, which subsequently leads to disorders such as reduced mobility, morphological changes, acrosome disturbances, and ultimately, damage to the nucleus and genetic material. This oxidative damage to DNA can result in the breakdown of both single-stranded and double-stranded DNA structures, culminating in fragmentation. If the DNA is not repaired and the damage accumulates, the sperm may undergo apoptosis. Damage to the sperm genome can ultimately impact fertility, potentially leading to infertility. Therefore, it is advisable to limit daily exposure to these sources to prevent irreversible damage caused by EMWs. Many men carry their cell phones in their trouser pockets or clipped to their belts, and the use of Bluetooth can increase their susceptibility to RF-EMR exposure. This exposure can induce changes in sperm quality through oxidative stress, potentially leading to infertility. Agarwal et al. [11] suggested that carrying a cell phone in a pocket could lead to a decline in sperm quality. However, it is important to note that the phone and male reproductive organs are separated by multiple tissue layers. Therefore, extrapolating these in vitro effects to real-life conditions requires further studies [11].

In July 2021, the European Parliament commissioned a research report titled “Health impact of 5G.” The report concluded that the commonly used RF-EMFs are likely carcinogenic to humans and have a definitive impact on male fertility. It also suggested potential adverse effects on the development of embryos, fetuses, and newborns. To mitigate these adverse effects, the organization proposed several strategies. These include favoring non-wireless connections, increasing distance from the source of RF-EMFs, switching off devices when not in use, and practicing safe phone usage [55].

Open access paper: https://ecerm.org/journal/view.php?doi=10.5653/cerm.2023.06121

--

Acute exposure of microwave impairs attention process by activating microglial inflammation

Jiang S, Ma Y, Shi Y, *et al.* Acute exposure of microwave impairs attention process by activating microglial inflammation. *Cell Biosci* **14**, 2 (2024). doi: 10.1186/s13578-023-01162-9.

Abstract

Background Attention provides the foundation for cognitions, which was shown to be affected by microwave (MW) radiation. With the ubiquitous of microwaves, public concerns regarding the impact of MW radiation on attention has hence been increased. Our study aims to investigate the potential effect and mechanism of acute microwave exposure on attention.

Results We identified obvious impairment of attention in mice by the five-choice serial reaction time (5-CSRT) task. Proteomic analysis of the cerebrospinal fluid (CSF) revealed neuroinflammation and microglial activation potentially due to acute MW exposure. Moreover, biochemical analysis further confirmed microglial activation in the prefrontal cortex (PFC) of mice subjected to acute MW exposure. Finally, minocycline, a commercially available anti-inflammatory compound, attenuated neuroinflammation, inhibited the upregulation of N-methyl-D-aspartic acid receptor (NMDAR) including NR2A and NR2B, and also accelerated the attentional recovery after MW exposure.

Conclusions We believe that microglial activation and NMDAR upregulation likely contribute to inattention induced by acute MW exposure, and we found that minocycline may be effective in preventing such process

Open access paper: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10768366/

--

Detrimental effects of electromagnetic radiation emitted from cell phone on embryo morphokinetics and blastocyst viability in mice

Seify M, Khalili MA, Anbari F, Koohestanidehaghi Y. Detrimental effects of electromagnetic radiation emitted from cell phone on embryo morphokinetics and blastocyst viability in mice. Zygote. Published online 2024:1-5. doi:10.1017/S0967199424000042

Abstract

Electromagnetic radiation (EMR) has deleterious effects on sperm motility and viability, as well as oocyte membrane and organelle structure. The aim was to assess the effects of cell phone radiation on preimplantation embryo morphokinetics and blastocyst viability in mice. For superovulation, 20 female mice were treated with intraperitoneal (IP) injections of 10 IU pregnant mare's serum gonadotropin (Folligon® PMSG), followed by 10 IU of human chorionic gonadotropin (hCG) after 48 h. The zygotes (n = 150) from the control group were incubated for 4 days. The experimental zygotes (n = 150) were exposed to a cell phone emitting EMR with a frequency range 900–1800 MHz for 30 min on day 1. Then, all embryos were cultured in the time-lapse system and annotated based on time points from the 2-cell stage (t2) to hatched blastocyst (tHDyz), as well as abnormal cleavage patterns. Blastocyst viability was assessed using Hoechst and propidium iodide staining. Significant increases ($P < 0.05$) were observed in the cleavage division time points of t2, t8, t10, and t12 of the experimental group compared with the controls. In terms of blastocyst formation parameters, a delay in embryo development was observed in the experimental group compared with the controls. Data analysis of the time intervals between the two groups showed a significant difference in the s3 time interval ($P < 0.05$). Also, the rates of fragmentation, reverse cleavage, vacuole formation, and embryo arrest were significantly higher in the experimental group ($P < 0.05$). Furthermore, the cell survival rate in the experimental group was lower than the control group ($P < 0.05$). Exposure to EMR has detrimental consequences for preimplantation embryo

development in mice. These effects can manifest as defects in the cleavage stage and impaired blastocyst formation, leading to lower cell viability.

https://www.cambridge.org/core/journals/zygote/article/abs/detrimental-effects-of-electromagnetic-radiation-emitted-from-cell-phone-on-embryo-morphokinetics-and-blastocyst-viability-in-mice/BC3E03A74EC2E5E9C97D6CF6E22253A3

--

Radio-tracking systems emit pulsed waves that could affect the health and alter the orientation of animals

Balmori A. Radio-tracking systems emit pulsed waves that could affect the health and alter the orientation of animals. Journal for Nature Conservation. Vol. 77. 2024, doi: 10.1016/j.jnc.2023.126520.

Abstract

Any study carried out with a given technique must avoid causing harmful effects on the experimental subjects, in order to avoid altering the survival rates of the animals or causing bias in the results of the study. Radio tagging is undoubtedly a very useful technique, and in many cases is the only option available to obtain information on the ecology and movements of wild animals, but unfortunately it has been used indiscriminately, without regard to the problems it can cause. The most well-known of these are the effects from the weight of the device and the harness, but there are others that have not yet been taken into account and that have not been sufficiently studied, such as the non-thermal effects of modulated and/or pulsed electromagnetic fields on the health and orientation of radio-tracked animals. This commentary provides a brief overview of the scientific evidence on the effects of electromagnetic fields, and underlines the importance of carrying out relevant studies in order to check for the possible effects of radio transmitters on animals.

https://www.sciencedirect.com/science/article/pii/S1617138123001917

--

A study on effects of cell phone tower-emitted non-ionizing radiations in an Allium cepa test system

Sharma A, Sharma S, Bahel S, Katnoria JK. A study on effects of cell phone tower-emitted non-ionizing radiations in an Allium cepa test system. Environ Monit Assess. 2024 Feb 13;196(3):261. doi: 10.1007/s10661-024-12435-2.

Abstract

Considering enormous growth in population, technical advancement, and added reliance on electronic devices leading to adverse health effects, *in situ* simulations were made to evaluate effects of non-ionizing radiations emitted from three cell phone towers (T1, T2, and T3) of frequency bands (800, 1800, 2300 MHz), (900, 1800, 2300 MHz), and (1800 MHz), respectively. Five sites (S1–S5) were selected near cell phone towers exhibiting different power densities. The site with zero power density was considered

as control. Effects of radiations were studied on morphology; protein content; antioxidant enzymes like ascorbate peroxidase (APX), superoxide dismutase (SOD), glutathione-S-transferase (GST), guaiacol peroxidase (POD), and glutathione reductase (GR); and genotoxicity using *Allium cepa*. Mean power density ($\mu W/cm^2$) was recorded as 1.05, 1.18, 1.6, 2.73, and 12.9 for sites 1, 2, 3, 4, and 5, respectively. A significant change in morphology, root length, fresh weight, and dry weight in *Allium cepa* was observed under the exposure at different sites. Protein content of roots showed significant difference for samples at all sites while bulbs at sites S4 and S5 when compared to control. Antioxidant activity for root in terms of APX, GST, and POD showed significant changes at S4 and S5 and GR at site S5 and SOD at S1, S2, S3, S4, and S5. Similarly, bulbs showed significant changes at sites S4 and S5 for APX while at sites S3, S4, and S5 for POD and S2, S3, S4, and S5 for SOD and S5 for GR and GST. Genotoxicity study has shown induction of abnormalities at different stages of the cell cycle in *Allium cepa* root tips. The samples under exposure to radiation with maximum power density have shown maximum induction of oxidative stress and genotoxicity.

https://pubmed.ncbi.nlm.nih.gov/38349609/

Excerpts

Specific locations with cell phone towers were identified using the Tarang Sanchar portal. The power density ($\mu W/cm^2$) at different sites near towers was noted manually using an electromagnetic radiation detector (model: PCE-EM 29; make: PCE instruments, UK). Five sites were selected following the criteria of high-power density exposure and site with convenient experimental setup. Cell phone towers emitting radiations of different frequency bands, viz., 800, 1800, and 2300 MHz, for Reliance Jio 900, 1800 and 2300 MHz for Airtel, and 1800 MHz for Vodafone Idea (VI), were located at the sites of experiment. Depending upon the distance from the cell towers, five sites near towers showed the increasing order of their power density with decreasing distance. Site 1 showed a mean power density of 1.05 $\mu W/cm^2$ at a distance of 20 m from cell towers, with an angle of $\angle 16.67°$. Meanwhile, site 2 recorded a mean power density of 1.18 $\mu W/cm^2$ at a distance of 15 m, with an angle of $\angle 21.77°$. Site 3 exhibited a mean power density of 1.6 $\mu W/cm^2$ at a distance of 10 m and an angle of $\angle 30.94°$. Site 4 showed a mean power density of 2.73 $\mu W/cm^2$ at a distance of 5 m and an angle of $\angle 50.19°$. In contrast, site 5 displayed a higher mean power density of 12.9 $\mu W/cm^2$ at distance of 1 m and an angle of $\angle 80.21°$ with an average temperature recorded to be 20–24 °C and relative humidity of 45–50%. The specifics regarding the selection of the exposure sites are provided in Fig. 1. Keeping all other environmental conditions constant and altering only one parameter that is power density, the site with zero power density is designated as the control at a distance of 50 m and an angle of $\angle 6.818°$.

Conclusion

The present study clearly revealed that plant samples under exposure to electromagnetic radiations generated from cell phone towers at varied frequencies and maximum power densities had altered morphological and biochemical characteristics. The enhanced level of antioxidative enzymes like APX, GST, GR, POD, and SOD during the present study indicate the enhancement of oxidative stress and genotoxicity. The harmful effects of exposure to non-ionizing radiations were also witnessed through reduced protein content in test samples as compared to that of control. The genotoxicity studies revealed that the damage by exposure to radiations was high for root samples as compared to bulb samples indicating that root tip cells of *Allium cepa* were sensitive bioindicators to reveal toxicity of radiation pollution. This study is the first of its kind conducted in the field and reveals that further

advance research is the need of an hour for in-depth understanding and devising the effective protective measures for mitigation of the detrimental effects of radiations.

--

A systematic review on cellular responses of Escherichia coli to nonthermal electromagnetic irradiation

Askaripour K, Żak A. A systematic review on cellular responses of Escherichia coli to nonthermal electromagnetic irradiation. Bioelectromagnetics. 2024 Jan;45(1):16-29. doi: 10.1002/bem.22484.

Highlights

- The present review aims to systematically evaluate literature to reach a firm conclusion on the causality between nonthermal electromagnetic radiation of alternating current and biological effects in *Escherichia coli*.
- The systematic review methodology by OHAT and the risk of bias tool evaluating the validity of experimental design and conduct within research works were followed.
- With 25 of the 114 records screened meeting the eligibility criteria, the evidence summation was performed regarding growth rate, morphology, and gene expression under exposure of extremely low frequency, intermediate frequency, or radio frequency.
- With 85% of experiments in the included studies focusing on the extremely low-frequency range, effects on growth rate, morphology, and gene expression were supported in 74%, 80%, and 33% of the experiments, respectively.

Abstract

Investigation of Escherichia coli under electromagnetic fields is of significance in human studies owing to its short doubling time and human-like DNA mechanisms. The present review aims to systematically evaluate the literature to conclude causality between 0 and 300 GHz electromagnetic fields and biological effects in E. coli. To that end, the OHAT methodology and risk of bias tool were employed. Exponentially growing cells exposed for over 30 min at temperatures up to 37◦C with fluctuations below 1◦C were included from the Web-of-Knowledge, PubMed, or EMF-Portal databases. Out of 904 records identified, 25 articles satisfied the selection criteria, with four excluded during internal validation. These articles examined cell growth (11 studies), morphology (three studies), and gene regulation (11 studies). Most experiments (85%) in the included studies focused on the extremely low-frequency (ELF) range, with 60% specifically at 50 Hz. Changes in growth rate were observed in 74% of ELF experiments and 71% of radio frequency (RF) experiments. Additionally, 80% of ELF experiments showed morphology changes, while gene expression changes were seen in 33% (ELF) and 50% (RF) experiments. Due to the limited number of studies, especially in the intermediate frequency and RF ranges, establishing correlations between EMF exposure and biological effects on E. coli is not possible.

https://pubmed.ncbi.nlm.nih.gov/37807247/

Conclusions

The present review systemically evaluates whether the current literature, passing through the risk of bias tool, while fulfilling the selection criteria defined in the PECO statement, supports an association between nonthermal electromagnetic fields and alterations in the biological composition and functioning of *E. coli* strains. The findings from the 21 included studies are dependent on the specific endpoints assessed, which include cell growth, morphology, and gene expression. The conclusions are subject to certain limitations, as the review focuses on exponentially growing strains that were exposed to nonthermal EMFs of frequencies 0–300 GHz for a minimum duration of 30 min.

In each study, it is possible that multiple experiments were conducted. Among these studies, 85% of experiments (16 articles) focused on the extremely low-frequency range, with 60% specifically examining 50 Hz. Additionally, 5% of experiments (one article) addressed the IF range, while 10% of experiments (four articles) addressed the RF range. In the investigation of nonthermal effects of extremely low-frequency electromagnetic fields (ELF-EMF), 74% of the experiments supported the effects on growth rate (investigated in nine articles), 80% supported the effects on morphology (investigated in three articles), and 33% supported the effects on gene expression (investigated in eight articles). Within the IF range, nonthermal effects on growth rate were observed in 100% of experiments (conducted in one article), but no experiments supported such effects on gene expression (investigated in one article). In the RF range, 71% of experiments supported nonthermal effects on growth rate (conducted in two articles), while 50% supported nonthermal effects on gene expression (conducted in two articles). Therefore, due to the limited number of studies available on the IF and RF ranges, the reliability of the results may primarily apply to ELF-EMF, particularly at 50 Hz.

When experiments demonstrated biological effects, the observed pattern varied as either beneficial or detrimental, depending on the exposure conditions and experimental setup. For example, out of 38 experiments on growth rate, 12 showed beneficial effects while 16 showed detrimental effects. This controversy can be explained by the concept of the *biological window*, in which EMF may contribute to positive physiological processes. However, if the accumulated exposure exceeds the cell-specific energy limit, detrimental effects may prevail. Several sources exhibited inconsistencies when observing these effects.

In terms of exposure parameters, continuity, referring to intermittent or continuous status, had a significant effect on the results, with pulsed signals showing a greater impact compared to sinusoidal signals. Extremely low frequencies and radio frequencies had a major impact on the investigated endpoints, while intermediate frequencies have received limited research attention. The intensity and duration of exposure influenced the absorbed energy levels, resulting in varying effects ranging from beneficial to neutral or detrimental. Modulation may introduce significant differences compared to unmodulated signals, depending on the modulation frequency. The optimal measurement time point depends on the specific endpoint and necessitates mechanistic understanding for accurate identification.

Special attention is needed regarding sufficient and clear reports on the amount of electromagnetic irradiation absorbed by the strains, as it can make the quantitative comparison between studies possible, and discriminate between nondetrimental or detrimental effects. Besides, special attention to the background electromagnetic irradiation as a confounder and the measurement time point for allowing the outcome to develop is required. Deficiencies associated with the risk of bias including *research personnel blinded to the study group*, *randomization of exposure*, *allocation concealment*, *attrition/exclusion bias*, and their possible effect on the outcome should be considered. More mechanistic investigations regarding cellular processes and their related genes are demanded.

Based on the limited number of studies encompassing three different frequency ranges and three different endpoints, the systematic review does not establish a conclusive correlation between nonthermal exposure to electromagnetic fields and cellular alterations in *E. coli* strains. However, future research could explore this correlation more specifically, considering the similarity of endpoints and levels of exposure between *E. coli* and humans. This exploration would help shed light on the potential implications of these effects on human health.

--

On the Quasistationarity of the Ambient Electromagnetic Field Generated by Wi-Fi Sources

Tuță L, Roșu G, Andone A, Spandole-Dinu S, Fichte LO. On the Quasistationarity of the Ambient Electromagnetic Field Generated by Wi-Fi Sources. *Electronics*. 2024; 13(2):301. doi:10.3390/electronics13020301.

Abstract

In recent decades, the widespread use of mobile phones and wireless technologies has led to a significant increase in radiofrequency electromagnetic fields (RF-EMFs), raising concerns about continuous RF-EMF exposure among the general population. Recent research indicates that real-life RF signals are more biologically active than controlled laboratory signals with a low variability, suggesting that living organisms can adapt to EMF exposure when the pattern has a low variability. However, using real-life sources with unpredictable variation signals in biological experiments contradicts the principle of experiment controllability. This paper aims to investigate the nature of signals generated by current sources of ambient EMFs in terms of stationarity, with the goal of replicating them in biological experiments to study the effects of EMF exposure. Employing a range of statistical methodologies, starting with descriptive statistical analysis and progressing to the advanced APDP and APTF methods, an examination is conducted on a collection of Wi-Fi signal recordings across various operating modes, with particular attention given to video streaming. The chosen datasets are scrutinized with respect to their adherence to a Gaussian distribution and the concept of stationarity. The results indicate that the observed Wi-Fi signals lack stationarity in both the time and frequency domains. However, based on the analytical findings, it is possible to generate signals in frequency that authentically replicate Wi-Fi signals, accounting for nonstationarity considerations.

Conclusions

The stationarity of electromagnetic emission sources, particularly those generating time-varying EMFs, can significantly impact living organisms. Exposure to nonstationary sources with dynamic signals limits an organism's ability to adapt to the changing stimulus. The consideration that real-life source signals may be more biologically active raises concerns about the accuracy and relevance of laboratory sources in experiments studying the effects of RF-EMF exposure. However, using real-life sources with unpredictable variations contradicts the principle of experiment controllability. The investigation into the impact of Wi-Fi routers on living organisms is crucial, as these emission sources exhibit either weak stationarity or nonstationarity, potentially leading to adverse effects on human health.

The paper describes the examination of temporal variations in ambient electromagnetic exposure emanating from Wi-Fi communication devices which involves the application of diverse statistical

methodologies. The objective is to ascertain the stationarity of these signals. The ultimate outcome aims to produce laboratory-controlled signals that faithfully replicate the authentic variability observed in real-life signals generated by Wi-Fi communication devices.

Several datasets were obtained from various frequency-domain measurements with a spectrum analyzer. A stationarity analysis was performed for the measured datasets, in both time and frequency domains, using the ADPD and APTF methods.

The findings derived from the analysis using APTF and APDP indicate that the observed Wi-Fi signals lack stationarity in both the time and frequency domains. This outcome poses a challenge with respect to simulating Wi-Fi signals, suggesting that the incorporation of modulations exclusive to the 802.11 standards (OFDM and DSSS), along with the introduction of Gaussian noise, may prove insufficient for an accurate emulation of authentic Wi-Fi signals.

We suggest, in this case, simulating the spectrum analyzer measurement by generating vectors from the Pearson system of distributions, with statistical parameters that correspond to practical measurements. The quality of these simulated Wi-Fi waveforms will depend on the measurements used to train the algorithm. These waveforms could be used for further EMF investigation by generating them using a portable SDR device, without requiring a Wi-Fi router with Internet access.

Open access paper: https://www.mdpi.com/2079-9292/13/2/301

--

Changes in honey bee nutrition after exposure to radiofrequency electromagnetic field

Migdal P, Plotnik M, Bieńkowski P, Murawska A, Berbeć E, Sobkiewicz P, Zarębski K, Latarowski K (2024). Changes in honey bee nutrition after exposure to radiofrequency electromagnetic field. The European Zoological Journal, 91:1, 172-179. doi: 10.1080/24750263.2024.2308550.

Abstract

Urban beekeeping has gained interest in recent years. Bees placed in cities, on the roofs of buildings, are exposed to many different sources of electromagnetic fields of different frequencies and intensities. Knowledge about the impact of electromagnetic fields on the physiology and behavior of insects (including honey bees) is limited. Hence, one of the first steps was to check how radiofrequency electromagnetic fields affect honey bees' nutrition. The level of total proteins, glucose, triglycerides, and TAS activity in bee hemolymph was analyzed. This is because they indicate the honey bee's nutrition level. The experiment involved 2-day-old Carniolan worker honey bees (*Apis mellifera carnica* L.). The bees were exposed to fields of 12 V/m, 28 V/m, and 61 V/m for 0.25, 1 and 3 h. Hemolymph was taken from 40 alive worker honey bees from each group immediately after exposure, by removing the antennae with sterile tweezers. The activity of biochemical markers (total proteins, TAS, and triglycerides) in the bee hemolymph in 12 V/m and 28 V/m groups showed statistically significant differences compared to the control group in most cases. According to our own research, radiofrequency electromagnetic fields disturb honey bee nutrition. The changes in the analyzed indicators may have long-term effects.

Open access paper: https://www.tandfonline.com/doi/full/10.1080/24750263.2024.2308550

--

Studies of the Electromagnetic Background at the Antarctic Vostok Station

Shurshakov VA, Belov EV, Ilyin EA, *et al.* Studies of the Electromagnetic Background at the Antarctic Vostok Station. *Hum Physiol* **49**, 906–909 (2023). doi: 10.1134/S0362119723070216.

Abstract

The electromagnetic background was measured in the living quarters of the Vostok Station located on the Antarctic plateau distant from the potential sources of technogenic electromagnetic fields. The level of E-radiation in the range from 0.8 to 8 GHz was determined using a MERA integral dosimeter. Based on the results of long-term continuous (one to 4 months) measurements, the mean flux density was equal to 1.5 ± 0.8 nWt/cm, which is nearly 7000 times lower than the admissible level for population adopted in Russia. The Antarctic Vostok Station can be considered as a plausible location for biomedical investigations into the effects of low-frequency electromagnetic radiation.

Conclusions

(1) The EFD values of technogenic EMR in the living quarters of the Vostok Station are 7 thousand times below the regulatory standards for EMR in the Russian Federation. This fact indicates that the long-term stay of people at the Vostok Station is safe as regards the sanitary standards for EMR. [The Russian regulatory standards for EMR 0.3–300 GHz are 1000 μW/cm^2 for personnel and 10 μW/cm^2 for the population.]

(2) The daily dynamics of the main frequencies in EMR EFD variability and its dependence on solar activity have been revealed.

(3) Since the members of polar expeditions at the Vostok Station perform daily outdoor work (up to 1.5 h), it would be reasonable for future expeditions to measure the EFD of EMR in an open area, which will allow estimating the natural (background) level of EMR EFD in the vicinity of the station.

The Antarctic Vostok Station can be considered as one of the possible places for conducting biomedical research under the conditions of reduced electromagnetic radiation flux.

Open access paper: https://link.springer.com/article/10.1134/S0362119723070216

--

Shortwave radiation-induced reproductive organ damage in male rats by enhanced expression of molecules associated with the calpain/Cdk5 pathway and oxidative stress

Yao B, Men J, Liu S, Bai Y, Yu C, Gao Y, Xu X, Zhao L, Zhang J, Wang H, Li Y, Peng R. Shortwave radiation-induced reproductive organ damage in male rats by enhanced expression of molecules associated with the calpain/Cdk5 pathway and oxidative stress. Electromagn Biol Med. 2023 Oct 2;42(4):150-162. doi: 10.1080/15368378.2023.2296896.

Abstract

Shortwave radiation has been reported to have harmful effects on several organs in humans and animals. However, the biological effects of 27 MHz shortwave on the reproductive system are not clear. In this study, we investigated the effects of shortwave whole-body exposure at a frequency of 27 MHz on structural and functional changes in the testis. Male Wistar rats were exposed to **27 MHz continuous shortwaves at average power densities of 0, 5, 10, or 30 mW/cm^2 for 6 min**. The levels of insulin-like factor 3 (INSL3) and anti-sperm antibodies (AsAb) in the peripheral serum, sperm motility, sperm malformation rate, and testicular tissue structure of rats were analyzed. Furthermore, the activity of superoxide dismutase (SOD), catalase (CAT), malondialdehyde (MDA) content, calpain, and Cdk5 expression were analyzed at 1, 7, 14, and 28 days after exposure. We observed that the rats after radiation had decreased serum INSL3 levels ($p < 0.01$), increased AsAb levels ($p < 0.05$), decreased percentage of class A+B sperm ($p < 0.01$ or $p < 0.05$), increased sperm malformation ($p < 0.01$ or $p < 0.05$), injured testicular tissue structure, decreased SOD and CAT activities ($p < 0.01$ or $p < 0.05$), increased MDA content ($p < 0.01$), and testicular tissue expressions of calpain1, calpain2, and Cdk5 were increased ($p < 0.01$ or $p < 0.05$). In conclusion, Shortwave radiation caused functional and structural damage to the reproductive organs of male rats. Furthermore, oxidative stress and key molecules in the calpain/Cdk5 pathway are likely involved in this process.

Plain language summary

Shortwave radiation has been used in communications, medical and military applications, and its damaging effects on several organs of the human body have been reported in the literature. However, the biological effects of shortwave radiation on the male reproductive system are unknown. The present study, by constructing an animal model of short-wave radiation and analyzing the experimental results, revealed that shortwave radiation could cause functional and structural damage to the reproductive organs of male rats, and that oxidative stress and key molecules in the calpain/Cdk5 pathway might be involved in this process. It will provide organizational data for further studies on the mechanisms of male reproductive damage by shortwave radiation.

https://pubmed.ncbi.nlm.nih.gov/38155529/

--

Effects of 3.5-GHz radiofrequency radiation on energy-regulatory hormone levels in the blood and adipose tissue

Bektas H, Dasdag S, Altindag F, Akdag MZ, Yegin K, Algul S. Effects of 3.5-GHz radiofrequency radiation on energy-regulatory hormone levels in the blood and adipose tissue. Bioelectromagnetics. 2024 Feb 18. doi: 10.1002/bem.22498.

Abstract

In recent years exposure of living beings to radiofrequency radiation (RFR) emitted from wireless equipment has increased. In this study, we investigated the effects of 3.5-GHz RFR on hormones that regulate energy metabolism in the body. Twenty-eight rats were divided into four groups: healthy sham (n = 7), healthy RFR (n = 7), diabetic sham (n = 7), and diabetic RFR (n = 7). Over a month, each group

spent 2 h/day in a Plexiglas carousel. The rats in the experimental group were exposed to RFR, but the sham groups were not. At the end of the experiment, blood and adipose tissues were collected from euthanized rats. Total antioxidant, total oxidant, hydrogen peroxide, ghrelin, nesfatin-1, and irisin were determined. Insulin expression in pancreatic tissues was examined by immunohistochemical analysis. Whole body specific absorption rate was 37 mW/kg. For the parameters analyzed in blood and fat, the estimated effect size varied within the ranges of 0.215-0.929 and 0.503-0.839, respectively. The blood and adipose nesfatin-1 ($p = 0.002$), blood and pancreatic insulin are decreased, ($p = 0.001$), gherelin ($p = 0.020$), irisin ($p = 0.020$), and blood glucose ($p = 0.040$) are increased in healthy and diabetic rats exposed to RFR. While nesfatin-1 are negatively correlated with oxidative stress, hyperglycemia and insulin, ghrelin and irisin are positively correlated with oxidative stress and hyperglycemia. Thus, RFR may have deleterious effects on energy metabolism, particularly in the presence of diabetes.

https://pubmed.ncbi.nlm.nih.gov/38369591/

--

Evidence of spatial clustering of childhood acute lymphoblastic leukemia cases in Greater Mexico City: report from the Mexican Inter-Institutional Group for the identification of the causes of childhood leukemia

Duarte-Rodríguez DA, Flores-Lujano J, McNally RJQ, Pérez-Saldivar ML, Jiménez-Hernández E, Martín-Trejo JA, Espinoza-Hernández LE, Medina-Sanson A, Paredes-Aguilera R, Merino-Pasaye LE, Velázquez-Aviña MM, Torres-Nava JR, Espinosa-Elizondo RM, Amador-Sánchez R, Dosta-Herrera JJ, Mondragón-García JA, González-Ulibarri JE, Martínez-Silva SI, Espinoza-Anrubio G, Paz-Bribiesca MM, Salcedo-Lozada P, Landa-García RÁ, Ramírez-Colorado R, Hernández-Mora L, Santamaría-Ascencio M, López-Loyola A, Godoy-Esquivel AH, García-López LR, Anguiano-Ávalos AI, Mora-Rico K, Castañeda-Echevarría A, Rodríguez-Jiménez R, Cibrian-Cruz JA, Solís-Labastida KA, Cárdenas-Cardos R, López-Santiago N, Flores-Villegas LV, Peñaloza-González JG, González-Ávila AI, Sánchez-Ruiz M, Rivera-Luna R, Rodríguez-Villalobos LR, Hernández-Pérez F, Olvera-Durán JÁ, García-Cortés LR, Mata-Rocha M, Sepúlveda-Robles OA, Bekker-Méndez VC, Jiménez-Morales S, Meléndez-Zajgla J, Rosas-Vargas H, Vega E, Núñez-Enríquez JC, Mejía-Aranguré JM. Evidence of spatial clustering of childhood acute lymphoblastic leukemia cases in Greater Mexico City: report from the Mexican Inter-Institutional Group for the identification of the causes of childhood leukemia. Front Oncol. 2024 Feb 14;14:1304633. doi: 10.3389/fonc.2024.1304633

Abstract

Background: A heterogeneous geographic distribution of childhood acute lymphoblastic leukemia (ALL) cases has been described, possibly, related to the presence of different environmental factors. The aim of the present study was to explore the geographical distribution of childhood ALL cases in Greater Mexico City (GMC).

Methods: A population-based case-control study was conducted. Children <18 years old, newly diagnosed with ALL and residents of GMC were included. Controls were patients without leukemia recruited from second-level public hospitals, frequency-matched by sex, age, and health institution with the cases. The residence address where the patients lived during the last year before diagnosis (cases) or the interview (controls) was used for geolocation. Kulldorff's spatial scan statistic was used to detect spatial clusters (SCs). Relative risks (RR), associated p-value and number of cases included for each cluster were obtained.

Results: A total of 1054 cases with ALL were analyzed. Of these, 408 (38.7%) were distributed across eight SCs detected. A relative risk of 1.61 ($p<0.0001$) was observed for the main cluster. Similar results were noted for the remaining seven ones. Additionally, a proximity between SCs, electrical installations and petrochemical facilities was observed.

Conclusions: The identification of SCs in certain regions of GMC suggest the possible role of environmental factors in the etiology of childhood ALL.

Open access paper:
https://www.frontiersin.org/journals/oncology/articles/10.3389/fonc.2024.1304633/full

--

International study of childhood leukemia in residences near electrical transformer rooms

Crespi CM, Sudan M, Juutilainen J, Roivainen P, Hareuveny R, Huss A, Kandel S, Karim-Kos HE, Thuróczy G, Jakab Z, Spycher BD, Flueckiger B, Vermeulen R, Vergara X, Kheifets L. International study of childhood leukemia in residences near electrical transformer rooms. Environ Res. 2024 Feb 10:118459. doi: 10.1016/j.envres.2024.118459.

Abstract

Objectives: New epidemiologic approaches are needed to reduce the scientific uncertainty surrounding the association between extremely low frequency magnetic fields (ELF-MF) and childhood leukemia. While most previous studies focused on power lines, the Transformer Exposure study sought to assess this association using a multi-country study of children who had lived in buildings with built-in electrical transformers. ELF-MF in apartments above built-in transformers can be 5 times higher than in other apartments in the same building. This novel study design aimed to maximize the inclusion of highly exposed children while minimising the potential for selection bias.

Methods: We assessed associations between residential proximity to transformers and risk of childhood leukemia using registry based matched case-control data collected in five countries. Exposure was based on the location of the subject's apartment relative to the transformer, coded as high (above or adjacent to transformer), intermediate (same floor as apartments in high category), or unexposed (other apartments). Relative risk (RR) for childhood leukemia was estimated using conditional logistic and mixed logistic regression with a random effect for case-control set.

Results: Data pooling across countries yielded 16 intermediate and 3 highly exposed cases. RRs were 1.0 (95% CI: 0.5, 1.9) for intermediate and 1.1 (95% CI: 0.3, 3.8) for high exposure in the conditional logistic model. In the mixed logistic model, RRs were 1.4 (95% CI: 0.8, 2.5) for intermediate and 1.3 (95% CI: 0.4, 4.4) for high. Data of the most influential country showed RRs of 1.1 (95% CI: 0.5, 2.4) and 1.7 (95% CI: 0.4, 7.2) for intermediate (8 cases) and high (2 cases) exposure.

Discussion: Overall, evidence for an elevated risk was weak. However, small numbers and wide confidence intervals preclude strong conclusions and a risk of the magnitude observed in power line studies cannot be excluded.

https://pubmed.ncbi.nlm.nih.gov/38346482/

--

The role of curcumin during pregnancy on the exposed fetuses' tissues of Wistar rats to electromagnetic field

Kalantar MH, Bayat PD, Ghaffari Khaligh S, Soleimani H. The role of curcumin during pregnancy on the exposed fetuses' tissues of Wistar rats to electromagnetic field. Electromagn Biol Med. 2024 Feb 17:1-10. doi: 10.1080/15368378.2024.2315214.

Abstract

To investigate curcumin (CUR) as the protector against the harmful effects of low-frequency electromagnetic field(LF- EMF, 50 Hz) during pregnancy period, 5 males and 15 females of Wistar rat mated and vaginal plaques were observed. Then, the pregnant rats were divided into six groups. During pregnancy(21 days), the EMF group was exposed to EMF for 30 min/day, the CUR group received a single dose of 50 mg/kg/daily CUR intraperitoneal, the EMF+CUR group was injected CUR and exposed to EMF daily. The DMSO(dimethyl sulfoxide) group was injected solvent of CUR (DMSO) intraperitoneal with the same volume of CUR solvent, the sham group was placed through the solenoid in the same conditions as the first group without exposure and the control group was kept in their cage in normal condition. After four weeks, babies born were divided according to the mother groups and sacrificed. Then, the three tissues injuries were investigated. EMF exposure led to an increase in outstanding necrotic areas in hippocampal tissue, an increase in the amount of hyperemia(p = 0.017) and necrotic(p = 0.005) in kidneys, and degeneration in liver tissue(p = 0.007) in the EMF group compared with EMF+CUR groups. A single dose of CUR daily during pregnancy can protect these tissues from injuries caused by LF-EMF exposure in rat fetuses.

Plain language summary

Electromagnetic fields (EMFs) are able to penetrate and be absorbed by the body. The researchers showed that these radiations might be harmful and lead to cancers, cardiovascular diseases, mental disorders, and fetal abnormalities. Curcumin as an active component in turmeric has anti-inflammatory, antioxidant and anti-hyperlipidemia properties. It can protect the body against diseases such as arthritis, anxiety, and metabolic syndrome. This study examined the effects of curcumin as the protector against the harmful effects of EMF (50Hz) during pregnancy period. So the pregnant rats were divided into six groups. During pregnancy, a group was exposed to EMF for 30 min/day, the second group was injected a dose of curcumin 50mg/kg/daily, the third group was injected curcumin and exposed to EMF daily. The fourth group was injected a curcumin solvent dose, the sham group was placed through the field generator in the same conditions as the first group without exposure and the control group was kept in their cage in normal condition. After four weeks, babies born were divided according to the mother groups and sacrificed. Then, the liver, kidney, and hippocampal tissues were investigated. EMF exposure led to an outstanding increase in necrotic areas in hippocampal tissue, a notable increase in the amount of hyperemia and necrosis in kidneys, and degeneration in liver tissue(p=0.007) in the EMF group compared with the third group that was exposed to EMF and received curcumin. A single dose of curcumin daily during pregnancy can protect these tissues from injuries caused by EMF(50Hz) exposure in rat fetuses.

https://pubmed.ncbi.nlm.nih.gov/38366892/

--

Extremely low-frequency EMF facilitate both osteoblast and osteoclast activity through Wnt/β-catenin signaling in the zebrafish scale

Kobayashi-Sun J, Kobayashi I, Kashima M, Hirayama J, Kakikawa M, Yamada S, Suzuki N. Extremely low-frequency electromagnetic fields facilitate both osteoblast and osteoclast activity through Wnt/β-catenin signaling in the zebrafish scale. Front Cell Dev Biol. 2024 Feb 7;12:1340089. doi: 10.3389/fcell.2024.1340089.

Abstract

Electromagnetic fields (EMFs) have received widespread attention as effective, noninvasive, and safe therapies across a range of clinical applications for bone disorders. However, due to the various frequencies of devices, their effects on tissues/cells are vary, which has been a bottleneck in understanding the effects of EMFs on bone tissue. Here, we developed an *in vivo* model system using zebrafish scales to investigate the effects of extremely low-frequency EMFs (ELF-EMFs) on fracture healing. Exposure to 10 millitesla (mT) of ELF-EMFs at 60 Hz increased the number of both osteoblasts and osteoclasts in the fractured scale, whereas 3 or 30 mT did not. Gene expression analysis revealed that exposure to 10 mT ELF-EMFs upregulated *wnt10b* and Wnt target genes in the fractured scale. Moreover, *β*-catenin expression was enhanced by ELF-EMFs predominantly at the fracture site of the zebrafish scale. Inhibition of Wnt/*β*-catenin signaling by IWR-1-endo treatment reduced both osteoblasts and osteoclasts in the fractured scale exposed to ELF-EMFs. These results suggest that ELF-EMFs promote both osteoblast and osteoclast activity through activation of Wnt/*β*-catenin signaling in fracture healing. Our data provide *in vivo* evidence that ELF-EMFs generated with a widely used commercial AC power supply have a facilitative effect on fracture healing.

Open access paper: https://www.frontiersin.org/articles/10.3389/fcell.2024.1340089/full

--

Influence of Magnetic Fields, Including the Planetary Magnetic Field, on Complex Life Forms: How Do Biological Systems Function in This Field and in Electromagnetic Fields?

Hart DA. The Influence of Magnetic Fields, Including the Planetary Magnetic Field, on Complex Life Forms: How Do Biological Systems Function in This Field and in Electromagnetic Fields? *Biophysica*. 2024; 4(1):1-21. doi: 10.3390/biophysica4010001.

Abstract
Life on Earth evolved to accommodate the biochemical and biophysical boundary conditions of the planet millions of years ago. The former includes nutrients, water, and the ability to synthesize other needed chemicals. The latter includes the 1 g gravity of the planet, radiation, and the geomagnetic field (GMF) of the planet. How complex life forms have accommodated the GMF is not known in detail, considering that Homo sapiens evolved a neurological system, a neuromuscular system, and a cardiovascular system that developed electromagnetic fields as part of their functioning. Therefore, all

of these could be impacted by magnetic fields. In addition, many proteins and physiologic processes utilize iron ions, which exhibit magnetic properties. Thus, complex organisms, such as humans, generate magnetic fields, contain significant quantities of iron ions, and respond to exogenous static and electromagnetic fields. Given the current body of literature, it remains somewhat unclear if Homo sapiens use exogenous magnetic fields to regulate function and what can happen if the boundary condition of the GMF no longer exerts an effect. Proposed deep space flights to destinations such as Mars will provide some insights, as space flight could not have been anticipated by evolution. The results of such space flight "experiments" will provide new insights into the role of magnetic fields on human functioning. This review will discuss the literature regarding the involvement of magnetic fields in various normal and disturbed processes in humans while on Earth and then further discuss potential outcomes when the GMF is no longer present to impact host systems, as well as the limitations in the current knowledge. The GMF has been present throughout evolution, but many details of its role in human functioning remain to be elucidated, and how humans have adapted to such fields in order to develop and retain function remains to be elucidated. Why this understudied area has not received the attention required to elucidate the critical information remains a conundrum for both health professionals and those embarking on space flight. However, proposed deep space flights to destinations such as Mars may provide the environments to test and assess the potential roles of magnetic fields in human functioning.

Open access paper: https://www.mdpi.com/2673-4125/4/1/1

--

Do electromagnetic fields from subsea power cables effect benthic elasmobranch behaviour? A risk-based approach for the Dutch Continental Shelf

Hermans A, Winter HV, Gill AB, Murk AJ. Do electromagnetic fields from subsea power cables effect benthic elasmobranch behaviour? A risk-based approach for the Dutch Continental Shelf. Environ Pollut. 2024 Feb 13:123570. doi: 10.1016/j.envpol.2024.123570.

Abstract

Subsea power cables cause electromagnetic fields (EMFs) into the marine environment. Elasmobranchs (rays, skates, sharks) are particularly sensitive to EMFs as they use electromagnetic-receptive sensory systems for orientation, navigation, and locating conspecifics or buried prey. Cables may intersect with egg laying sites, mating, pupping, and nursery grounds, foraging habitat and migration routes of elasmobranchs and the effects of encountering EMFs on species of elasmobranchs are largely unknown. Demonstrated behavioural effects are attraction, disturbance and indifference, depending on EMF characteristics, exposed life stage, exposure level and duration. We estimated exposure levels of elasmobranchs to subsea power cable EMFs, based on modelled magnetic fields in the Dutch Continental Shelf and compared these to reported elasmobranch sensory sensitivity ranges and experimental effect levels. We conclude that the risk from subsea power cables has a large uncertainty and varies per life stage and species ecology. Based on estimated no-observed effect levels (from 10^{-3} to 10^{-1} μT) we discuss what will probably be the most affected species and life stage for six common benthic elasmobranchs in the Southern North Sea. We then identify critical knowledge gaps for reducing the uncertainty in the risk assessments for EMFs effects on benthic elasmobranchs.

https://pubmed.ncbi.nlm.nih.gov/38360387/

--

A Boundary Element Method of Bidomain Modeling for Predicting Cellular Responses to Electromagnetic Fields

Czerwonky DM, Aberra AS, Gomez LJ. A Boundary Element Method of Bidomain Modeling for Predicting Cellular Responses to Electromagnetic Fields. bioRxiv [Preprint]. 2023 Dec 16:2023.12.15.571917. doi: 10.1101/2023.12.15.571917. (preprint, not peer-reviewed)

Abstract
Objective: Commonly used cable equation-based approaches for determining the effects of electromagnetic fields on excitable cells make several simplifying assumptions that could limit their predictive power. Bidomain or "whole" finite element methods have been developed to fully couple cells and electric fields for more realistic neuron modeling. Here, we introduce a novel bidomain integral equation designed for determining the full electromagnetic coupling between stimulation devices and the intracellular, membrane, and extracellular regions of neurons.

Methods: Our proposed boundary element formulation offers a solution to an integral equation that connects the device, tissue inhomogeneity, and cell membrane-induced E-fields. We solve this integral equation using first-order nodal elements and an unconditionally stable Crank-Nicholson time-stepping scheme. To validate and demonstrate our approach, we simulated cylindrical Hodgkin-Huxley axons and spherical cells in multiple brain stimulation scenarios.

Main results: Comparison studies show that a boundary element approach produces accurate results for both electric and magnetic stimulation. Unlike bidomain finite element methods, the bidomain boundary element method does not require volume meshes containing features at multiple scales. As a result, modeling cells, or tightly packed populations of cells, with microscale features embedded in a macroscale head model, is made computationally tractable, and the relative placement of devices and cells can be varied without the need to generate a new mesh.

Significance: Device-induced electromagnetic fields are commonly used to modulate brain activity for research and therapeutic applications. Bidomain solvers allow for the full incorporation of realistic cell geometries, device E-fields, and neuron populations. Thus, multi-cell studies of advanced neuronal mechanisms would greatly benefit from the development of fast-bidomain solvers to ensure scalability and the practical execution of neural network simulations with realistic neuron morphologies.

Open access paper: https://www.biorxiv.org/content/10.1101/2023.12.15.571917v1

--

Static magnetic fields as a factor in modification of tissue and cell structure: a review

Saletnik BA, Puchalska-Sarna A, Saletnik A, Lipa T, Dobrzański B, Puchalski C. Static magnetic fields as a factor in modification of tissue and cell structure: a review. *Int. Agrophys.* 2024;38(1):43-75. doi:10.31545/intagr/176998.

Highlights

- The static magnetic field (SMF) is an indispensable factor in the natural environment.
- A moderate SMF in the most commonly used range of 2–80 mT has a potential application in the formation and re-modeling of plants and animals, including human cells.
- SMFs can significantly change the potential of the cell membrane, and thus can have a significant impact on the properties of the cell.

Abstract

This review is intended to contribute to the evidence of the effects of static magnetic field on cells and tissue, as well as to present research results that will elucidate the complex matters involved in the formation and remodeling of cells. The cell characteristics studied in the papers that are reviewed include cell viability and proliferation, aggregation and their differentiation, structure and membrane potential. A moderate static magnetic field in the most commonly used range of 2-80 mT has potential application in the formation and remodeling of plant and human cells. However, in the case of cancer cells, the range of fields commonly used was 0.2-9 T. Magnetism promotes changes in plant cell growth, which prompts the cell to proliferate, thereby ensuring an increased rate of biomass production. Some research presented the enhancement of the differentiation of plant cells and skeletal muscle tissue by over 30% at 80 mT static magnetic field. Changes in the cell cycle and growth reflect directly on the cell number and viability and provide useful information to detect modifications in the cell machinery. Static magnetic field, depending on its intensity, enhances cell proliferation and thus may improve, among other processes, tissue regeneration, wound healing and the inhibition of cancer cell proliferation. Researchers showed, among other things, that cells under the influence of static magnetic field changed their shape, had a larger chloroplast, stiffer cell wall, density of the cytoskeleton and cytoplasm contained several mitochondria. Numerous studies also discussed the behavior of the cell membrane of plant and animal organisms, including humans, under the influence of a static magnetic field. The effects of static magnetic field on the cell membrane of plant and human cells were similar. The research results indicate that static magnetic fields can significantly change membrane depolarization and its potential that regulates ion movement and thus can have a significant impact on the properties and biological functionality of the cell. Studies have shown that continuous application of static magnetic field caused deformation and damage of cell membrane. Based on the theoretical analyses presented also in this review, it can be concluded that static magnetic field affects cells and tissue, giving them changes in properties and behaviors and modulates, e.g. in the activity of ion channels. Thus it may produce effects leading to changes in the functioning of the cell. It is possible to formulate directions for further research aimed at using static magnetic fields for the non-invasive remodeling and formation of plant and human cells.

Open access paper: http://www.international-agrophysics.org/Static-magnetic-fields-as-a-factor-in-modification-of-tissue-and-cell-structure-a,176998,0,2.html

--

The biological effects of terahertz wave radiation-induced injury on neural stem cells

Wang Y, Xiong Y, Chen M, Liu F, He H, Ma Q, Gao P, Xiang G, Zhang L. The biological effects of terahertz wave radiation-induced injury on neural stem cells. iScience. 2023 Jul 18;26(10):107418. doi: 10.1016/j.isci.2023.107418.

Abstract

Terahertz (THz) is an electromagnetic wave with a radiation wavelength range of 30-3000 μm and a frequency of 0.1-10 THz. With the development of new THz sources and devices, THz has been widely applied in various fields. However, there are few studies on biological effects of THz irradiation on the human neural stem cells (hNSCs) and mouse neural stem cells (mNSCs), which need to be further studied. We studied the biological effects of THz radiation on hNSCs and mNSCs. The effects of THz irradiation time and average output power on the proliferation, apoptosis, and DNA damage of NSCs were analyzed by flow cytometry and immunofluorescence. The results showed that the proliferation and apoptosis of NSCs were dose-dependently affected by THz irradiation time and average output power. The proliferation of hNSCs was more vulnerable to damage and apoptosis was more serious under the same terahertz irradiation conditions compared to those of mNSCs.

Open access paper: https://www.cell.com/iscience/fulltext/S2589-0042(23)01495-5